



AHCCCS Fee-For-Service Program Pharmacy Prior  
Authorization Criteria

**Effective for  
Banner Health Plans  
August 1, 2025**

Table of Contents

Acetaminophen (Dose > 4 gm)..... 17

Acthar Gel, Cortrophin Gel..... 19

Actimmune ..... 22

Adacel TDAP vaccine ..... 26

Adakveo (crizanlizumab-tmca) ..... 28

Adalimumab ..... 31

Adbry (tralokinumab-ldrm) ..... 44

ADHD Agents - AZM ..... 49

Adstiladrin (nadofaragene firadenovec-vncg)..... 55

Aduhelm (aducanumab-avwa) ..... 58

Adzynma (ADAMTS13, recombinant-krhn) ..... 69

Aemcolo ..... 71

Afinitor, Torpenz (everolimus) ..... 73

Afrezza..... 88

Agamree (vamorolone)..... 91

Airsupra (albuterol-budesonide) ..... 95

Aldurazyme - Arizona ..... 98

Alinia ..... 100

Alpha Interferons - AZM..... 102

Alpha-1 Proteinase Inhibitors..... 106

Alyftrek (vanzacaftor/tezacaftor/deutivacaftor) ..... 108

Alzheimer's Agents - AZM ..... 113

Amvuttra (vutrisiran) ..... 116

Anthelmintics - Arizona ..... 121

Anti-Parkinson's Agents (Crexont, Duopa, Vyalev) ..... 124

Anticonvulsants - AZM ..... 128

Antidepressants - AZM ..... 145

Antiemetics - AZM ..... 150

Antiglaucoma Agents - AZM.....	152
Antipsoriatic Agents.....	154
Antipsychotics - AZM .....	157
Anxiolytics - AZM .....	177
Apomorphine products (Apokyn, Onapgo) .....	180
Aqneursa (levacetylleucine) .....	183
Aquadeks.....	186
Arcalyst (riloncept) .....	188
Arikayce.....	192
Atorvaliq (atorvastatin oral suspension) .....	196
Attruby (acoramidis) .....	199
Austedo (deutetrabenazine) .....	202
Azole Antifungals.....	207
Baclofen products (Fleqsuvy, Lyvispah, Ozobax/Ozobax DS).....	219
Baxdela.....	221
Belbuca, Butrans - AZM.....	226
Benlysta (belimumab).....	234
Benznidazole.....	238
Bevacizumab Products .....	240
Bimzelx (bimekizumab-bkzx) .....	242
Bonjesta and Diclegis .....	255
Brand Over Generic Not Covered .....	257
Breast Cancer - Arizona .....	259
Breo Ellipta .....	263
Brexafemme (ibrexafungerp) .....	266
Breztri, Trelegy Ellipta - AZM.....	268
Brilinta and Effient .....	271
Buprenorphine Sublingual Tablet .....	274
Bylvay (odevixibat) .....	277

Cablivi .....	282
Cabotegravir Containing Agents .....	284
Calcium/Vitamin D .....	288
Camzyos (mavacamten) .....	290
Carbaglu (carglumic acid) .....	293
Casgevy (exagamglogene autotemcel injection).....	297
Cayston .....	304
CGRP Inhibitors - AZM.....	306
Cholbam.....	325
Cialis for BPH - AZM.....	328
Cibinqo (abrocitinib) .....	330
Cimzia (certolizumab).....	335
Colony Stimulating Factors - AZM .....	351
Compounds and Bulk Powders.....	362
Constipation Agents - AZM .....	373
Continuous Blood Glucose Monitoring Devices (CGM) .....	381
Copper Chelating Agents .....	392
Corlanor .....	396
Cosentyx (secukinumab).....	399
Coverage of Off-Label Non-FDA Approved Indications.....	412
Crenessity (crinecerfont).....	421
Cuvrior (trientine hydrochloride) .....	424
Cystaran, Cystadrops .....	427
Daliresp (roflumilast) .....	429
Daraprim .....	431
Daxxify (botulinum toxin type a injection) .....	435
Daybue (trofinetide).....	438
DDAVP (desmopressin) tablets - AZM.....	441
Declomycin - Arizona .....	443

Dificid .....	446
Dofetilide - Arizona .....	448
Dojolvi (triheptanoin) .....	450
DPP-4 Inhibitors - AZM .....	453
Drugs for Compassionate Use .....	456
Dry Eye Disease - AZM .....	458
Duexis and Vimovo - Arizona .....	462
Dupilixent (dupilumab) .....	467
Durezol .....	486
Duvyzat (givinostat) .....	488
Ebglyss (lebrikizumab-lbkz) .....	491
Ecoza (econazole) .....	496
Eculizumab (Soliris, Bkemv, Epysqli) .....	498
Elaprase - Arizona .....	511
Elevidys (delandistrogene moxeparvovec-rokl) .....	513
Elidel (pimecrolimus) cream, tacrolimus ointment.....	516
Elmiron .....	518
Emflaza (deflazacort) .....	520
Emrosi (minocycline) .....	523
Enbrel (etanercept) .....	525
Encelto (revakinagene taroretcel-lwey).....	533
Endari .....	535
Enjaymo (sutimlimab-jome) .....	537
Entocort EC.....	541
Entresto .....	543
Entyvio (vedolizumab).....	547
Epaned .....	557
Epinephrine Products .....	559
Eplerenone- Arizona .....	561

Epsolay (benzoyl peroxide) cream .....	563
Erythropoietic Agents - AZM .....	565
Estrogens- Arizona.....	579
Eucrisa .....	581
Evkeeza (evinacumab-dgnb).....	583
Evrysdi (risdiplam) .....	588
Exondys - AZM.....	594
Ezallor Sprinkle (rosuvastatin) .....	598
Fabry Disease Agents .....	601
Fasenra (benralizumab) .....	604
Fecal Microbiota Agents.....	613
Fentanyl IR .....	617
Fexmid (cyclobenzaprine 7.5mg)- Arizona .....	621
Filspari (sparsentan) .....	623
Filsuvez (birch triterpenes) .....	627
Firdapse.....	631
Flucytosine- Arizona .....	633
Forteo, Prolia, Teriparatide, Tymlos - AZM .....	635
Furoscix (furosemide injection).....	642
Galafold.....	644
Gamifant (emapalumab-lzsg) .....	646
Gattex (teduglutide) .....	649
Gaucher's Disease Agents - AZM.....	652
Generic fluticasone-salmeterol diskus, Wixela Inhub (authorized generic of Advair Diskus) .....	656
Generic tretinoin cream and gel, generic Avita cream and gel, generic atralin gel.....	658
Global Quantity Limits .....	660
GLP-1 Agonists - AZM .....	664
Glycopyrrolate Products .....	669
Gonadotropin-Releasing Hormone Agonists.....	671

Gralise, Horizant - AZM .....	686
Growth Hormone, Growth Stimulating Agents - AZM .....	688
HCG .....	719
Hemangeol .....	721
Hemophilia Clotting Factors.....	723
Hepatitis C - AZM.....	745
Hereditary Angioedema (HAE) Agents - AZM .....	755
Hetlioz, Hetlioz LQ (tasimelteon) .....	760
High Cost Oral Atypical Antipsychotics (Caplyta, Cobenfy, Fanapt, Lybalvi, Rexulti, Vraylar) .....	767
HIV (Fuzeon, Selzentry) .....	777
Hydroxychloroquine.....	780
Hyftor (sirolimus) topical gel.....	782
Igalmi (dexmedetomidine).....	784
Ilaris (canakinumab) .....	786
Imcivree (setmelanotide) .....	798
Immune Globulins - AZM.....	802
Inbrija .....	865
Infliximab Products.....	868
Ingrezza (valbenazine).....	882
Inhaled Corticosteroids - AZM .....	886
Injectable Oncology Agents.....	888
Inlyta .....	890
Insulins, Concentrated- Arizona.....	894
Inzirco (hydrochlorothiazide) oral suspension.....	896
Iqirvo (elafibranor) .....	898
Iron Chelators .....	901
Irritable Bowel Syndrome-Diarrhea .....	905
Isotretinoin - AZM .....	908
Isturisa (osilodrostat).....	913

Izervay (avacincaptad pegol) .....	916
Jesduvroq (daprodustat) .....	918
Joenja (leniolisib).....	921
Juxtapid - AZ.....	924
Jynarque .....	928
Kalydeco (ivacaftor) .....	930
Katerzia, Norliqva (amlodipine oral solution).....	934
Kepivance (palifermin) .....	936
Kerendia (finerenone).....	938
Keveyis .....	941
Kevzara (sarilumab) .....	943
Kineret (anakinra) .....	951
Korsuva (difelikefalin).....	960
Krystexxa (pegloticase) .....	963
Kuvan .....	966
LAMA, LABA - AZM .....	968
Lamzede (velmanase alfa-tycv) .....	970
Lantidra (donislecel-jujn) .....	972
Lenmeldy (atidarsagene autotemcel) .....	975
Leqembi (lecanemab-irmb).....	978
Leqvio (inclisiran) .....	984
Leucovorin- Arizona .....	990
Libervant, Nayzilam, Valtoco .....	992
Lidoderm (lidocaine) 5% patches.....	995
Likmez (metronidazole) oral suspension .....	997
Livdelzi (seladelpar) .....	1000
Livmarli (maralixibat) .....	1003
Livtencity (maribavir) .....	1007
Lodoco (colchicine) .....	1010

Long-Acting Opioid Products - AZM.....	1014
Lonhala and Yupelri - AZM .....	1034
Lucemyra (lofexidine).....	1037
Lumizyme -Arizona .....	1040
Lupkynis (voclosporin) .....	1042
Lyfgenia (lovotibeglogene autotemcel) .....	1046
Lyrica.....	1049
Lysteda .....	1055
Makena- AZM .....	1057
Marinol, Syndros .....	1059
Mepron .....	1064
Metformin products - AZM.....	1066
Migranal.....	1071
Miplyffa (arimoclomol).....	1073
Monurol .....	1077
Mozobil.....	1079
MS Agents - AZM.....	1081
Multaq .....	1084
Myalept.....	1086
Myfembree (relugolix, estradiol, and norethindrone acetate), Oriahnn (elagolix, estradiol, and norethindrone acetate capsules; elagolix capsules) .....	1089
Mytesi.....	1093
Nadolol.....	1095
Namzaric (memantine/donepezil).....	1097
Natpara .....	1099
Nemludio (nemolizumab-ilto) .....	1102
Nexiclon XR (clonidine ER) .....	1109
Nexletol (bempedoic acid) and Nexlizet (bempedoic acid-ezetimibe) .....	1111
Niktimvo (axatilimab-csfr) .....	1116
Nityr- Arizona .....	1118

Nocdurna .....	1120
Non-Preferred Drugs - AZM.....	1123
Non-Preferred Prenatal Vitamins .....	1127
Nothera .....	1133
Nourianz.....	1136
Nucala (mepolizumab) .....	1138
Nuedexta .....	1149
Nuplazid .....	1151
Nuzyra .....	1153
OAB - Overactive Bladder Agents - AZM.....	1158
Ocaliva .....	1160
Octreotide Products - AZM .....	1163
Ohtuvayre (ensifentrine).....	1169
Ojjaara (momelotinib).....	1174
Olumiant (baricitinib).....	1176
Opzelura (ruxolitinib).....	1189
Oral Oncology Agents .....	1195
Orencia (abatacept).....	1197
Orfadin (nitisinone).....	1202
Orilissa .....	1204
Orkambi - AZM .....	1208
Osphena - Arizona .....	1211
Otezla (apremilast) .....	1213
Oxervate .....	1218
Palforzia [Peanut (Arachis hypogaea)] .....	1220
Palynziq .....	1224
Panretin .....	1227
Pediculicides - AZM .....	1230
Pedmark (sodium thiosulfate injection, solution) .....	1232

Piasky (crovalimab).....	1234
Pombiliti (cipaglucosidase alfa-atga) .....	1237
Pradaxa Pellet Packs .....	1240
Praluent .....	1242
Preferred Drugs- Arizona .....	1251
Presbyopia Agents .....	1253
Pretomanid .....	1256
Prevymis (letermovir) .....	1258
Primary Hyperoxaluria (PH1) Agents [Oxlumo (lumasiran), Rivfloza (nedosiran)].....	1261
Procysbi.....	1266
Progesterone - Non-Oral .....	1268
Proton Pump Inhibitors (PPIs) .....	1270
Provigil, Nuvigil.....	1272
Pulmonary Arterial Hypertension (PAH) Agents .....	1276
Pulmozyme .....	1282
Pyrukynd (mitapivat) .....	1284
Qalsody (tofersen) .....	1287
Qutenza (capsaicin).....	1290
Radicava (edaravone) .....	1293
Ranolazine products .....	1296
Rayos .....	1299
Reblozyl (luspatercept-aamt) .....	1302
Recorlev (levoketoconazole).....	1307
Rectiv .....	1310
Regranex.....	1312
Relyvrio (sodium phenylbutyrate and taurursodiol) .....	1314
Repatha .....	1317
Respiratory Syncytial Virus (RSV) Vaccines .....	1326
Retinal Vascular Disease Agents.....	1329

Reyvow - Arizona .....	1337
Rezdiffra (resmetirom) .....	1341
Rezurock (belumosudil) .....	1344
Rezzayo (rezafungin) .....	1347
Rhofade .....	1349
Rinvoq (upadacitinib) .....	1351
Rystiggo (rozanolixizumab) .....	1382
Rytelo (imetelstat) .....	1385
Samsca .....	1388
Sedative Hypnotics - AZM.....	1390
Serevent Diskus - Arizona.....	1394
SGLT-2 Inhibitors - AZM .....	1397
Shingrix (zoster vaccine recombinant, adjuvanted) .....	1403
Short-Acting Opioid Products - AZM .....	1405
Signifor .....	1419
Siliq (brodalumab) .....	1421
Skyclarys (omaveloxolone).....	1438
Skyrizi (risankizumab-rzaa).....	1440
Sodium Oxybate Products (Lumryz, Xyrem, Xywav) .....	1453
Sofdra (sofpironium) .....	1460
Sohonos (palovarotene) .....	1463
Somatuline Depot (lanreotide).....	1466
Somavert .....	1471
Soriatane .....	1474
Sotyktu (deucravacitinib) .....	1477
Spinraza- Arizona.....	1485
Spiriva (generic tiotropium) products .....	1493
Spravato, ketamine - AZM.....	1495
Strensiq .....	1504

Sublingual Immunotherapy (SLIT) .....	1509
Sublocade, Brixadi - AZM .....	1517
Suboxone - AZM .....	1521
Sucraid .....	1525
Sunlenca (lenacapavir sodium) .....	1528
Sunosi (solriamfetol).....	1530
Sutent.....	1535
Syfovre (pegcetacoplan).....	1543
Symdeko.....	1545
Symlin .....	1548
Synagis .....	1550
Systane, Refresh, Gonak, Genteal, Tears Naturale.....	1564
Talicia and rifabutin .....	1566
Taltz (ixekizumab) .....	1572
Targretin.....	1585
Tarpeyo (budesonide) .....	1588
Tasigna .....	1591
Tecelra (afamitresgene autoleucel) .....	1596
Tegsedi .....	1599
Tepezza (teprotumumab-trbw).....	1603
Test Strips .....	1605
Testosterone - AZM .....	1609
Tezruly (terazosin) oral solution.....	1616
Tezspire (tezepelumab-ekko) .....	1618
Thalomid .....	1622
Thrombopoiesis Stimulating Agents.....	1631
Tobramycin Inhalation - ARIZONA .....	1638
Tocilizumab (Actemra, Tofidence, Tyenne).....	1641
Topical Capsaicin Products .....	1648

Topical NSAIDs.....	1650
Trastuzumab Products.....	1655
Tremfya (guselkumab).....	1657
Tretinoin Topical .....	1670
Trikafta (elexacaftor/tezacaftor/ivacaftor).....	1672
Triptans - AZM .....	1676
Tryngolza (olezarsen sodium) .....	1687
Tryvio (aprocitentan) .....	1690
Twynéo (tretinoin-benzoyl peroxide 0.1-3% cream) .....	1694
Tykerb .....	1696
Tysabri (natalizumab).....	1703
Ultomiris (ravulizumab-cwvz) .....	1709
Uplizna (inebilizumab-cdon) .....	1715
Urea Cycle Disorder Agents.....	1719
Ustekinumab .....	1723
Valchlor .....	1734
Valsartan oral solution .....	1737
Vecamyl.....	1739
Velphoro (sucroferric oxyhydroxide), Auryxia (ferric citrate).....	1741
Velsipity (etrasimod) .....	1743
Vemlidy.....	1749
Veopoz (pozelimab) .....	1752
Veozah (fezolinetant) .....	1755
Verkazia (cyclosporine ophthalmic emulsion 0.1%).....	1757
Vijoice (alpelisib) .....	1760
Vitamin B-12.....	1763
Vitamin C.....	1765
Vitamin D.....	1767
Vivjoa (oteseconazole).....	1769

Vonjo (pacritinib) .....	1772
Vonoprazan Containing Agents .....	1775
Voxzogo (vosoritide) .....	1780
Vtama (tapinarof) .....	1784
Vyjuvek (beremagene geperpavec-svdt) .....	1789
Vykat XR (diazoxide choline) .....	1792
Vyndaqel and Vyndamax .....	1795
Vyvgart (efgartigimod alfa-fcab) .....	1800
Wainua (eplontersen).....	1805
Wakix (pitolisant) .....	1808
Wegovy (semaglutide).....	1812
Xalkori .....	1817
Xdemvy (lotilaner).....	1822
Xeljanz, Xeljanz XR (tofacitinib) .....	1824
Xenazine .....	1832
Xenleta .....	1836
Xenpozyme (olipudase alfa).....	1839
Xermelo .....	1842
Xhance (fluticasone nasal spray) .....	1844
Xolair (omalizumab).....	1846
Xolremdi (mavorixafor).....	1858
Xopenex Respules .....	1861
Xphozah (tenapanor).....	1863
Xromi (hydroxyurea) .....	1866
Xuriden .....	1869
Yorvipath (palopegteriparatide).....	1871
Zeposia (ozanimod) .....	1873
Zinplava (bezlotoxumab) .....	1879
Zolgensma (onasemnogene abeparvovec-xioi) .....	1881

Zontivity.....	1884
Zortress - ARIZONA.....	1886
Zoryve (roflumilast).....	1888
Ztalmy (ganaxolone) .....	1893
Zunveyl (benzgalantamine).....	1896
Zurzuvae (zuranolone).....	1898
Zynteglo (betibeglogene autotemcel) .....	1900
Zyvox.....	1904

Acetaminophen (Dose > 4 gm)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144743
<b>Guideline Name</b>	Acetaminophen (Dose > 4 gm)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/21/2024
-----------------	-----------

## 1 . Criteria

Product Name:Acetaminophen (Dose > 4gm)	
Diagnosis	DUR Reject 88 (Total APAP > 4 g)
Guideline Type	DUR Reject 88
<b>Approval Criteria</b>  1 - Requests for acetaminophen dosages greater than 4000mg per day should be denied. The total dose of acetaminophen (cumulative total daily dose of 4000mg) is not supported by the Food and Drug Administration (FDA).	
Notes	Note: Reject message: "DUR1:APAP = Total APAP >4g; Verify dose; EnterO/R -"

## 2 . Revision History

Date	Notes
3/21/2024	Guideline type changed from Administrative to DUR Reject 88

Acthar Gel, Cortrophin Gel

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-224208
<b>Guideline Name</b>	Acthar Gel, Cortrophin Gel
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Acthar Gel vial	
Diagnosis	Infantile spasm (i.e., West Syndrome)*
Approval Length	4 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of infantile spasms (i.e., West Syndrome)*  <b>AND</b>	

**2** - Patient is less than 2 years old

**AND**

**3** - Both of following:

**3.1** Initial dose: 75 units per meters squared intramuscular (IM) twice daily for 2 weeks

**AND**

**3.2** After 2 weeks, dose should be tapered according to the following schedule: 30 units per meters squared IM in the morning for 3 days; 15 units per meters squared IM in the morning for 3 days; 10 units per meters squared IM in the morning for 3 days; 10 units per meters squared IM every other morning for 6 days (3 doses)

Notes	*Note: Acthar Gel is not medically necessary for treatment of acute exacerbations of multiple sclerosis.
-------	--

Product Name: Acthar Gel, Cortrophin	
Diagnosis	Opsoclonus-myoclonus syndrome (i.e., OMS, Kinsbourne Syndrome)*
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Opsoclonus-myoclonus syndrome (i.e., OMS, Kinsbourne Syndrome)*	
<b>AND</b>	
2 - For Cortrophin requests ONLY: Trial and failure or intolerance to Acthar Gel (verified via paid pharmacy claims or submission of medical records/chart notes)	
Notes	*Note: Acthar Gel is not medically necessary for treatment of acute exacerbations of multiple sclerosis.

## 2 . Revision History

Date	Notes
3/26/2025	Added new GPIs for Acthar and Cortrophin

Actimmune

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99673
<b>Guideline Name</b>	Actimmune
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Actimmune	
Diagnosis	Chronic Granulomatous Disease (CGD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic granulomatous disease	

Product Name:Actimmune	
Diagnosis	Chronic Granulomatous Disease (CGD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Actimmune</p>	

Product Name:Actimmune	
Diagnosis	Severe, Malignant Osteopetrosis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of severe, malignant osteopetrosis</p>	

Product Name:Actimmune	
Diagnosis	Severe, Malignant Osteopetrosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Actimmune</p>	

Product Name:Actimmune	
Diagnosis	Primary Cutaneous Lymphomas
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has ONE of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Mycosis fungoides (MF)</li> <li>• Sézary syndrome (SS)</li> </ul>	

Product Name:Actimmune	
Diagnosis	Primary Cutaneous Lymphomas
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Actimmune</p>	

Product Name:Actimmune	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Actimmune will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.

<b>Product Name:Actimmune</b>	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Actimmune therapy</p>	

## 2 . Revision History

Date	Notes
6/7/2021	7.1 Implementation

Adacel TDAP vaccine

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-124866
<b>Guideline Name</b>	Adacel TDAP vaccine
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Adacel	
Diagnosis	Pregnant Patients 19 years of age and older*
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 2 - Vaccine is being used to prevent pertussis in infants younger than 2 months of age  <b>AND</b>	

1 - Patient is 19 years of age or older

**AND**

3 - Both of the following:

- Patient is pregnant
- Vaccine is being administered during 3rd trimester of pregnancy

Notes

\*Note: Patients under 19 years of age must get immunization from PC P or pediatrician through the VFC (Vaccines For Children) Program

## 2 . Revision History

Date	Notes
4/20/2023	New program

Adakveo (crizanlizumab-tmca)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-226191
<b>Guideline Name</b>	Adakveo (crizanlizumab-tmca)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Adakveo	
Diagnosis	Sickle cell disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of sickle cell disease, identified by any genotype	

**AND**

**2** - ONE of the following:

**2.1** BOTH of the following:

- Age 16 to 20 years
- Prescriber attests the service is medically necessary to correct or ameliorate a defect, a condition, or a physical or mental illness in an eligible patient

**OR**

**2.2** Age greater than or equal to 21 years

**AND**

**3** - Patient has experienced at least two vaso-occlusive crises within the past 12 months

**AND**

**4** - Trial and failure or inadequate response, contraindication, or intolerance to hydroxyurea

Product Name:Adakveo	
Diagnosis	Sickle cell disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy (e.g., reduction in annual rate of vaso-occlusive events, increased time between each vaso-occlusive event)	

## 2 . Revision History

Date	Notes
3/26/2025	Added step through HXU to initial auth, added reauth criteria.

Adalimumab

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-325188
<b>Guideline Name</b>	Adalimumab
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

#### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

#### 1 . Criteria

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderately to severely active rheumatoid arthritis  <b>AND</b>	

**2** - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to a 3 month trial of ONE non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima

Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

### Approval Criteria

**1** - Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis (PJIA)

**AND**

**2** - Paid claims or submission of medical records (e.g., chart notes) confirming a history of failure to a minimum duration of a 6-week trial, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:

- leflunomide
- methotrexate

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima

Diagnosis	Ankylosing Spondylitis (AS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of active ankylosing spondylitis (AS)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to TWO NSAIDs (non-steroidal anti-inflammatory drugs) (e.g., ibuprofen, naproxen) at maximally indicated doses, each used for at least 4 weeks within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a rheumatologist</p>	

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Rheumatoid Arthritis (RA), Polyarticular Juvenile Idiopathic Arthritis (PJIA), Ankylosing Spondylitis (AS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p>	

2 - Prescribed by or in consultation with a rheumatologist

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of active psoriatic arthritis (PsA)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with ONE of the following:</p> <ul style="list-style-type: none"><li>• Rheumatologist</li><li>• Dermatologist</li></ul>	

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of moderate to severe chronic plaque psoriasis (PsO)</p> <p><b>AND</b></p> <p>2 - Greater than or equal to 3% body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis</p> <p><b>AND</b></p> <p>3 - Both of the following:</p> <p><b>3.1</b> Paid claims or submission of medical records (e.g., chart notes) documenting history of</p>	

failure to ONE of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Vtama
- Zoryve 0.3% cream

**AND**

**3.2** Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**4** - Prescribed by or in consultation with a dermatologist

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Hidradenitis Suppurativa (HS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of moderate to severe hidradenitis suppurativa (i.e., Hurley Stage II or III)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to at least ONE oral antibiotic (e.g., doxycycline, clindamycin, rifampin) at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced</p>	

**AND**

**3** - Prescribed by or in consultation with a dermatologist

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Plaque Psoriasis (PsO), Hidradenitis Suppurativa (HS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a dermatologist	

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Adult Crohn's Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active Crohn's disease	
<b>AND</b>	

**2 - ONE of the following:**

**2.1** Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)
- Azathioprine (Imuran)
- 6-mercaptopurine (Purinethol)
- Methotrexate (Rheumatrex, Trexall)

**OR**

**2.2** Patient has lost response or intolerant to infliximab (e.g., Remicade, Inflectra, Renflexis)

**AND**

**3 - Prescribed by or in consultation with a gastroenterologist**

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Pediatric Crohn's Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active Crohn's disease	
<b>AND</b>	
2 - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:	
<ul style="list-style-type: none"><li>• Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)</li></ul>	

- Azathioprine (Imuran)
- 6-mercaptopurine (Purinethol)
- Methotrexate (Rheumatrex, Trexall)

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Ulcerative Colitis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of moderately to severely active ulcerative colitis</p> <p><b>AND</b></p> <p><b>2</b> - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)</li> <li>• 6-mercaptopurine (Purinethol)</li> <li>• Azathioprine (Imuran)</li> <li>• Aminosalicylates (e.g., mesalamine, sulfasalazine)</li> </ul> <p><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with a gastroenterologist</p>	

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Adult Crohn's Disease, Pediatric Crohn's Disease, Ulcerative Colitis

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a gastroenterologist</p>	

Product Name: Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Uveitis (UV)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of non-infectious uveitis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Uveitis is classified as ONE of the following:</p> <ul style="list-style-type: none"> <li>• intermediate</li> <li>• posterior</li> <li>• panuveitis</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Paid claims or submission of medical records (e.g., chart notes) documenting history of</p>	

failure to at least ONE corticosteroid (e.g., prednisolone, prednisone) at maximally indicated dose within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to at least ONE systemic non-biologic immunosuppressant (e.g., methotrexate, cyclosporine, azathioprine, mycophenolate) at a maximally indicated dose within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**5** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Ophthalmologist

Product Name:Preferred: Unbranded Adalimumab-fkjp (Biocon manufacturer), Hadlima	
Diagnosis	Uveitis (UV)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with ONE of the following:	
<ul style="list-style-type: none"><li>• Rheumatologist</li><li>• Ophthalmologist</li></ul>	

Product Name:Non-Preferred*: Brand Humira, Abrilada, Amjevita, Cyltezo, Brand Adalimumab-adbm, Hulio, Hyrimoz, Brand Adalimumab-adaz, Idacio, Brand Adalimumab-aacf, Brand Adalimumab-ryvk, Simlandi, Yuflyma, Brand Adalimumab-aaty, Yusimry, and newly launched adalimumab products	
Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred products are Unbranded Adalimumab-fkjp and Hadlima.  **Please note: The drug(s) listed above may require additional review.</p>	
Notes	*Patients must use preferred adalimumab biosimilar (refer to background table for list of preferred adalimumab biosimilars).

## 2 . Background

Benefit/Coverage/Program Information			
Preferred Adalimumab Biosimilars			
NDC	Product Label	GPI-14	GPI-14 Description
49502041602	ADALIMU-FKJP KIT 40/0.8ML	6627001535F520	ADALIMUMAB-FKJP AUTO-INJECTOR KIT 40 MG/0.8ML
83257002232	ADALIMU-FKJP KIT 40/0.8ML	6627001535F520	ADALIMUMAB-FKJP AUTO-INJECTOR KIT 40 MG/0.8ML
49502041702	ADALIMU-FKJP KIT 20/0.4ML	6627001535F810	ADALIMUMAB-FKJP PREFILLED SYRINGE KIT 20 MG/0.4ML
83257002042	ADALIMU-FKJP KIT 20/0.4ML	6627001535F810	ADALIMUMAB-FKJP PREFILLED SYRINGE KIT 20 MG/0.4ML
49502041802	ADALIMU-FKJP KIT 40/0.8ML	6627001535F820	ADALIMUMAB-FKJP PREFILLED SYRINGE KIT 40 MG/0.8ML

83257002142	ADALIMU-FKJP KIT 40/0.8ML	6627001535F820	ADALIMUMAB-FKJP PREFILLED SYRINGE KIT 40 MG/0.8ML
78206018701	HADLIMA PUSH INJ 40/0.4ML	6627001520D510	ADALIMUMAB-BWWD SOLN AUTO-INJECTOR 40 MG/0.4ML
78206018401	HADLIMA PUSH INJ 40/0.8ML	6627001520D520	ADALIMUMAB-BWWD SOLN AUTO-INJECTOR 40 MG/0.8ML
78206018601	HADLIMA INJ 40/0.4ML	6627001520E510	ADALIMUMAB-BWWD SOLN PREFILLED SYRINGE 40 MG/0.4ML
78206018301	HADLIMA INJ 40/0.8ML	6627001520E520	ADALIMUMAB-BWWD SOLN PREFILLED SYRINGE 40 MG/0.8ML

### 3 . Revision History

Date	Notes
7/16/2025	Updated preferred agents, updated criteria throughout, updated NP s ection verbiage.

Adbry (tralokinumab-ldrm)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-150052
<b>Guideline Name</b>	Adbry (tralokinumab-ldrm)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Adbry	
Diagnosis	Atopic Dermatitis
Approval Length	6 Months*
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderate to severe atopic dermatitis	

**AND**

**2** - Submission of documentation (e.g., chart notes) demonstrating one of the following:

- Involvement of at least 10% body surface area (BSA)
- SCORing Atopic Dermatitis (SCORAD) index value of at least 25 [A]

**AND**

**3** - Patient is 12 years of age or older

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist/Immunologist

**AND**

**5** - History of failure, contraindication, or intolerance to BOTH of the following topical therapies: (document drug, date of trial, and/or contraindication to medication)\*\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole) ointment

Notes

\*QL Override (For new starts only): Enter 2 PAs as follows: First PA: Approve 6 syringes per 28 days for one month; Second PA: Approve 4 syringes per 28 days (no overrides needed) for the remaining 11 months. (Adbry is hard-coded with a quantity of 4 syringes per 28 days).  
\*\*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication

Product Name:Adbry

Diagnosis

Atopic Dermatitis

Approval Length

12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of documentation (e.g., chart notes) demonstrating positive clinical response to therapy as evidenced by at least ONE of the following:</p> <ul style="list-style-type: none"> <li>• Reduction in body surface area involvement from baseline</li> <li>• Reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline [A]</li> </ul>	

## 2 . Background

Clinical Practice Guidelines			
Table 1. Relative potencies of topical corticosteroids [2]			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment, gel	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream, lotion	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25

	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05
	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment, lotion	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream, lotion	0.1
	Triamcinolone acetonide	Cream, ointment, lotion	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

### 3 . Revision History

Date	Notes
7/20/2024	Added new GPI

ADHD Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242227
<b>Guideline Name</b>	ADHD Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Adderall, generic amphetamine/dextroamphetamine tablets, Brand Adderall XR, generic amphetamine/dextroamphetamine ER capsules, Brand Aptensio XR, generic atomoxetine, Brand Concerta, Brand Daytrana, generic dexamethylphenidate tablets, generic dexamethylphenidate ER, generic dextroamphetamine tablets, Brand Focalin, Brand Focalin XR, generic lisdexamfetamine capsules and chewables, Brand Methylin solution, generic methylphenidate solution, generic methylphenidate tablets, generic methylphenidate ER tablets, generic methylphenidate ER (CD) capsules, generic methylphenidate ER (LA) capsules, generic methylphenidate ER (XR) capsules, generic methylphenidate patch, Brand Ritalin, Brand Ritalin LA, Brand Strattera, Brand Vyvanse capsules and chewables, Brand Zenzedi	
Diagnosis	PA Required for Children Under 6 Years Old
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - The requesting clinician has documented that the child has a diagnosis of attention deficit hyperactivity disorder (ADHD)

**AND**

2 - The requesting clinician has documented that psychosocial issues have been evaluated before request for ADHD medications

**AND**

3 - The requesting clinician has documented non-medication alternatives that have been attempted before request for ADHD medications

**AND**

4 - The requested dose does NOT exceed the Food and Drug Administration (FDA) recommended maximum daily dosage unless the provider has submitted clinical justification for the dose exceeding the FDA maximum

Product Name: Brand Intuniv, generic guanfacine IR/ER, Brand Kapvay, generic clonidine IR/ER

Diagnosis	PA Required for Children Under 6 Years Old
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - One of the following:

1.1 All of the following:

**1.1.1** The requesting clinician has documented that the child has a diagnosis of attention deficit hyperactivity disorder (ADHD)

**AND**

**1.1.2** The requesting clinician has documented that psychosocial issues have been evaluated before request for ADHD medications

**AND**

**1.1.3** The requesting clinician has documented non-medication alternatives that have been attempted before request for ADHD medications

**AND**

**1.1.4** The requested dose does NOT exceed the Food and Drug Administration (FDA) recommended maximum daily dosage unless the provider has submitted clinical justification for the dose exceeding the FDA maximum

**OR**

**1.2** Both of the following:

**1.2.1** Diagnosis of insomnia

**AND**

**1.2.2** Trial and failure, contraindication, or intolerance to melatonin

Product Name:NON-PREFERRED DRUGS: Xelstrym patch	
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - The patient has a history of failure, contraindication, or intolerance to a trial to THREE of the following preferred products\*:

- Brand Adderall
- generic amphetamine/dextroamphetamine tablets
- Brand Concerta ER
- generic dexamethylphenidate tablets
- Brand Focalin XR
- Brand Methylin solution
- generic methylphenidate tablets
- Brand Ritalin LA
- generic methylphenidate ER (CD) capsules
- Vyvanse capsules
- generic atomoxetine
- generic clonidine ER
- generic guanfacine ER
- generic dextroamphetamine tablets

**AND**

2 - The patient has a history of failure, contraindication, or intolerance to Daytrana

Notes

\*Alternatives may require prior authorization

Product Name:NON-PREFERRED DRUGS: Onyda XR

Approval Length 12 month(s)

Guideline Type Prior Authorization

**Approval Criteria**

1 - The patient has a history of failure, contraindication, or intolerance to a trial to THREE of the following preferred products\*:

- Brand Adderall
- generic amphetamine/dextroamphetamine tablets
- Brand Concerta ER
- generic dexamethylphenidate tablets
- Brand Focalin XR
- Brand Methylin solution
- generic methylphenidate tablets
- Brand Ritalin LA

- generic methylphenidate ER (CD) capsules
- Vyvanse capsules
- generic atomoxetine
- generic clonidine ER
- generic guanfacine ER
- generic dextroamphetamine tablets

**AND**

**2** - The patient has a history of failure, contraindication, or intolerance to generic clonidine patch

Notes

\*Alternatives may require prior authorization

Product Name:NON-PREFERRED DRUGS: Brand Adderall XR, Brand Adhansia XR, Brand Adzenys XR-ODT, generic amphetamine IR tablets, generic amphetamine ER suspension, generic amphetamine/dextroamphetamine ER capsules, Brand Aptensio XR, Brand Azstarys, Brand Cotempla XR-ODT, Brand Desoxyn, Brand Dexedrine, generic dextroamphetamine oral solution, generic dextroamphetamine IR tablet, generic dextroamphetamine ER, Brand Dyanavel XR (oral suspension and chewable tablets), Brand Evekeo, Brand Evekeo ODT, Brand Focalin, Brand Focalin XR, Brand Intuniv, Brand Jornay PM, Brand Kapvay, generic lisdexamfetamine capsules and chewables, generic methamphetamine , generic methylphenidate chewable, generic methylphenidate patch, generic methylphenidate soln, generic methylphenidate ER tablets, generic methylphenidate ER (LA) capsules, generic methylphenidate ER (XR) capsules, Brand Mydayis, Brand Procentra, Brand Qelbree, Brand Quillichew ER, Brand Quillivant XR, Relexxii, Brand Ritalin, Brand Strattera, Brand Vyvanse chewables, Brand Zenzedi

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - The patient has a history of failure, contraindication, or intolerance to a trial to FOUR of the following preferred products\*:

- Brand Adderall
- generic amphetamine/dextroamphetamine tablets
- Brand Concerta ER
- Daytrana
- generic dexmethylphenidate tablets
- Brand Focalin XR
- Brand Methylin solution
- generic methylphenidate tablets

- Brand Ritalin LA
- generic methylphenidate ER (CD) capsules
- Vyvanse capsules
- generic atomoxetine
- generic clonidine ER
- generic guanfacine ER
- generic dextroamphetamine tablets

**AND**

**2 -** If the request is for a multi-source brand medication (i.e., MSC O) ONE of the following:

**2.1** BOTH of the following:

- The brand is being requested because of an adverse reaction, allergy or sensitivity to the generic and the prescriber must attest to submitting the FDA MedWatch Form for allergic reactions to the medications
- The patient has tried three generic formulations (if available) of the requested brand product

**OR**

**2.2** ONE of the following:

- The brand is being requested due to a therapeutic failure with the generic (please provide reason for therapeutic failure)
- The brand is being requested because transition to the generic could result in destabilization of the patient (rationale must be provided)
- Special clinical circumstances exist that preclude the use of the generic equivalent of the multi-source brand medication for the patient (rationale must be provided)

Notes	*Alternatives may require prior authorization
-------	---

## 2 . Revision History

Date	Notes
4/24/2025	Updated step req for NP section, aligns with MSC O section of NP admin guideline

Adstiladrin (nadofaragene firadenovec-vncg)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-133805
<b>Guideline Name</b>	Adstiladrin (nadofaragene firadenovec-vncg)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Adstiladrin	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting ALL of the following: 1.1 Diagnosis of high-risk, non-Muscle Invasive Bladder Cancer (NMIBC)	

**AND**

**1.2** One of the following:

- Tumor is carcinoma in situ (CIS)
- Ta/T1 high grade disease

**AND**

**1.3** Patient is not eligible for or has elected not to undergo cystectomy

**AND**

**1.4** Patient has received an adequate course of Bacillus Calmette Guérin (BCG) therapy defined as the administration of at least 5 of 6 doses of an initial induction course plus one of the following:

- At least two of three doses of maintenance therapy
- At least two of six doses of a second induction course

**AND**

**1.5** Tumor is BCG unresponsive as defined by one of the following:

- Persistent disease following adequate BCG therapy
- Disease recurrence after an initial tumor-free state following adequate BCG therapy
- T1 disease following a single induction course of BCG

**AND**

**1.6** The patient has had all resectable disease (Ta and T1 components) removed

**AND**

**1.7** The patient does not have extra-vesical (i.e., urethra, ureter, or renal pelvis), muscle invasive (T2-T4), or metastatic urothelial carcinoma

Product Name:Adstiladrin	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on therapy	

## 2 . Revision History

Date	Notes
9/26/2023	New Program

Aduhelm (aducanumab-avwa)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-107262
<b>Guideline Name</b>	Aduhelm (aducanumab-avwa)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/17/2022
-----------------	-----------

## 1 . Criteria

Product Name:Aduhelm	
Diagnosis	Alzheimer's Disease - MEDICARE PART B*
Approval Length	6 month(s)
Guideline Type	Medicare Part B
<b>Approval Criteria</b> 1 - Requested medication is billed through Medicare Part B  <b>AND</b>	

<b>2 - Submission of documentation confirming patient is enrolled in a CMS approved prospective comparative study</b>	
Notes	*Note: THIS SECTION SHOULD ONLY BE USED FOR DUAL ELIGIBLE MEMBERS (WILL HAVE AZMDUAL PLAN CODE) COVERED UNDER MEDICARE PART B THAT ARE REQUESTING SECONDARY COVERAGE.

<b>Product Name:Aduhelm</b>	
Diagnosis	Alzheimer's Disease - MEDICARE PART D*
Approval Length	None
Guideline Type	Prior Authorization requests from providers from Medicare Part D for Dual Eligible Members
<p><b>Approval Criteria</b></p> <p>1 - Requested medication is billed through Medicare Part D</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Requests for coverage of Aduhelm (aducanumab) are not authorized and will not be approved under Part D</p>	
Notes	*Note: THIS SECTION SHOULD ONLY BE USED FOR DUAL ELIGIBLE MEMBERS (WILL HAVE AZMDUAL PLAN CODE). APPROVAL LENGTH: NONE - REQUESTS FOR ADUHELM ARE NOT COVERED UNDER MEDICARE PART D AND SHALL BE DENIED AS A BENEFIT EXCLUSION.

<b>Product Name:Aduhelm</b>	
Diagnosis	Alzheimer's Disease - FEE-FOR-SERVICE
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Diagnosis of one of the following:

- Mild cognitive impairment (MCI) due to Alzheimer's Disease (AD)
- Mild dementia due to Alzheimer's Disease (AD)

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values, examination histories) documenting the basis for diagnosis, including all of the following:

**2.1** Documentation of a comprehensive history and neurological examination, inclusive of a description of the nature and duration of cognitive symptoms within the previous 3 months

**AND**

**2.2** Medical records documenting baseline (within the previous three months) cognitive function based on ONE of the following objective assessments:

- Mini-Mental State Examination (MMSE) score  $\geq 24$
- Montreal Cognitive Assessment (MoCA) score  $\geq 15$

**AND**

**2.3** Medical records documenting confirmed evidence of clinically significant AD neuropathology based on ONE of the following:

- Cerebral Spinal Fluid (CSF) biomarkers
- Amyloid positron emission tomography (PET)

**AND**

**3** - Patient has received recent (within the previous 3 months) baseline brain magnetic resonance imaging (MRI) prior to initiating treatment

**AND**

**4** - Patient does not have significant cerebrovascular disease as established by brain MRI showing any of the following:

- Acute or sub-acute hemorrhage
- Prior macro-hemorrhage or prior subarachnoid hemorrhage (unless finding is not due to an underlying structural or vascular hemorrhage)
- 4 or more brain microhemorrhages
- Cortical infarct
- More than 1 lacunar infarct
- Superficial siderosis
- History of diffuse white matter disease

**AND**

**5** - Patient does not have any of the following non-AD neurodegenerative disorders:

- Probable dementia with Lewy bodies by consensus criteria
- Suspected frontotemporal degeneration
- Dementia in down syndrome

**AND**

**6** - Patient does not have any of the following exclusionary neurological or psychiatric conditions:

- Uncontrolled seizure disorder
- Uncontrolled mood disorder, anxiety disorder, or psychosis
- Substance use disorder active in the past 2 years

**AND**

**7** - Patient does not have any of the following cardiovascular conditions:

- Uncontrolled hypertension
- Coronary artery disease (including unstable angina and myocardial infarction)
- Heart failure
- Arrhythmia
- Clinically significant carotid atherosclerosis and/or peripheral arterial disease

**AND**

**8** - Both of the following:

- Patient is not currently taking an anticoagulant or antiplatelet agent (unless aspirin 325 mg/day or less)
- Patient has no history of transient ischemic attack (TIA), stroke, or unexplained loss of consciousness within previous year prior to initiating treatment

**AND**

**9** - Patient does not have any uncontrolled clinically significant chronic medical condition (e.g., liver disease, kidney disease, pulmonary disease, autoimmune disease requiring chronic immunosuppression, malignant neoplasm, active chronic infection [HIV, HCV], poorly controlled diabetes mellitus)

**AND**

**10** - Prescribed dosing is in accordance with the United States Food and Drug Administration approved labeling

**AND**

**11** - Prescribed by or in consultation with one of the following:

- Neurologist
- Geriatrics specialist

**AND**

**12** - Prescriber attests that the patient and/or authorized representative (e.g., power of attorney, invoked health care proxy) has shared in decision-making and has been informed on the known and potential risks and lack of established clinical benefit associated with Aduhelm (aducanumab-avwa) treatment

**AND**

**13** - Therapy should be discontinued permanently and the request should be denied if one or more of the following apply:

<ul style="list-style-type: none"> <li>• If the patient has had <math>\geq 10</math> new incident microhemorrhages, regardless of clinical severity (including asymptomatic)</li> <li>• If the patient had a serious event [Serious events include concern for immediate risk of death (a life-threatening event); inpatient hospitalization or prolongation of existing hospitalization due to symptoms; new persistent or significant disability/incapacity]</li> <li>• If the patient has had <math>\geq 3</math> new incident areas of superficial siderosis, regardless of clinical severity (including asymptomatic) therapy should be discontinued permanently and the request should be denied</li> </ul>	
Notes	<p>*NOTE: If the patient has had <math>\geq 10</math> new incident microhemorrhages, regardless of clinical severity (including asymptomatic) therapy should be discontinued permanently and the request should be denied.</p> <p>*NOTE: If the patient had a serious event, therapy should be discontinued. †</p> <p>*NOTE: If the patient has had <math>\geq 3</math> new incident areas of superficial siderosis, regardless of clinical severity (including asymptomatic) therapy should be discontinued permanently and the request should be denied.</p> <p>†Serious events include concern for immediate risk of death (a life-threatening event); inpatient hospitalization or prolongation of existing hospitalization due to symptoms; new persistent or significant disability/incapacity.</p> <p>‡Requests should be evaluated case-by-case with clinical review and MD advisor.</p>

Product Name:Aduhelm	
Diagnosis	Alzheimer's Disease - FEE-FOR-SERVICE
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Prescribed dosing is in accordance with the United States Food and Drug Administration approved labeling</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Follow-up MRIs have been conducted at the following timeframes:</p>	

- Week 14 (after 4th infusion, prior to first 6 mg/kg dose)
- Week 22 (after 6th infusion, prior to first 10 mg/kg dose)
- Week 30 (after 8th infusion, prior to third 10 mg/kg dose)
- Week 42 (after 11th infusion, prior to sixth 10 mg/kg dose)
- Every 6 months thereafter

**AND**

**3** - Patient's diagnosis continues to be mild cognitive impairment or mild dementia stage due to Alzheimer's disease as established by one of the following examination scales:

**3.1** One of the following:

- Mini Mental State Exam (MMSE) score  $\geq 24$
- Montreal Cognitive Assessment (MoCA) score  $\geq 15$

**OR**

**3.2** Both of the following:

- MMSE  $<24$  or MoCA  $<15$
- Rate of decline was slower than expected ( $<2$  points/year)

**AND**

**4** - ONE of the following (ARIA-H, microhemorrhages):

- Patient has had no new incident microhemorrhage
- Patient has had 1 to 4 new incident microhemorrhage(s) AND microhemorrhages are asymptomatic (no clinical symptoms)
- Patient has had 5 to 9 new incident microhemorrhages AND microhemorrhages are asymptomatic (no clinical symptoms) AND the microhemorrhages have been stabilized
- Patient has had 1 to 9 new incident microhemorrhages AND microhemorrhages resulted in mild, moderate or severe clinical symptoms AND the microhemorrhages have been stabilized

**AND**

**5** - ONE of the following (ARIA-H, superficial siderosis)

- Patient has had no new incident areas of superficial siderosis
- Patient has had 1 new incident area of superficial siderosis AND superficial siderosis is asymptomatic (no clinical symptoms)
- Patient has had 2 new incident areas of superficial siderosis AND superficial siderosis is asymptomatic (no clinical symptoms) AND the superficial siderosis has been stabilized
- Patient has had 1 to 2 new incident areas of superficial siderosis AND superficial siderosis resulted in mild, moderate or severe clinical symptoms AND the superficial siderosis has been stabilized

**AND**

**6 - ONE of the following (ARIA-E)**

- Patient has had no new ARIA-E
- Patient has mild ARIA-E on MRI AND ARIA-E is asymptomatic (no clinical symptoms)
- Patient has had moderate or severe ARIA-E on MRI AND ARIA-E is asymptomatic (no clinical symptoms) AND the ARIA-E is stable
- Patient has had mild, moderate or severe ARIA-E on MRI AND ARIA-E resulted in mild, moderate or severe clinical symptoms AND the ARIA-E is stable

**AND**

**7 - One of the following:**

**7.1 Patient does not meet ANY of the following:**

- Initiation of anticoagulation
- Development of active immune-mediated/autoimmune conditions (e.g., Crohn's disease, SLE, aplastic anemia, myasthenia gravis, meningitis/encephalitis)
- Initiation of immunomodulatory medications (e.g., cancer immunotherapies, rituximab, azathioprine)
- Development of other neurologic conditions (e.g., intracerebral bleeds, TBI, stroke)

**OR**

**7.2 BOTH of the following:**

- Patient does meet one of the above
- Prescriber documents clinical rationale for continued use of aducanumab†

**AND**

**8** - Prescribed by or in consultation with one of the following:

- Neurologist
- Geriatric specialist

**AND**

**9** - Therapy should be discontinued permanently and the request should be denied if one or more of the following apply:

- If the patient has had  $\geq 10$  new incident microhemorrhages, regardless of clinical severity (including asymptomatic)
- If the patient had a serious event [Serious events include concern for immediate risk of death (a life-threatening event); inpatient hospitalization or prolongation of existing hospitalization due to symptoms; new persistent or significant disability/incapacity]
- If the patient has had  $\geq 3$  new incident areas of superficial siderosis, regardless of clinical severity (including asymptomatic) therapy should be discontinued permanently and the request should be denied

Notes

\*NOTE: If the patient has had  $\geq 10$  new incident microhemorrhages, regardless of clinical severity (including asymptomatic) therapy should be discontinued permanently and the request should be denied.

\*NOTE: If the patient had a serious event, therapy should be discontinued. †

\*NOTE: If the patient has had  $\geq 3$  new incident areas of superficial siderosis, regardless of clinical severity (including asymptomatic) therapy should be discontinued permanently and the request should be denied.

†Serious events include concern for immediate risk of death (a life-threatening event); inpatient hospitalization or prolongation of existing hospitalization due to symptoms; new persistent or significant disability/incapacity.

‡Requests should be evaluated case-by-case with clinical review and MD advisor.

## 2 . Background

**Appendix**

**ARIA - H (Microhemorrhages)**

		New Incident Microhemorrhages		
		Radiographic Severity		
		Mild (1 to 4)	Moderate (5 to 9)	Severe (≥10)
Clinical Symptom Severity	Asymptomatic	Continue treatment; MRI q4w until stable	Suspend treatment; MRI q4w until stable; Restart once stable	Stop Permanently
	Mild	Suspend treatment; MRI q4w until stable Restart once stable and clinical symptoms resolved		Stop Permanently
	Moderate			
	Severe			
Serious	Stop Permanently			

**ARIA - H (Superficial Siderosis)**

		New Incident Areas of Superficial Siderosis (Central Read)		
		Radiographic Severity		
		Mild (1)	Moderate (2)	Severe (≥3)
Clinical Symptom Severity	Asymptomatic	Continue treatment; MRI q4w until stable	Suspend treatment; MRI q4w until stable; Restart once stable	Stop Permanently
	Mild	Suspend treatment; MRI q4w until stable Restart once stable and clinical symptoms resolved		Stop Permanently
	Moderate			
	Severe			
Serious	Stop Permanently			

**ARIA - E**

		ARIA-E Severity on MRI (Central Read)
		Radiographic Severity

		Mild	Moderate	Severe
<b>Clinical Symptom Severity</b>	Asymptomatic	Continue treatment; MRI q4w until stable	Suspend treatment; MRI q4w until stable; Restart once stable	
	Mild	Suspend treatment; MRI q4w until stable Restart once stable and clinical symptoms resolved		
	Moderate			
	Severe			
	Serious	Stop Permanently		

### 3 . Revision History

Date	Notes
5/17/2022	Updated Medicare sections for clarification.

Adzyna (ADAMTS13, recombinant-krhn)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-143519
<b>Guideline Name</b>	Adzyna (ADAMTS13, recombinant-krhn)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Adzyna	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting ALL of the following: 1.1 Diagnosis of congenital thrombotic thrombocytopenic purpura (cTTP)	

**AND**

**1.2** Molecular genetic testing confirms mutations in the ADAMTS13 gene

**AND**

**1.3** Trial and inadequate response, contraindication or intolerance to plasma-based infusions

Product Name:Adzynma	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy	
<b>AND</b>	
2 - Trial and inadequate response, contraindication or intolerance to plasma-based infusions [B, 11]	

## 2 . Revision History

Date	Notes
2/23/2024	New program

Aemcolo

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99426
<b>Guideline Name</b>	Aemcolo
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Aemcolo	
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of travelers' diarrhea  <b>AND</b>  2 - History of failure, contraindication, or intolerance to ONE of the following:	

- Azithromycin (generic Zithromax)
- Ciprofloxacin (generic Cipro)
- Levofloxacin (generic Levaquin)
- Ofloxacin (generic Floxin)

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copy C&S Arizona to Arizona Standard

Afinitor, Torpenz (everolimus)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-150053
<b>Guideline Name</b>	Afinitor, Torpenz (everolimus)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz	
Diagnosis	Neuroendocrine tumors
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of one of the following: <ul style="list-style-type: none"><li>Neuroendocrine tumors of pancreatic origin</li></ul>	

- Neuroendocrine tumors of gastrointestinal origin
- Neuroendocrine tumors of lung origin
- Neuroendocrine tumors of thymic origin

**AND**

**2** - Disease is progressive

**AND**

**3** - One of the following:

- Disease is unresectable
- Disease is locally advanced
- Disease is metastatic

**AND**

**4** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz	
Diagnosis	Renal cell cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of renal cell cancer</p> <p><b>AND</b></p>	

**2 - One of the following:**

**2.1 Disease has relapsed**

**OR**

**2.2 BOTH of the following**

- Medically or surgically unresectable tumor
- Diagnosis of Stage IV disease

**AND**

**3 - One of the following:**

**3.1 Patient with non-clear cell histology**

**OR**

**3.2 Both of the following:**

**3.2.1 Patient with predominantly clear cell histology**

**AND**

**3.2.2 History of failure, contraindication, or intolerance to at least one prior systemic therapy [e.g., Nexavar (sorafenib), Sutent (sunitinib), Opdivo (nivolumab), Cabometyx (cabozantinib)]**

**AND**

**4 - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)**

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Renal Angiomyolipoma with Tuberous Sclerosis Complex
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of renal angiomyolipoma and tuberous sclerosis complex (TSC), not requiring immediate surgery</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)</p>	

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz	
Diagnosis	Subependymal Giant Cell Astrocytoma Associated with Tuberous Sclerosis Complex
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis (TS)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is not a candidate for curative surgical resection</p>	

**AND**

**3** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Waldenströms Macroglobulinemia or Lymphoplasmacytic Lymphoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of one of the following:

- Waldenströms macroglobulinemia
- Lymphoplasmacytic lymphoma

**AND**

**2** - One of the following:

- Disease is non-responsive to primary treatment
- Disease is progressive
- Disease has relapsed

**AND**

**3** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Breast Cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Diagnosis of breast cancer

**AND**

2 - One of the following:

2.1 Disease is recurrent

**OR**

2.2 Disease is metastatic

**AND**

3 - One of the following:

3.1 Disease is hormone receptor positive (HR+) [i.e., estrogen-receptor-positive (ER+) or progesterone-receptor-positive (PR+)]

**OR**

3.2 BOTH of the following:

- Disease is hormone receptor negative (HR-)
- Disease has clinical characteristics that predict a HR+ tumor

**AND**

**4** - Disease is human epidermal growth factor receptor 2 (HER2)-negative

**AND**

**5** - One of the following:

**5.1** Patient is a postmenopausal woman

**OR**

**5.2** BOTH of the following:

- Patient is a premenopausal woman
- Patient is being treated with ovarian ablation/suppression

**OR**

**5.3** Patient is male

**AND**

**6** - One of the following:

**6.1** Both of the following:

**6.1.1** Used in combination with Aromasin (exemestane)

**AND**

**6.1.2** One of the following:

**6.1.2.1** Disease progressed while on or within 12 months of non-steroidal aromatase inhibitor [e.g., Arimidex (anastrozole), Femara (letrozole)] therapy

**OR**

**6.1.2.2** Patient was treated with tamoxifen at any time

**OR**

**6.2** Used in combination with ONE of the following:

- Fulvestrant
- Tamoxifen

**AND**

**7** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Hodgkin Lymphoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of classical Hodgkin lymphoma

**AND**

**2** - ONE of the following:

- Disease is refractory

- Disease has relapsed

**AND**

**3** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	PEComa (perivascular epithelioid cell tumor), recurrent angiomyolipoma, lymphangiomyomatosis, or gastrointestinal stromal tumor (GIST)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - One of the following

**1.1** One of the following diagnoses:

- PEComa (perivascular epithelioid cell tumor)
- Recurrent angiomyolipoma
- Lymphangiomyomatosis

**OR**

**1.2** All of the following:

**1.2.1** Diagnosis of Gastrointestinal Stromal Tumor (GIST)

**AND**

**1.2.2** Disease has progressed after single agent therapy with ONE of the following:

- Gleevec (imatinib)
- Sutent (sunitinib)
- Stivarga (regorafenib)

**AND**

**1.2.3** Used in combination with ONE of the following:

- Gleevec (imatinib)
- Sutent (sunitinib)
- Stivarga (regorafenib)

**AND**

**2** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz	
Diagnosis	Thymic Carcinoma or Thymoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - One of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of thymic carcinoma</li> <li>• Diagnosis of thymoma</li> </ul> <p><b>AND</b></p> <p><b>2</b> - ONE of the following:</p>	

**2.1** History of failure, contraindication, or intolerance to at least one prior first-line chemotherapy regimen

**OR**

**2.2** Patient has extrathoracic metastatic disease

**AND**

**3** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Follicular carcinoma, Hürthle cell carcinoma, or papillary carcinoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of ONE of the following:

- Follicular carcinoma
- Hürthle cell carcinoma
- Papillary carcinoma

**AND**

**2** - ONE of the following:

- Unresectable locoregional recurrent disease
- Persistent disease
- Metastatic disease

**AND**

**3** - ONE of the following:

- Patient has symptomatic disease
- Patient has progressive disease

**AND**

**4** - Disease is refractory to radioactive iodine treatment

**AND**

**5** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Meningioma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of meningioma

**AND**

**2** - Disease is recurrent or progressive

**AND**

**3** - Surgery and/or radiation is not possible

**AND**

**4** - Used in combination with bevacizumab (e.g., Avastin, Mvasi, Zirabev)

**AND**

**5** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Endometrial Carcinoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of endometrial carcinoma

**AND**

**2** - Used in combination with letrozole

**AND**

**3** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	Tuberous Sclerosis Complex associated Partial-Onset Seizures
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of tuberous sclerosis complex associated partial-onset seizures

**AND**

2 - Used as adjunctive therapy

**AND**

**3** - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name: Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz

Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Use supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.

**AND**

2 - For Brand Afinitor/Afinitor Disperz or Torpenz requests only, history of failure, intolerance, or contraindication to generic everolimus tablet or everolimus tablet for oral suspension (Applies to requests for Brand Afinitor/Afinitor Disperz or Torpenz ONLY)

Product Name:Brand Afinitor, Generic everolimus tablet, Brand Afinitor Disperz, Generic everolimus tablet for oral suspension, Torpenz	
Diagnosis	All indications listed above
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on therapy	

**2 . Revision History**

Date	Notes
7/25/2024	Added GPIs for Torpenz (NP), step though generic everolimus. Consolidated all reauth sections into one criteria box for all covered indications.

Afrezza

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99427
<b>Guideline Name</b>	Afrezza
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Afrezza	
Diagnosis	Type 1 or Type 2 diabetes mellitus
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Diagnosis of type 1 diabetes mellitus and used in combination with a basal insulin or continuous insulin pump	

**OR**

**1.2** Diagnosis of type 2 diabetes mellitus

**AND**

**2** - Patient is unable to self-inject medications (e.g. Humalog, Lantus, Levemir) due to ONE of the following:

- Physical impairment
- Visual impairment
- Lipohypertrophy
- Documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-5 for specific phobia diagnostic criteria)

**AND**

**3** - Forced Expiratory Volume (FEV1) within the last 60 days is greater than or equal to 70% of expected normal as determined by the physician

**AND**

**4** - Afrezza will not be approved in patients with ONE of the following:

- Who smoke cigarettes
- Who recently quit smoking (within the past 6 months)
- With chronic lung disease (e.g. asthma, chronic obstructive pulmonary disease)

Product Name: Afrezza	
Diagnosis	Type 1 or Type 2 diabetes mellitus
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Repeat pulmonary function test confirms that patient has NOT experienced a decline of 20% or more in Forced Expiratory Volume (FEV1)

**AND**

2 - Patient continues to be unable to self-inject short-acting insulin due to ONE of the following:

- Physical impairment
- Visual impairment
- Lipohypertrophy
- Documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-5 for specific phobia diagnostic criteria)

**AND**

3 - Patient continues to not smoke cigarettes

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copy C&S Arizona to Arizona Standard

Agamree (vamorolone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144824
<b>Guideline Name</b>	Agamree (vamorolone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Agamree	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of Duchenne muscular dystrophy (DMD)	

**AND**

**2** - Patient is 2 years of age or older

**AND**

**3** - Patient has received genetic testing for a mutation of the dystrophin gene

**AND**

**4** - Submission of medical records (e.g., chart notes) documenting one of the following:

**4.1** Patient has a confirmed mutation of the dystrophin gene

**OR**

**4.2** Muscle biopsy confirmed an absence of dystrophin protein

**AND**

**5** - Submission of medical records (e.g., chart notes) or paid claims confirming patient has had a trial and failure or intolerance to prednisone or prednisolone given at a dose of 0.75 mg/kg/day or 10 mg/kg/weekend

**AND**

**6** - Prescribed by or in consultation with a neurologist who has experience treating children

**AND**

**7** - One of the following:

**7.1** For patients less than or equal to 50kg, dose will not exceed 6mg/kg of body weight once daily

**OR**

**7.2** For patients greater than 50kg, dose will not exceed 300mg/day

Product Name:Agamree	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting patient has experienced a benefit from therapy (e.g., improvement in preservation of muscle strength)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p><b>2.1</b> For patients less than or equal to 50kg, dose will not exceed 6mg/kg of body weight once daily</p> <p style="text-align: center;"><b>OR</b></p> <p><b>2.2</b> For patients greater than 50kg, dose will not exceed 300mg/day</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Submission of medical records (e.g., chart notes) or paid claims confirming patient has had a trial and failure or intolerance to prednisone or prednisolone given at a dose of 0.75 mg/kg/day or 10 mg/kg/weekend</p>	

## 2 . Revision History

Date	Notes
3/25/2024	New program

Airsupra (albuterol-budesonide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-133833
<b>Guideline Name</b>	Airsupra (albuterol-budesonide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Airsupra	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of asthma  <b>AND</b>	

2 - Patient is 18 years of age or older

**AND**

3 - Trial and failure, contraindication, or intolerance to treatment with ALL of the following preferred products:

- Advair Diskus (brand) or Advair HFA
- Dulera
- Brand Symbicort

**AND**

4 - Trial, failure, contraindication or intolerance to BOTH of the following:

- Generic albuterol inhaler
- A preferred inhaled corticosteroid (e.g, Pulmicort, Brand Flovent, Asmanex)

**AND**

5 - Physician has provided rationale for needing to use fixed-dose combination therapy with Airsupra instead of taking individual products in combination (i.e., albuterol inhaler and Pulmicort)

Product Name: Airsupra	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	

## 2 . Revision History

Date	Notes
9/28/2023	New program

Aldurazyme - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99428
<b>Guideline Name</b>	Aldurazyme - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Aldurazyme	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Confirmed diagnosis of Hurler and Hurler-Scheie forms of Mucopolysaccharidosis I (MPS I)	

**OR**

**1.2** Both the following:

**1.2.1** Confirmed diagnosis of Scheie form of Mucopolysaccharidosis I (MPS I)

**AND**

**1.2.2** Have moderate to severe symptoms

## **2 . Revision History**

Date	Notes
3/10/2021	Bulk Copy C&S Arizona to Arizona Standard

Alinia

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99429
<b>Guideline Name</b>	Alinia
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Alinia, generic nitazoxanide	
Diagnosis	Diarrhea caused by Giardia lamblia
Approval Length	3 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of giardiasis  <b>AND</b>	

2 - History of failure, contraindication, or intolerance to metronidazole

Product Name:Brand Alinia, generic nitazoxanide	
Diagnosis	Diarrhea caused by Cryptosporidium parvum
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of cryptosporidiosis	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copy C&S Arizona to Arizona Standard

Alpha Interferons - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-105169
<b>Guideline Name</b>	Alpha Interferons - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

## 1 . Criteria

Product Name: Intron A	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of hairy cell leukemia  <b>OR</b>	

**2** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of condylomata acuminata (genital or perianal)

**OR**

**3** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of AIDS-related Kaposi's sarcoma

**OR**

**4** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of leptomeningeal metastases

**OR**

**5** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of meningiomas

**OR**

**6** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of kidney cancer

**OR**

**7** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting treatment of myeloproliferative neoplasms (MPNs) such as essential thrombocythemia (ET), polycythemia vera (PV), or primary myelofibrosis (PM)

**OR**

**8** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of follicular lymphoma

**OR**

**9** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of adult T-cell leukemia, lymphoma

**OR**

**10** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of mycosis fungoides, Sézary syndrome

**OR**

**11** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of desmoid tumors/aggressive fibromatosis

**OR**

**12** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of giant cell tumor of the bone

**OR**

**13** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of malignant melanoma

Product Name:Alferon N	
Approval Length	8 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting	

treatment of refractory or recurring external condylomata acuminata (genital or venereal warts) due to the human papillomavirus (HPV) infection

## 2 . Revision History

Date	Notes
3/24/2022	Removed Sylatron from guideline, Added Submission of Medical Records

Alpha-1 Proteinase Inhibitors

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-138189
<b>Guideline Name</b>	Alpha-1 Proteinase Inhibitors
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	2/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Aralast NP, Glassia, Prolastin-C, Prolastin-C liquid, Zemaira	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has clinically evident emphysema</p> <p style="text-align: center;"><b>AND</b></p>	

**2** - Submission of medical records (e.g., chart notes) documenting a diagnosis of severe congenital deficiency of Alpha1- proteinase inhibitor (alpha1 antitrypsin deficiency)

**AND**

**3** - For Glassia requests ONLY: Paid claims or submission of medical records (e.g., chart notes) (document drug, duration, and date of use) confirming trial and failure, contraindication or intolerance to ALL of the following:

- Aralast NP
- Prolastin-C or Prolastin-C liquid
- Zemaira

## **2 . Revision History**

Date	Notes
1/23/2024	Added Glassia (NP), Prolastin-C, and Zemaira as targets.

Alyftrek (vanzacaftor/tezacaftor/deutivacaftor)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-226189
<b>Guideline Name</b>	Alyftrek (vanzacaftor/tezacaftor/deutivacaftor)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Alyftrek	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of cystic fibrosis (CF)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting presence of at least one of the following mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene as detected by a U.S. Food and Drug Administration (FDA)-cleared cystic fibrosis mutation test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA):

- F508del mutation
- A mutation in the CFTR gene that is responsive based on clinical, in vitro, or extrapolated data<sup>^</sup>

**AND**

**3** - Patient is 6 years of age or older

**AND**

**4** - Prescribed by or in consultation with a specialist affiliated with a CF care center

**AND**

**5** - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure or inadequate response to Trikafta (elexacaftor/tezacaftor/ivacaftor)

Notes	<sup>^</sup> Please consult Background section for table of CFTR gene mutations responsive to Alyftrek.
-------	---

Product Name:Alyftrek	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) documenting that patient demonstrates positive clinical response to therapy (e.g., stable or improved lung function [percent predicted forced expiratory volume in one second {PPFEV1}], decreased number of pulmonary exacerbations)

**AND**

2 - Prescribed by or in consultation with a specialist affiliated with a CF care center

## 2 . Background

### Clinical Practice Guidelines

#### List of CFTR Gene Mutations that are Responsive to Alyftrek [1]

^Intent of table is to provide a quick reference; PA team members should still review at point of request for clinical appropriateness as off label support continuously evolves.

[Last Reviewed: 2/17/25]

#### Based on Clinical Data\*

A455E	G551D	L1077P†	R352Q	S549N	V754M	
D1152H	G85E†	L206W	R75Q	S549R	W1098C†	
F508del†	H1054D	M1101K†	S1159F	S945L	W1282R	
G1244E	I336K	R1066H	S1251N	V562I	Y563N†	

#### Based on in vitro Data‡

1507_1515del9	E116Q	G424S	I556V	P140S	R334L	T1053I
2183A→G	E193K	G463V	I601F	P205S	R334Q	T1086I
3141del9	E292K	G480C	I618T	P499A	R347H	T1246I
3195del6	E403D	G480S	I807M	P5L	R347L	T1299I
3199del6	E474K	G551A	I980K	P574H	R347P	T338I
546insCTA	E56K	G551S	K1060T	P67L	R352W	T351I
A1006E	E588V	G576A	K162E	P750L	R516G	T604I
A1067P	E60K	G576A;R668C§	K464E	P99L	R516S	V1153E
A1067T	E822K	G622D	L1011S	Q1100P	R553Q	V1240G

A107G	E92K	G628R	L102R	Q1291R	R555G	V1293 G
A120T	F1016S	G91R	L1065P	Q1313K	R560S	V201M
A234D	F1052V	G970D	L1324P	Q237E	R560T	V232D
A309D	F1074L	G970S	L1335P	Q237H	R668C	V392G
A349V	F1099L	H1085P	L137P	Q359R	R709Q	V456A
A46D	F1107L	H1085R	L1480P	Q372H	R74Q	V456F
A554E	F191V	H1375P	L15P	Q452P	R74W	V520F
A559T	F200I	H139R	L165S	Q493R	R74W;D1270N§	V603F
A559V	F311del	H199R	L320V	Q552P	R74W;V201M§	W361R
A561E	F311L	H199Y	L333F	Q98R	R74W;V201M;D 1270N§	Y1014C
A613T	F508C	H609R	L333H	R1048G	R75L	Y1032C
A62P	F508C;S12 51N§	H620P	L346P	R1066C	R751L	Y109N
A72D	F575Y	H620Q	L441P	R1066L	R792G	Y161D
C491R	F587I	H939R	L453S	R1066M	R933G	Y161S
D110E	G1047R	H939R;H9 49L	L619S	R1070Q	S1045Y	Y301C
D110H	G1061R	I1027T	L967S	R1070W	S108F	Y569C
D1270N	G1069R	I105N	L997F	R1162L	S1118F	Y913C
D1445N	G1123R	I1139V	M1101R	R117C	S1159P	
D192G	G1247R	I1234Vdel 6aa	M1137V	R117C;G576A; R668C	S1235R	
D443Y	G1249R	I125T	M150K	R117G	S1255P	
D443Y;G576A; R668C§	G126D	I1269N	M152V	R117H	S13F	
D513G	G1349D	I331N	M265R	R117L	S341P	
D565G	G149R	I1366N	M952I	R117P	S364P	
D579G	G178E	I1398S	M952T	R1283M	S492F	
D614G	G178R	I148N	N1088D	R1283S	S549I	
D836Y	G194R	I148T	N1303I	R170H	S589N	
D924N	G194V	I175V	N1303K§	R258G	S737F	
D979V	G27E	I502T	N186K	R297Q	S912L	
D993Y	G27R	I506L	N187K	R31C	S977F	
E116K	G314E	I506T	N418S	R31L	T1036N	
<b>Based on Extrapolation¶</b>						
1341G→A	2789+2ins A	3041- 15T→G	3849+10kb C→T	3850-3T→G	5T;TG13	711+3A →G

1898+3A→G	2789+5G→A	3272-26A→G	3849+4A→G	4005+2T→C	621+3A→G	E831X
2752-26A→G	296+28A→G	3600G→A	3849+40A→G	5T;TG12		
<p>* Clinical data is obtained from Trials 1 and 2.</p> <p>† This mutation is also predicted to be responsive by FRT assay with ALYFTREK.</p> <p>‡ The N1303K mutation is predicted to be responsive only by HBE assay. All other mutations predicted to be responsive with in vitro data are supported by FRT assay.</p> <p>§ Complex/compound mutations where a single allele of the <i>CFTR</i> gene has multiple mutations; these exist independent of the presence of mutations on the other allele.</p> <p>¶ Efficacy is extrapolated to certain non-canonical splice mutations because clinical trials in all mutations in this subgroup are infeasible and these mutations are not amenable to interrogation by FRT system.</p>						

### 3 . Revision History

Date	Notes
3/26/2025	New program

Alzheimer's Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-109871
<b>Guideline Name</b>	Alzheimer's Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/27/2022
-----------------	-----------

## 1 . Criteria

Product Name:Brand Aricept, generic donepezil, Brand Namenda/Namenda XR, generic memantine/memantine XR, Brand Razadyne, generic galantamine hydrobromide, Brand Razadyne ER, generic galantamine ER	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of dementia of the Alzheimer's type	

Product Name:Brand Exelon, generic rivastigmine
---

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of dementia of the Alzheimer's type</p> <p style="text-align: center;"><b>OR</b></p> <p>2 - Diagnosis of dementia associated with Parkinson's disease</p>	

Product Name:Adlarity	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of dementia of the Alzheimer's type</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p><b>2.1</b> History of failure, contraindication or intolerance to ALL of the following preferred drugs* (verified via paid pharmacy claims):</p> <ul style="list-style-type: none"> <li>• generic donepezil</li> <li>• generic galantamine IR/ER</li> <li>• generic memantine</li> <li>• generic oral rivastigmine</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>2.2</b> Both of the following:</p>	

**2.2.1** History of failure, contraindication or intolerance to generic rivastigmine patch\* (verified via paid pharmacy claims)

**AND**

**2.2.2** Patient is unable to swallow oral formulations or has documented swallowing difficulties

Notes

\*PA may be required

## 2 . Revision History

Date	Notes
7/27/2022	Added XR formulations of Namenda/memantine to product name section. No change to criteria.

Amvuttra (vutrisiran)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-240226
<b>Guideline Name</b>	Amvuttra (vutrisiran)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Amvuttra	
Diagnosis	Hereditary transthyretin-mediated amyloidosis with polyneuropathy (hATTR-PN)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of documentation (e.g., chart notes) confirming diagnosis of hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) with polyneuropathy	

**AND**

**2** - Patient has a transthyretin (TTR) mutation (e.g., V30M)

**AND**

**3** - Two of the following:

- Patient has a baseline polyneuropathy disability (PND) score less than or equal to IIIb
- Patient has a baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2
- Patient has a baseline neuropathy impairment score (NIS) greater than or equal to 5 and less than or equal to 130
- Patient has a baseline Karnofsky Performance Status score greater than or equal to 60%

**AND**

**4** - Presence of clinical signs and symptoms of the disease (e.g., peripheral/autonomic neuropathy, walking ability, quality of life)

**AND**

**5** - Patient has not had a liver transplant

**AND**

**6** - Requested drug is not used in combination with a TTR silencer (e.g., Tegsedi) or a TTR stabilizer (e.g., Vyndaqel)

**AND**

**7** - Prescribed by or in consultation with a neurologist

Product Name: Amvuttra

Diagnosis	Hereditary transthyretin-mediated amyloidosis with polyneuropathy (hATTR-PN)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of documentation (e.g., chart notes) confirming positive clinical response to therapy as evidenced by an improvement in clinical signs and symptoms from baseline (e.g., neuropathy, quality of life, gait speed, nutritional status, decrease in serum TTR level)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Two of the following:</p> <ul style="list-style-type: none"> <li>• Patient continues to have a polyneuropathy disability (PND) score less than or equal to IIIb</li> <li>• Patient continues to have a familial amyloidotic polyneuropathy (FAP) stage of 1 or 2</li> <li>• Patient continues to have a neuropathy impairment score (NIS) greater than or equal to 5 and less than or equal to 130</li> <li>• Patient continues to have a Karnofsky Performance Status score greater than or equal to 60%</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Requested drug is not used in combination with a TTR silencer (e.g., Tegsedi) or a TTR stabilizer (e.g., Vyndaqel)</p>	

Product Name: Amvuttra	
Diagnosis	Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Submission of documentation (e.g., chart notes) confirming diagnosis of transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM)

**AND**

**2** - One of the following:

**2.1** Patient has a transthyretin (TTR) mutation (e.g., V122I)

**OR**

**2.2** Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits

**OR**

**2.3** Both of the following:

- Cardiac magnetic resonance imaging or scintigraphy scan suggestive of amyloidosis
- Absence of light-chain amyloidosis

**AND**

**3** - Patient has New York Heart Association (NYHA) Functional Class I, II, or III heart failure

**AND**

**4** - Requested drug is not used in combination with a TTR silencer (e.g., Onpattro)

**AND**

5 - Prescribed by or in consultation with a cardiologist

Product Name:Amvuttra	
Diagnosis	Transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure	
<b>AND</b>	
2 - Requested drug is not used in combination with a TTR silencer (e.g., Onpattro)	

## 2 . Revision History

Date	Notes
4/24/2025	Added criteria for new ATTR-CM indication. Added criterion regarding concomitant use of TTR silencer/stabilizer to hATTR-PN sections.

Anthelmintics - Arizona

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-99431
<b>Guideline Name</b>	Anthelmintics - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Brand Albenza, generic albendazole	
Diagnosis	See Note section*
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Enterobius vermicularis (pinworm)  <b>OR</b>	

**2 - Diagnosis of Hydatid Disease [Echinococcosis (Tapeworm)]**

**OR**

**3 - Diagnosis of Ancylostoma/Necatoriasis (Hookworm)**

**OR**

**4 - Diagnosis of Ascariasis (Roundworm)**

**OR**

**5 - Diagnosis of Mansonella perstans (Filariasis)**

**OR**

**6 - Diagnosis of Toxocariasis (Roundworm)**

**OR**

**7 - Diagnosis of Trichinellosis**

**OR**

**8 - Diagnosis of Trichuriasis (Whipworm)**

**OR**

**9 - Diagnosis of Capillariasis**

Notes

\* Enterobius vermicularis (pinworm), Hydatid Disease [Echinococcosis (Tapeworm)]  
Ancylostoma/Necatoriasis (Hookworm), Ascariasis (Roundworm), Ma

	nsonella perstans (Filariasis), Toxocariasis (Roundworm), Trichinellois, Trichuriasis (Whipworm), Capillariasis
--	---

Product Name:Brand Albenza, generic albendazole	
Diagnosis	Neurocysticercosis
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of neurocysticercosis</p>	

Product Name:Brand Stromectol, generic ivermectin	
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of intestinal strongyloidiasis due to the nematode parasite Strongyloides stercoralis</p> <p style="text-align: center;"><b>OR</b></p> <p>2 - Diagnosis of onchocerciasis due to the nematode parasite Onchocerca volvulus</p>	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copy C&S Arizona to Arizona Standard

Anti-Parkinson's Agents (Crexont, Duopa, Vyalev)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-163934
<b>Guideline Name</b>	Anti-Parkinson's Agents (Crexont, Duopa, Vyalev)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Crexont	
Diagnosis	Parkinson's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following diagnoses: <ul style="list-style-type: none"><li>Parkinson's disease</li><li>Post-encephalitic parkinsonism</li></ul>	

- Symptomatic parkinsonism that may follow carbon monoxide and/or manganese intoxication

**AND**

**2** - History of failure (after a minimum 30 day trial) or adverse reaction to one of the following:

- Generic carbidopa-levodopa immediate release
- Generic carbidopa-levodopa extended release

Product Name: Duopa	
Diagnosis	Parkinson's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of advanced Parkinson's disease</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is levodopa-responsive</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Patient experiences disabling "off" periods for a minimum of 3 hours per day</p> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Disabling "off" periods occur despite therapy with BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Oral levodopa-carbidopa</li> </ul>	

- One drug from a different class of anti-Parkinson's disease therapy (e.g., COMT [catechol-O-methyltransferase] inhibitor [entacapone, tolcapone], MAO-B [monoamine oxidase-B] inhibitor [selegiline, rasagiline], dopamine agonist [pramipexole, ropinirole])

**AND**

**5** - Has undergone or has planned placement of a procedurally-placed tube

**AND**

**6** - Prescribed by or in consultation with a neurologist

Product Name:Vyalev	
Diagnosis	Parkinson's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of advanced Parkinson's disease</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is levodopa-responsive</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Patient experiences disabling "off" periods for a minimum of 2.5 hours per day</p> <p style="text-align: center;"><b>AND</b></p>	

4 - Disabling "off" periods occur despite therapy with BOTH of the following:

- Oral levodopa-carbidopa
- One drug from a different class of anti-Parkinson's disease therapy (e.g., COMT [catechol-O-methyltransferase] inhibitor [entacapone, tolcapone], MAO-B [monoamine oxidase-B] inhibitor [selegiline, rasagiline], dopamine agonist [pramipexole, ropinirole])

**AND**

5 - Prescribed by or in consultation with a neurologist

Product Name: Crexont, Duopa, Vyalev	
Diagnosis	Parkinson's disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	

## 2 . Revision History

Date	Notes
1/31/2025	Added reauth criteria for Crexont

Anticonvulsants - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164382
<b>Guideline Name</b>	Anticonvulsants - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name: PREFERRED: generic lacosamide, Xcopri; NON-PREFERRED: Aptiom, Briviact, Brand Vimpat	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:  1.1 All of the following:  1.1.1 Diagnosis of partial-onset seizures	

**AND**

**1.1.2** History of greater than or equal to 8 week trial of at least TWO of the following (any release formulation qualifies)\*: (APPLIES TO APTIOM, BRIVIACT, AND BRAND VIMPAT ONLY)

- Carbamazepine
- Divalproex
- Gabapentin
- Fycompa
- generic lacosamide
- Lamotrigine
- Levetiracetam
- Oxcarbazepine
- Phenytoin
- Pregabalin
- Topiramate
- Valproic acid
- Xcopri
- Zonisamide

**AND**

**1.1.3** One of the following: (APPLIES TO APTIOM, BRIVIACT, AND BRAND VIMPAT ONLY)

**1.1.3.1** Both of the following:

- Documented history of persisting seizures after titration to the highest tolerated dose with each medication trial of preferred formulary alternatives
- Lack of compliance as a reason for treatment failure has been ruled out

**OR**

**1.1.3.2** Both of the following:

- Documentation of failure of preferred formulary alternatives due to intolerable side effects
- Reasonable efforts were made to minimize the side effect (e.g. change timing of dosing, divide dose out for more frequent but smaller doses, etc.)

**AND**

**1.1.4** Trial and failure, contraindication, or intolerance to generic lacosamide (APPLIES TO BRAND VIMPAT ONLY)

**OR**

**1.2** For continuation of prior therapy for a seizure disorder

Notes

\*Preferred Drugs may require PA

Product Name: Motpoly XR

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:

**1.1** Both of the following:

**1.1.1** Diagnosis of one of the following:

- partial-onset seizures
- primary generalized tonic-clonic seizures

**AND**

**1.1.2** Patient weighs at least 50 kg

**OR**

**1.2** For continuation of prior therapy for a seizure disorder

Product Name:Fycompa	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:</p> <p>1.1 Diagnosis of partial-onset or primary generalized tonic-clonic seizures</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 For continuation of prior therapy for a seizure disorder</p>	

Product Name:Epidiolex	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:</p> <p>1.1 Diagnosis of seizures associated with Dravet syndrome</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 Diagnosis of seizures associated with Lennox-Gastaut syndrome</p> <p style="text-align: center;"><b>OR</b></p> <p>1.3 Diagnosis of seizures associated with tuberous sclerosis complex (TSC)</p>	

**OR**

**1.4** For continuation of prior therapy for a seizure disorder

Notes

\*Drug may require PA

Product Name:Diacomit

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of Dravet syndrome and currently taking clobazam

**OR**

**2** - For continuation of prior therapy for a seizure disorder

Product Name:Fintepla

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ALL of the following:

**1.1** Diagnosis of seizures associated with Dravet syndrome

**AND**

**1.2** History of greater than or equal to 8-week trial of at least TWO of the following (any release formulation qualifies)\*:

- Divalproex (e.g., generic Depakote)
- Epidiolex
- Levetiracetam (e.g., generic Keppra)
- Topiramate (e.g., generic Topamax)
- Valproic acid (e.g., generic Depakene)
- Zonisamide (generic Zonegran)

**AND**

**1.3** ONE of the following:

**1.3.1** BOTH of the following:

**1.3.1.1** Documented history of persisting seizures after titration to the highest tolerated dose with each medication trial of preferred formulary alternatives

**AND**

**1.3.1.2** Lack of compliance as a reason for treatment failure has been ruled out

**OR**

**1.3.2** BOTH of the following:

**1.3.2.1** Documentation of failure of preferred formulary alternatives due to intolerable side effects

**AND**

**1.3.2.2** Reasonable efforts were made to minimize the side effect (e.g., change timing of dosing, divide dose out for more frequent but smaller doses, etc.)

**OR**

**2** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ALL of the following:

**2.1** Diagnosis of seizures associated with Lennox-Gastaut syndrome

**AND**

**2.2** History of greater than or equal to 8 week trial, contraindication or intolerance of at least TWO of the following (any release formulation qualifies)\*:

- Banzel (rufinamide)
- Clobazam
- Divalproex
- Epidiolex
- Felbamate
- Lamotrigine
- Topiramate
- Valproic Acid

**AND**

**2.3** ONE of the following:

**2.3.1** BOTH of the following:

- Documented history of persisting seizures after titration to the highest tolerated dose with each medication trial of preferred formulary alternatives
- Lack of compliance as a reason for treatment failure has been ruled out

**OR**

**2.3.2** BOTH of the following:

- Documentation of failure of preferred formulary alternatives due to intolerable side effects
- Lack of compliance as a reason for treatment failure has been ruled out

**OR**

**3** - For continuation of prior therapy for a seizure disorder

Notes	*Drug may require PA
-------	----------------------

Product Name:PREFERRED: Brand Banzel tablets and suspension, generic rufinamide tablets; NON-PREFERRED: generic rufinamide solution	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Submission of medical records (e.g., chart notes, lab work, imaging) documenting both of the following:</p> <p>1.1.1 Diagnosis of seizures associated with Lennox-Gastaut syndrome</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.2 Trial and failure, contraindication, or intolerance to Brand Banzel suspension (APPLIES TO GENERIC RUFINAMIDE SUSPENSION ONLY)</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 For continuation of prior therapy for a seizure disorder</p>	

Product Name:PREFERRED: generic clobazam; NON-PREFERRED: Brand Onfi	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting one of the following:</p> <p>1.1 Both of the following:</p>	

- Diagnosis of seizures associated with Lennox-Gastaut syndrome
- Trial and failure, contraindication, or intolerance to generic clobazam (APPLIES TO BRAND ONFI ONLY)

**OR**

**1.2** All of the following:

- Diagnosis of Dravet syndrome
- Patient is currently taking Diacomit
- Trial and failure, contraindication, or intolerance to generic clobazam (APPLIES TO BRAND ONFI ONLY)

**OR**

**2** - For continuation of prior therapy for a seizure disorder

Product Name: Sympazan	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:</p> <p><b>1.1</b> ALL of the following:</p> <p><b>1.1.1</b> Diagnosis of seizures associated with Lennox-Gastaut syndrome (LGS)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.1.2</b> BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Used as adjunctive therapy (defined as accessory treatment used in combination to enhance primary treatment.)</li> </ul>	

- Not used as primary treatment

**AND**

**1.1.3** History of greater than or equal to 8 week trial, contraindication or intolerance of at least TWO of the following (any release formulation qualifies)\*:

- Brand Banzel suspension/tablets or runfinamide tablets
- Divalproex
- Felbamate
- Lamotrigine
- Topiramate
- Valproic acid

**AND**

**1.1.4** Prescriber provides a reason or special circumstance the patient cannot use generic clobazam tablets or suspension

**OR**

**1.2** ALL of the following:

**1.2.1** Diagnosis of refractory partial onset seizures (four or more uncontrolled seizures per month after an adequate trial of at least two antiepileptic drugs)

**AND**

**1.2.2** BOTH of the following:

- Used as adjunctive therapy (defined as accessory treatment used in combination to enhance primary treatment.)
- Not used as primary treatment

**AND**

**1.2.3** History of greater than or equal to 8 week trial of at least TWO of the following (any release formulation qualifies)\*:

- Carbamazepine
- Divalproex
- Fycompa
- Gabapentin
- Lacosamide
- Lamotrigine
- Levetiracetam
- Oxcarbazepine
- Phenytoin
- Pregabalin
- Topiramate
- Valproic acid
- Xcopri
- Zonisamide

**AND**

**1.2.4** Prescriber provides a reason or special circumstance the patient cannot use generic clobazam tablets or suspension

**OR**

**1.3** ALL of the following:

**1.3.1** Diagnosis of Dravet syndrome

**AND**

**1.3.2** Patient is currently taking Diacomit

**AND**

**1.3.3** Prescriber provides a reason or special circumstance the patient cannot use generic clobazam tablets or suspension

**OR**

**1.4** For continuation of prior therapy for a seizure disorder

Notes	*Drug may require PA
-------	----------------------

Product Name: PREFERRED: generic tiagabine; NON-PREFERRED: Brand Gabitril

Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:

**1.1** All of the following:

**1.1.1** Diagnosis of partial-onset seizures

**AND**

**1.1.2** Used as adjunctive therapy (defined as accessory treatment used in combination to enhance primary treatment)

**AND**

**1.1.3** Not used as primary treatment

**AND**

**1.1.4** Trial and failure, contraindication, or intolerance to generic tiagabine (APPLIES TO BRAND GABITRIL ONLY)

**OR**

**1.2** For continuation of prior therapy for a seizure disorder

Product Name: Brand Sabril Tablets, generic vigabatrin tablets

Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:

1.1 All of the following:

1.1.1 Diagnosis of complex partial seizures

**AND**

1.1.2 Used as adjunctive therapy (defined as accessory treatment used in combination to enhance primary treatment)

**AND**

1.1.3 Not used as primary treatment

**AND**

1.1.4 History of greater than or equal to 8 week trial of at least TWO of the following (any release formulation qualifies)\*:

- Carbamazepine
- Divalproex
- Fycompa
- Gabapentin
- Lacosamide
- Lamotrigine
- Levetiracetam
- Oxcarbazepine
- Phenytoin
- Pregabalin
- Topiramate
- Valproic acid
- Xcopri

- Zonisamide

**OR**

**1.2** For continuation of prior therapy for a seizure disorder

Notes	*Drug may require PA
-------	----------------------

Product Name: Brand Sabril Oral Solution, generic vigabatrin oral solution, generic vigadrone oral solution, Vigafyde

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of infantile spasms

**OR**

2 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting all of the following:

**2.1** Diagnosis of complex partial seizures

**AND**

**2.2** Used as adjunctive therapy (defined as accessory treatment used in combination to enhance primary treatment)

**AND**

**2.3** Not used as primary treatment

**AND**

**2.4** History of greater than or equal to 8 week trial of at least TWO of the following (any release formulation qualifies)\*:

- Carbamazepine
- Divalproex
- Fycompa
- Gabapentin
- Lacosamide
- Lamotrigine
- Levetiracetam
- Oxcarbazepine
- Phenytoin
- Pregabalin
- Topiramate
- Valproic acid
- Xcopri
- Zonisamide

**OR**

**3** - For continuation of prior therapy for a seizure disorder

Notes

\*Drug may require PA

Product Name: PREFERRED: Brand Trokendi XR; NON-PREFERRED: generic topiramate ER, Brand Qudexy XR, generic topiramate ER sprinkle

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:

**1.1** All of the following:

**1.1.1** Diagnosis of partial-onset seizures

**AND**

**1.1.2** Trial and failure, contraindication, or intolerance to generic topiramate immediate-release (IR) tablet or topiramate IR sprinkle capsule (APPLIES TO GENERIC TOPIRIMATE ER, BRAND QUDEXY XR, AND GENERIC TOPIRIMATE ER SPRINKLE ONLY)

**AND**

**1.1.3** Trial and failure, contraindication, or intolerance to Brand Trokendi XR (APPLIES TO GENERIC TOPIRIMATE ER, BRAND QUDEXY XR, AND GENERIC TOPIRIMATE ER SPRINKLE ONLY)

**OR**

**1.2** For continuation of prior therapy for a seizure disorder

Product Name: Gabarone	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p><b>1.1</b> All of the following:</p> <ul style="list-style-type: none"><li>• Patient is 3 years of age or older</li><li>• Submission of medical records (e.g., chart notes) documenting diagnosis of partial onset seizures</li><li>• Trial and failure or intolerance to generic immediate-release gabapentin (generic for Neurontin)</li></ul> <p><b>OR</b></p> <p><b>1.2</b> All of the following</p>	

- Patient is 18 years of age or older
- Submission of medical records (e.g., chart notes) documenting diagnosis of postherpetic neuralgia (PHN)
- Trial and failure or intolerance to generic immediate-release gabapentin (generic for Neurontin)
- Trial and failure or intolerance to Brand Gralise\*

Notes

\*Drug may require PA

## 2 . Revision History

Date	Notes
1/30/2025	Added criteria for Gabarone (NP).

Antidepressants - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269205
<b>Guideline Name</b>	Antidepressants - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:generic citalopram oral solution, generic fluoxetine oral solution, generic sertraline oral conc for solution	
Diagnosis	Requests for Patients greater than 12 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - The member is unable to swallow the oral tablet/capsule	

Product Name:Amitriptyline, amoxapine, bupropion tabs/SR tabs/XL tabs (150 and 300mg), citalopram tabs/oral soln, clomipramine, desipramine, doxepin caps/oral conc for solution, duloxetine capsules (20, 30, 60mg), escitalopram, fluoxetine caps/oral soln, fluvoxamine IR, generic mirtazapine tabs/ODT, imipramine tabs/caps, nortriptyline caps/oral soln, paroxetine tabs, protriptyline, sertraline tabs/oral soln, trazodone, trimipramine, venlafaxine tabs/ER capsules	
Diagnosis	PREFERRED DRUG Requests for patient 6 years of age or younger
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - The patient is unresponsive to other treatment modalities, unless contraindicated (i.e. other medications or behavioral modification attempted)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - The physician attests that the requested medication is medically necessary. (Document rationale for use)</p>	
Notes	Drug may require PA

Product Name:Brand Anafranil, Aplenzin, Auvelity, Brand Celexa, generic citalopram capsules, Brand Cymbalta, generic duloxetine 40mg caps, Drizalma , Brand Effexor XR, generic venlafaxine ER tabs, Emsam, Fetzima, fluvoxamine ER, Brand Lexapro, maprotiline, Marplan, Brand Nardil, generic phenelzine, nefazodone, Brand Norpramin, Brand Pamelor caps/oral soln, Brand Parnate, generic tranlycypromine, Brand Paxil, generic paroxetine capsules, Brand Paxil susp, generic paroxetine suspension, Brand Paxil CR, generic paroxetine ER, Pexeva, Brand Pristiq, generic desvenlafaxine ER, Brand Prozac, generic fluoxetine tablets, Brand Remeron SLTB, Brand Remeron, Trintellix, Viibryd, Brand Wellbutrin SR, Brand Wellbutrin XL/Forfivo, generic bupropion ER (XL) 450mg tabs, Brand Zoloft, generic sertraline capsules	
Diagnosis	Non-Preferred Drugs
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

**1** - The patient is unresponsive to other treatment modalities, unless contraindicated (i.e. other medications or behavioral modification attempted)

**AND**

**2** - The physician attests that the requested medication is medically necessary. (Document rationale for use)

**AND**

**3** - Patient has a history of failure, contraindication or intolerance to at least 3 preferred alternatives\*

- Bupropion (Generic Wellbutrin)
- Bupropion SR (Generic Wellbutrin SR)
- Bupropion XL (Generic Wellbutrin XL)
- Citalopram (Generic Celexa)
- Duloxetine 20mg, 30mg, or 60 mg capsules
- Escitalopram tablets (Generic Lexapro)
- Esketamine (Spravato)
- Fluoxetine capsules (Generic Prozac)
- Fluoxetine solution (Generic Prozac)
- Fluvoxamine tablets (Generic Luvox)
- Mirtazapine (Generic Remeron)
- Paroxetine tablets (Generic Paxil)
- Sertraline tablets (Generic Zoloft)
- Trazodone (Generic Desyrel)
- Venlafaxine (Generic Effexor)
- Venlafaxine ER capsules (Generic Effexor ER)

Notes

\*Drug may require PA

Product Name: Brand Venlafaxine besylate ER

Diagnosis

Non-Preferred Drugs

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

1 - The patient is unresponsive to other treatment modalities, unless contraindicated (i.e. other medications or behavioral modification attempted)

**AND**

2 - The physician attests that the requested medication is medically necessary. (Document rationale for use)

**AND**

3 - Patient has history of failure or intolerance to preferred generic venlafaxine or venlafaxine ER

**AND**

4 - Patient has a history of failure, contraindication or intolerance to at least 2 preferred alternatives\*

- Bupropion (Generic Wellbutrin)
- Bupropion SR (Generic Wellbutrin SR)
- Bupropion XL (Generic Wellbutrin XL)
- Citalopram (Generic Celexa)
- Duloxetine 20mg, 30mg, or 60 mg capsules
- Escitalopram tablets (Generic Lexapro)
- Esketamine (Spravato)
- Fluoxetine capsules (Generic Prozac)
- Fluoxetine solution (Generic Prozac)
- Fluvoxamine tablets (Generic Luvox)
- Mirtazapine (Generic Remeron)
- Paroxetine tablets (Generic Paxil)
- Sertraline tablets (Generic Zoloft)
- Trazodone (Generic Desyrel)

Notes	*Drug may require PA
-------	----------------------

Product Name: Raldesy	
Diagnosis	Non-Preferred Drugs
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Trial and failure, or intolerance to trazodone oral tablet

**OR**

2 - Patient is unable to swallow the oral tablet formulation of trazodone

**2 . Revision History**

Date	Notes
5/29/2025	Removed Raldesy from Preferred < 6 yo section

Antiemetics - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-148805
<b>Guideline Name</b>	Antiemetics - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Anzemet, granisetron tablet, granisetron injection	
Diagnosis	Nausea and vomiting associated with cancer chemotherapy
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Prevention or treatment of nausea and vomiting associated with cancer chemotherapy	

Product Name:Anzemet, granisetron tablet, granisetron injection
---

Diagnosis	Nausea and vomiting associated with radiotherapy
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Prevention or treatment of nausea and vomiting associated with radiotherapy (total body irradiation, single high-dose fraction to the abdomen, or daily fractions to the abdomen)</p>	

Product Name:Anzemet, granisetron tablet, granisetron injection	
Diagnosis	Postoperative nausea and/or vomiting
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Prevention of postoperative nausea and/or vomiting (administration prior to induction of anesthesia)</p>	

## 2 . Revision History

Date	Notes
6/27/2024	Updated targets (Anzemet, granisetron tablet/inj). Removed ondansetron 24mg as target

Antiglaucoma Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-163933
<b>Guideline Name</b>	Antiglaucoma Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:generic brinzolamide ophth susp, generic tafluprost ophth soln	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of elevated intraocular pressure due to ocular hypertension or open angle glaucoma	

## 2 . Revision History

Date	Notes
1/31/2025	Updated targets to generic Azopt and generic Zioptan

Antipsoriatic Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-99551
<b>Guideline Name</b>	Antipsoriatic Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

### 1 . Criteria

Product Name:Brand Dovonex cream, generic calcipotriene cream, Brand Calcitrene ointment, generic calcipotriene ointment, Brand Vectical, generic calcitriol ointment	
Diagnosis	Psoriasis
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of psoriasis	

**AND**

**2** - History of failure, contraindication, or intolerance to TWO medium to Very high potency corticosteroid topical treatments (see Table 1 in Background section)

## 2 . Background

### Benefit/Coverage/Program Information

**Table 1. Relative Potency of Selected Topical Corticosteroid Products**

<b>Drug</b>	<b>Dosage Form</b>	<b>Strength</b>
<b>Super High Potency</b>		
Augmented betamethasone dipropionate (Diprolene)	Gel, Ointment	0.05%
Clobetasol propionate (Temovate, Temovate E)	Cream, Solution	0.05%
Halobetasol propionate (Ultravate)	Cream	0.05%
<b>High Potency</b>		
Augmented betamethasone dipropionate (Diprolene, Diprolene AF)	Cream, Lotion	0.05%
Betamethasone dipropionate	Lotion, Ointment	0.05%
Fluocinonide (Lidex, Lidex E)	Cream, Solution	0.05%
Triamcinolone acetonide (Kenalog)	Cream, Ointment	0.5%
<b>Medium Potency</b>		

Betamethasone valerate (Beta-Val)	Cream	0.1%
Fluocinolone acetonide (Synalar)	Cream, Ointment	0.025%
Fluticasone propionate (Cutivate)	Cream, Lotion	0.05%
	Ointment	0.005%
Hydrocortisone butyrate (Locoid)	Ointment, Solution	0.1%
Mometasone furoate (Elocon)	Cream, Ointment, Solution	0.1%
Prednicarbate (Dermatop)	Cream	0.1%
Triamcinolone acetonide (Kenalog)	Cream, Lotion, Ointment	0.1%
	Ointment	0.025%

Antipsychotics - AZM

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-226192
<b>Guideline Name</b>	Antipsychotics - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	4/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Brand Abilify tablets, generic aripiprazole tablets and oral solution, Brand Geodon capsules, generic ziprasidone capsules, Brand Latuda, generic lurasidone, lithium carbonate (capsules, tablets, ER tablets, oral solution), Brand Lithobid, Brand Risperdal (tablets, solution) generic risperidone (tablets, ODT, solution), Brand Seroquel, generic quetiapine, Brand Zyprexa, Brand Zyprexa Zydys, generic olanzapine (tablets, ODT)	
Diagnosis	PA Required for Patients < 6 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1** - The patient has been diagnosed per current DSM (Diagnostic and Statistical Manual of Mental Disorders) criteria with one of the following disorders:

- Bipolar Spectrum Disorder
- Schizophrenic Spectrum Disorder
- Tourette's or other tic disorder
- Autism Spectrum Disorder

**AND**

**2** - The requesting clinician has documented that psychosocial issues have been evaluated before request for antipsychotic medications

**AND**

**3** - The requesting clinician has documented non-medication alternatives that have been attempted before request for antipsychotic medications

**AND**

**4** - The above documentation includes information on the expected outcomes and an evaluation of potential adverse events

**AND**

**5** - The patient does not have a known hypersensitivity to the requested agent

Product Name: chlorpromazine tablets, fluphenazine (tablets, oral concentrate, elixir), haloperidol tablets and oral concentrate, loxapine, molindone, perphenazine, pimozide, thioridazine, thiothixene, trifluoperazine

Diagnosis	PA Required for Patients < 12 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization

## Approval Criteria

**1** - The patient has been diagnosed per current DSM (Diagnostic and Statistical Manual of Mental Disorders) criteria with one of the following disorders:

- Bipolar Spectrum Disorder
- Schizophrenic Spectrum Disorder
- Tourette's or other tic disorder
- Autism Spectrum Disorder

**AND**

**2** - The requesting clinician has documented that psychosocial issues have been evaluated before request for antipsychotic medications

**AND**

**3** - The requesting clinician has documented non-medication alternatives that have been attempted before request for antipsychotic medications

**AND**

**4** - The above documentation includes information on the expected outcomes and an evaluation of potential adverse events

**AND**

**5** - The patient does not have a known hypersensitivity to the requested agent

Product Name: chlorpromazine injection, Brand Clozaril, generic clozapine (tablets, ODT), fluphenazine decanoate, Brand Haldol decanoate injection, generic haloperidol decanoate, Brand Haldol lactate injection, generic haloperidol lactate injection

Diagnosis	PA Required for Patients < 18 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - ONE of the following:

**1.1** BOTH of the following:

**1.1.1** ONE of the following:

**1.1.1.1** The requested medication must be used for an FDA (Food and Drug Administration) approved indication

**OR**

**1.1.1.2** The use of the drug is supported by information in ONE of the following appropriate compendia of literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmaco-economic studies
- Other drug reference resources

**AND**

**1.1.2** The patient meets the FDA minimum age limit or the prescriber attests they are aware of FDA labeling regarding the use of the antipsychotic medication and feels the treatment with the requested medication is medically necessary (Document rationale for use)

**OR**

**1.2** The patient is currently on the requested medication

Product Name: Abilify Asimtufii, Abilify Maintena	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has ONE of the following diagnoses:</p> <ul style="list-style-type: none"> <li>Schizophrenia or schizoaffective disorder</li> <li>Bipolar disorder</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <p>2.1 BOTH of the following:</p> <ul style="list-style-type: none"> <li>Patient is non-adherent with oral atypical antipsychotic dosage forms</li> <li>Patient has established tolerability with aripiprazole</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Patient is unable to take oral solid alternatives</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)</p>	

Product Name: Abilify Mycite	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - One of the following:</b></p> <p><b>1.1 All of the following:</b></p> <p><b>1.1.1 Patient is 18 years of age or older</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.1.2 Patient has ONE of the following:</b></p> <ul style="list-style-type: none"> <li>• Schizophrenia or schizoaffective disorder</li> <li>• Bipolar disorder</li> <li>• Autism</li> <li>• Major depressive disorder</li> <li>• Tourette's</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.1.3 Submission of medical records or claims history documenting the patient is currently prescribed aripiprazole and tolerates the medication</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.1.4 Submission of medical records or claims history documenting the patient's adherence to aripiprazole is less than 80 percent within the past 6 months (medication adherence percentage is defined as the number of pills absent in a given time period divided by the number of pills prescribed during that same time, multiplied by 100)</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.1.5 ALL of the following strategies (if applicable to the patient) to improve patient adherence have been tried without success:</b></p> <ul style="list-style-type: none"> <li>• Utilization of a pill box</li> </ul>	

- Utilization of a smart phone reminder (ex. alarm, application, or text reminder)
- Involving family members or friends to assist
- Coordinating timing of dose to coincide with dosing of another daily medication

**AND**

**1.1.6** Submission of medical records or claims history documenting patient has experienced life-threatening or potentially life-threatening symptoms, or has experienced a severe worsening of symptoms leading to a hospitalization which was attributed to the lack of adherence to aripiprazole

**AND**

**1.1.7** Prescriber acknowledges that Abilify MyCite has not been shown to improve patient adherence and attests that Abilify MyCite is medically necessary for the patient to maintain compliance, avoid life-threatening worsening of symptoms, and reduce healthcare resources utilized due to lack of adherence

**AND**

**1.1.8** Prescriber agrees to track and document adherence of Abilify MyCite through software provided by the manufacturer

**AND**

**1.1.9** The patient has a history of failure, contraindication, or intolerance or reason or special circumstance they cannot use TWO of the following: (Drug may require PA)

- Abilify Maintena
- Invega Sustenna
- Risperdal Consta
- Aristada
- Perseris

**OR**

**1.2** ONE of the following:

**1.2.1** The patient has been receiving treatment with the requested non-preferred behavioral health medication and is new to the plan (enrollment effective date within the past 90 days)

**OR**

**1.2.2** The patient is currently receiving treatment with the requested non-preferred behavioral health medication in the hospital and must continue upon discharge

Product Name: Abilify Mycite

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Documentation that patient is clinically stable on Abilify MyCite

**AND**

**2** - Submission of medical records or claims history documenting that the use of Abilify MyCite has increased adherence to 80 percent or more

**AND**

**3** - Prescriber attests that the patient requires the continued use of Abilify MyCite to remain adherent

Product Name: Aristada, Aristada Initio

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Patient has a diagnosis of schizophrenia or schizoaffective disorder

**AND**

2 - ONE of the following:

2.1 BOTH of the following:

- Patient is non-adherent with oral atypical antipsychotic dosage forms
- Patient has established tolerability with oral aripiprazole

**OR**

2.2 Patient is unable to take oral solid alternatives

**AND**

3 - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name:Erzofri, Invega Sustenna	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has a diagnosis of schizophrenia or schizoaffective disorder	
<b>AND</b>	
2 - ONE of the following:	

**2.1 BOTH** of the following:

- Patient is non-adherent with oral atypical antipsychotic dosage forms
- Patient has established tolerability with oral paliperidone or oral risperidone

**OR**

**2.2** Patient is unable to take oral solid alternatives

**AND**

**3** - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

**AND**

**4** - For Erzofri requests ONLY: History of failure or adverse reaction to Invega Sustenna (Applies to Erzofri requests only)

Product Name: Invega Trinza

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Patient has a diagnosis of schizophrenia or schizoaffective disorder

**AND**

**2** - Patient has been treated with Invega Sustenna for at least 4 months

**AND**

**3** - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name: Invega Hafyera

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Patient has a diagnosis of schizophrenia or schizoaffective disorder

**AND**

**2** - Patient has been treated with Invega Sustenna or Invega Trinza for at least 6 months

**AND**

**3** - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name: Opipza

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1 - ONE of the following:**

**1.1 Both of the following:**

- Patient is 13 years of age or older
- Diagnosis of schizophrenia

**OR**

**1.2 Both of the following:**

- Patient is 18 years of age or older
- Diagnosis of major depressive disorder (MDD)

**OR**

**1.3 Both of the following:**

- Patient is 6 years of age or older
- Diagnosis of autistic disorder

**OR**

**1.4 Both of the following:**

- Patient is 6 years of age or older
- Diagnosis of Tourette's disorder

**AND**

**2 - BOTH of the following:**

**2.1 ONE of the following:**

- History of failure or intolerance to aripiprazole oral tablets
- History of failure or intolerance to aripiprazole oral solution

**AND**

**2.2** Provider submits clinical rationale documenting why patient is unable to take aripiprazole ODT formulation

**AND**

**3** - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name:Perseris	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient has a diagnosis of schizophrenia or schizoaffective disorder</p> <p><b>AND</b></p> <p><b>2</b> - ONE of the following:</p> <p><b>2.1</b> BOTH of the following:</p> <ul style="list-style-type: none"><li>• Patient is non-adherent with oral atypical antipsychotic dosage forms</li><li>• Patient has established tolerability with oral risperidone</li></ul> <p><b>OR</b></p> <p><b>2.2</b> Patient is unable to take oral solid alternatives</p> <p><b>AND</b></p>	

**3** - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name:Risperdal Consta

Approval Length | 12 month(s)

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Patient has ONE of the following diagnoses:

- Schizophrenia or schizoaffective disorder
- Bipolar disorder

**AND**

**2** - ONE of the following:

**2.1** BOTH of the following:

- Patient is non-adherent with oral atypical antipsychotic dosage forms
- Patient has established tolerability with oral risperidone

**OR**

**2.2** Patient is unable to take oral solid alternatives

**AND**

**3** - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name: Rykindo, generic risperidone ER IM

Approval Length | 12 month(s)

Guideline Type | Prior Authorization

**Approval Criteria**

1 - Patient has ONE of the following diagnoses:

- Schizophrenia or schizoaffective disorder
- Bipolar disorder

**AND**

2 - ONE of the following:

2.1 BOTH of the following:

- Patient is non-adherent with oral atypical antipsychotic dosage forms
- Patient has established tolerability with oral risperidone

**OR**

2.2 Patient is unable to take oral solid alternatives

**AND**

3 - History of failure, contraindication or intolerance to BOTH of the following:

- Perseris
- Risperdal Consta

**AND**

4 - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

Product Name: Uzedy	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has a diagnosis of schizophrenia or schizoaffective disorder</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <p>2.1 BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Patient is non-adherent with oral atypical antipsychotic dosage forms</li> <li>• Patient has established tolerability with oral risperidone</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Patient is unable to take oral solid alternatives</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - History of failure, contraindication or intolerance to BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Perseris</li> <li>• Risperdal Consta</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - If the patient is less than 18 years of age, the prescriber attests they are aware of FDA (Food and Drug Administration) labeling regarding use of long acting injectable antipsychotic products in patients less than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)</p>	

Product Name: Brand Abilify tablets, Adasuve, aripiprazole ODT, chlorpromazine tablets, Brand Clozaril, fluphenazine (injection, tablets), Brand Geodon (capsules, injection) generic ziprasidone (capsules, injection), Brand Haldol decanoate injection, generic haloperidol injection, Brand Invega tablets, generic paliperidone ER tablets, Brand Latuda, Brand Lithobid, loxapine, perphenazine-amitriptyline, prochlorperazine injection, Brand Risperdal (tablets, solution), generic risperidone solution, Brand Saphris, generic asenapine sublingual tablets, Secuado, Brand Seroquel, Brand Seroquel XR, generic quetiapine ER tablets, Brand Symbyax, generic fluoxetine-olanzapine, Versacloz, Brand Zyprexa (tablets, injection), Zyprexa Relprevv, Brand Zyprexa Zydis

Diagnosis	Non-Preferred Drugs
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - One of the following:

1.1 All of the following:

1.1.1 ONE of the following:

1.1.1.1 Patient has a history of failure, contraindication or intolerance to at least FOUR of the following:

- Aripiprazole oral (generic Abilify)
- Aripiprazole injectable formulations (Abilify Maintena, Aristada, Aristada Initio)
- Clozapine/clozapine ODT
- Lurasidone
- Olanzapine/olanzapine ODT
- Paliperidone oral\*\* (DOES NOT APPLY TO REQUESTS FOR PALIPERIDONE ER TABLETS)
- Paliperidone injectable formulations (Invega Sustenna, Invega Trinza, Hafyera)
- Quetiapine
- Risperidone/risperidone ODT
- Risperidone injectable formulations (Perseris, Risperdal Consta)

**OR**

1.1.1.2 There are no preferred formulary alternatives for the requested drug

**AND**

**1.1.2** If the request is for a multi-source brand medication (i.e., MSC O), ONE of the following:

**1.1.2.1** BOTH of the following:

- The brand is being requested because of an adverse reaction, allergy or sensitivity to the generic and the prescriber must attest to submitting the FDA MedWatch Form for allergic reactions to the medications
- If there are generic product(s), the member has tried at least three (if available)

**OR**

**1.1.2.2** ONE of the following:

- The brand is being requested due to a therapeutic failure with the generic (please provide reason for therapeutic failure)
- The brand is being requested because transition to the generic could result in destabilization of the patient (rationale must be provided)
- Special clinical circumstances exist that preclude the use of the generic equivalent of the multi-source brand medication for the patient (rationale must be provided)

**AND**

**1.1.3** ONE of the following:

**1.1.3.1** The requested drug must be used for an FDA (Food and Drug Administration)-approved indication

**OR**

**1.1.3.2** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System

- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmacoeconomic studies
- Other drug reference resources

**AND**

**1.1.4** ONE of the following:

**1.1.4.1** The drug is being prescribed within the manufacturer's published dosing guidelines

**OR**

**1.1.4.2** The drug falls within dosing guidelines found in ONE of the following compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmacoeconomic studies
- Other drug reference resources

**AND**

**1.1.5** The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plans' program\*

**OR**

**1.2** The requested medication is a behavioral health medication and ONE of the following:

**1.2.1** The patient has been receiving treatment with the requested non-preferred behavioral health medication and is new to the plan (enrollment effective date within the past 90 days)

**OR**

**1.2.2** The patient is currently receiving treatment with the requested non-preferred behavioral health medication in the hospital and must continue upon discharge

Notes

\*Note: Medications used solely for anti-obesity/weight loss, cosmetic ( e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs ( Cialis/Tadalafil) are covered for clinical diagnoses other than ED.

\*\*If the request is for generic paliperidone ER tablets, please omit "paliperidone oral" as an alternative\*\*

## 2 . Revision History

Date	Notes
3/26/2025	Added step through Perseris to Rykindo section.

Anxiolytics - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-126970
<b>Guideline Name</b>	Anxiolytics - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/23/2023
-----------------	-----------

## 1 . Criteria

Product Name:buspirone, Brand Xanax tabs, generic alprazolam tabs, alprazolam ODT, alprazolam conc, Brand Xanax XR, generic alprazolam ER, chlordiazepoxide, Brand Tranxene T, generic clorazepate dipotassium, Brand Valium tabs, generic diazepam tabs, diazepam conc, diazepam oral soln, Brand Ativan, Loreev XR, generic lorazepam, lorazepam conc, generic oxazepam, Brand Klonopin tabs, generic clonazepam tabs, clonazepam ODT	
Diagnosis	Requests for Patients less than 6 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - The patient is unresponsive to other treatment modalities, unless contraindicated (i.e. other medications or behavioral modification attempted).

**AND**

2 - The physician attests that the requested medication is medically necessary (Document rationale for use)

Product Name:Loreev XR

Diagnosis	Requests for Patients 6 years of age and older
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Trial and failure, or contraindication to generic lorazepam

**AND**

2 - The physician attests that the requested medication is medically necessary (Document rationale for use)

Product Name:buspirone, Brand Xanax tabs, generic alprazolam tabs, alprazolam ODT, alprazolam conc, Brand Xanax XR, generic alprazolam ER, chlordiazepoxide, Brand Tranxene T, generic clorazepate dipotassium, Brand Valium tabs, generic diazepam tabs, diazepam conc, diazepam oral soln, Brand Ativan, Loreev XR, generic lorazepam, lorazepam conc, generic oxazepam, Brand Klonopin tabs, generic clonazepam tabs, clonazepam ODT

Diagnosis	Reject 75: Drug Utilization Review: Greater than 1 Anxiolytic in 30 days
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - The medication is being used to adjust the dose of the drug

**OR**

2 - The medication will be used in place of the previously prescribed drug, and not in addition to it

**OR**

3 - The medication dosage form will be used in place of the previously prescribed medication dosage form, and not in addition to it

**OR**

4 - The physician attests they are aware of the multiple anxiolytics prescribed to the patient and feels treatment with both medications is medically necessary (Document rationale for use)

## 2 . Revision History

Date	Notes
6/22/2023	Updated DUR Reject code from 88 to rej 75. No changes to clinical c riteria.

Apomorphine products (Apokyn, Onapgo)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-221195
<b>Guideline Name</b>	Apomorphine products (Apokyn, Onapgo)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Apokyn, generic apomorphine injection, Onapgo	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting all of the following:  1.1 Diagnosis of Parkinson's disease	

**AND**

**1.2** Medication will be used as intermittent treatment for OFF episodes

**AND**

**1.3** Patient is currently on a stable dose of a carbidopa/levodopa-containing medication and will continue receiving treatment with a carbidopa/levodopa-containing medication while on therapy

**AND**

**1.4** Patient continues to experience greater than or equal to 2 hours of OFF time per day despite optimal management of carbidopa/levodopa therapy including BOTH of the following:

- Taking carbidopa/levodopa on an empty stomach or at least one half-hour or more before or one hour after a meal or avoidance of high protein diet
- Dose and dosing interval optimization

**AND**

**1.5** History of failure, contraindication, or intolerance to TWO anti-Parkinson's disease therapies from the following adjunctive pharmacotherapy classes (trial must be from two different classes):

- Dopamine agonists (e.g., pramipexole, ropinirole)
- Catechol-O-methyl transferase (COMT) inhibitors (e.g., entacapone)
- Monoamine oxidase (MAO) B inhibitors (e.g., rasagiline, selegiline)

**AND**

**2** - Prescribed by or in consultation with a neurologist or specialist in the treatment of Parkinson's disease

Product Name: Brand Apokyn, generic apomorphine injection, Onapgo

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient will continue to receive treatment with a carbidopa/levodopa-containing medication</p>	

**2 . Revision History**

Date	Notes
3/26/2025	Added Onapgo, removed Kynmobi, updated guideline name.

Aqneursa (levacetylleucine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-240206
<b>Guideline Name</b>	Aqneursa (levacetylleucine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/15/2025
-----------------	-----------

## 1 . Criteria

Product Name:Aqneursa	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Niemann-Pick disease type C (NPC)  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) documenting the diagnosis has been confirmed by one of the following:

**2.1** Genetically confirmed (deoxyribonucleic acid [DNA] sequence analysis) by mutations in both alleles of NPC1 or NPC2

**OR**

**2.2** Mutation in only one allele of NPC1 or NPC2 plus either positive filipin staining or elevated cholestane triol/oxysterols (>2 x upper limit of normal)

**AND**

**3** - Patient has at least one neurological symptom of the disease (e.g., hearing loss, vertical supranuclear gaze palsy, ataxia, dementia, dystonia, seizures, dysarthria, or dysphagia)

**AND**

**4** - Patient weighs greater than or equal to 15kg

**AND**

**5** - Requested drug will not be used in combination with Miplyffa (arimoclomol). \*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.

**AND**

**6** - Prescribed by or in consultation with a specialist knowledgeable in the treatment of Niemann-Pick disease type C

Notes	*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.
-------	--

Product Name:Aqneursa

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy (e.g., slowing of disease progression, improvement in neurological symptoms of the disease)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Requested drug will not be used in combination with Miplyffa (arimoclomol). *The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.</p>	
Notes	*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.

**2 . Revision History**

Date	Notes
4/15/2025	Added verbiage regarding concomitant use with Miplyffa, no change to clinical criteria

Aquadeks

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99514
<b>Guideline Name</b>	Aquadeks
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Aquadeks	
Diagnosis	Cystic Fibrosis
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystic fibrosis	

## 2 . Revision History

Date	Notes
4/10/2021	7/1 Implementation

Arcalyst (rilonacept)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152659
<b>Guideline Name</b>	Arcalyst (rilonacept)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Arcalyst	
Diagnosis	Cryopyrin-Associated Periodic Syndromes (CAPS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting a diagnosis of Cryopyrin-Associated Periodic Syndromes (CAPS) [including Familial Cold Auto-inflammatory Syndrome (FCAS), Muckle-Wells Syndrome (MWS)]	

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Immunologist
- Allergist
- Dermatologist
- Rheumatologist
- Neurologist
- Specialist with expertise in the management of CAPS

**AND**

**3** - The medication will not be used in combination with another biologic agent

Product Name:Arcalyst	
Diagnosis	Cryopyrin-Associated Periodic Syndromes (CAPS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes) documenting patient has experienced disease stability or improvement in clinical symptoms while on therapy as evidenced by one of the following:	
<ul style="list-style-type: none"><li>• Improvement in rash, fever, joint pain, headache, or conjunctivitis</li><li>• Decreased number of disease flare days</li><li>• Normalization of inflammatory markers (C-reactive protein [CRP], erythrocyte sedimentation rate [ESR], serum amyloid A [SAA])</li><li>• Corticosteroid dose reduction</li><li>• Improvement in MD global score or active joint count</li></ul>	

Product Name:Arcalyst

Diagnosis	Deficiency of Interleukin-1 Receptor Antagonist (DIRA)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting ALL of the following:</p> <p>1.1 Diagnosis of deficiency of interleukin-1 receptor antagonist (DIRA)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Patient weighs at least 10 kg</p> <p style="text-align: center;"><b>AND</b></p> <p>1.3 Patient is currently in remission (e.g., no fever, skin rash, and bone pain; no radiological evidence of active bone lesions; C-reactive protein [CRP] less than 5 mg/L)</p>	

Product Name:Arcalyst	
Diagnosis	Recurrent Pericarditis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:</p> <p>1.1 Diagnosis of recurrent pericarditis as evidenced by at least 2 episodes that occur a minimum of 4 to 6 weeks apart</p>	

**AND**

**1.2** History of failure, contraindication, or intolerance to ALL of the following:

- nonsteroidal anti-inflammatory drugs (e.g., ibuprofen, naproxen)
- colchicine
- corticosteroids (e.g., prednisone)

**AND**

**2** - Prescribed by or in consultation with a cardiologist

Product Name:Arcalyst	
Diagnosis	Recurrent Pericarditis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy	

## 2 . Revision History

Date	Notes
8/27/2024	Added criteria for DIRA and recurrent pericarditis. Updated CAPS criteria.

Arikayce

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99710
<b>Guideline Name</b>	Arikayce
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Arikayce	
Diagnosis	Refractory Mycobacterium avium complex (MAC) lung disease
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of refractory Mycobacterium avium complex (MAC) lung disease	

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) or claims history documenting respiratory cultures positive for MAC within the previous 6 months

**AND**

**3** - Submission of medical records (e.g., chart notes, laboratory values) or claims history documenting the patient has been receiving a multidrug background regimen containing at least TWO of the following agents for a minimum of 6 consecutive months within the past 12 months (prescription claims history may be used in conjunction as documentation of medication use, dose, and duration):

- Macrolide antibiotic\* (e.g., azithromycin, clarithromycin)
- Ethambutol\*
- Rifamycin antibiotic\* (e.g., rifampin, rifabutin)

**AND**

**4** - Patient will continue to receive a multidrug background regimen

**AND**

**5** - Documentation that the patient has not achieved negative sputum cultures after receipt of a multidrug background regimen for a minimum of 6 consecutive months

**AND**

**6** - In vitro susceptibility testing of recent (within 6 months) positive culture documents that the MAC isolate is susceptible to amikacin with a minimum inhibitory concentration (MIC) of less than or equal to 64 micrograms per milliliter (mcg/mL)

**AND**

**7** - Prescribed by or in consultation with one of the following:

<ul style="list-style-type: none"> <li>• Infectious disease specialist</li> <li>• Pulmonologist</li> </ul>	
Notes	*Drug may require PA)

Product Name:Arikayce	
Diagnosis	Refractory Mycobacterium avium complex (MAC) lung disease
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Documentation that the patient has achieved negative respiratory cultures</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 ALL of the following:</p> <p>1.2.1 Patient has not achieved negative respiratory cultures while on Arikayce</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2.2 Physician attestation that patient has demonstrated clinical benefit while on Arikayce</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2.3 In vitro susceptibility testing of most recent (within 6 months) positive culture with available susceptibility testing documents that the Mycobacterium avium complex (MAC) isolate is susceptible to amikacin with an minimum inhibitory concentration (MIC) of less than 64 micrograms per milliliter (mcg/mL)</p>	
--	--

**AND**

**1.2.4** Patient has NOT received greater than 12 months of Arikayce therapy with continued positive respiratory cultures

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) or claims history documenting that the patient continues to receive a multidrug background regimen containing at least TWO of the following agents (prescription claims history may be used in conjunction as documentation of medication use, dose, and duration):

- Macrolide antibiotic\* (e.g., azithromycin, clarithromycin)
- Ethambutol\*
- Rifamycin antibiotic\* (e.g., rifampin, rifabutin)

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Infectious disease specialist
- Pulmonologist

Notes

\*Drug may require PA

## 2 . Revision History

Date	Notes
5/12/2021	Arizona Medicaid 7.1 Implementation

Atorvaliq (atorvastatin oral suspension)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-125916
<b>Guideline Name</b>	Atorvaliq (atorvastatin oral suspension)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/20/2023
-----------------	-----------

## 1 . Criteria

Product Name:Atorvaliq	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Both of the following:  1.1.1 Patient is less than 10 years of age	

**AND**

**1.1.2** Prescribed by or in consultation with a cardiologist

**OR**

**1.2** Both of the following:

**1.2.1** Medication is being used for one of the following:

**1.2.1.1** To reduce the risk of one of the following:

- Myocardial infarction (MI), stroke, revascularization procedures, and angina in adults with multiple risk factors for coronary heart disease (CHD) but without clinically evident CHD
- MI and stroke in adults with type 2 diabetes mellitus with multiple risk factors for CHD but without clinically evident CHD
- Non-fatal MI, fatal and non-fatal stroke, revascularization procedures, hospitalization for congestive heart failure, and angina in adults with clinically evident CHD

**OR**

**1.2.1.2** As an adjunct to diet to reduce low-density lipoprotein cholesterol (LDL-C) in one of the following:

- Adults with primary hyperlipidemia
- Adults and pediatric patients aged 10 years and older with heterozygous familial hypercholesterolemia (HeFH)

**OR**

**1.2.1.3** As an adjunct to other LDL-C-lowering therapies, or alone if such treatments are unavailable, to reduce LDL-C in adults and pediatric patients aged 10 years and older with homozygous familial hypercholesterolemia (HoFH)

**OR**

**1.2.1.4** As an adjunct to diet for the treatment of adults with one of the following:

- Primary dysbetalipoproteinemia
- Hypertriglyceridemia

**AND**

**1.2.2** One of the following:

**1.2.2.1** Trial and failure, contraindication, or intolerance to generic atorvastatin tablets (verified via paid pharmacy claims or submitted chart notes)

**OR**

**1.2.2.2** Patient is unable to swallow oral tablets

## 2 . Revision History

Date	Notes
5/19/2023	Revised verbiage for patients under 10 yo

Attruby (acoramidis)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-193195
<b>Guideline Name</b>	Attruby (acoramidis)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Attruby	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of transthyretin-mediated amyloidosis with cardiomyopathy (ATTR-CM)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting one of the following:

**2.1** Presence of a transthyretin (TTR) mutation (e.g., V122I) as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)

**OR**

**2.2** Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of TTR amyloid deposits

**OR**

**2.3** Both of the following:

- Cardiac magnetic resonance imaging or scintigraphy scan suggestive of amyloidosis
- Absence of light-chain amyloidosis

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting patient has New York Heart Association (NYHA) Functional Class I, II, or III heart failure

**AND**

**4** - Requested drug is not used in combination with a TTR silencer (e.g., Amvuttra) or a TTR stabilizer (e.g., Diflunisal)

**AND**

**5** - Prescribed by or in consultation with a cardiologist

Product Name:Attruby	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Requested drug is not used in combination with a TTR silencer (e.g., Amvuttra) or a TTR stabilizer (e.g., Diflunisal)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with a cardiologist</p>	

## 2 . Revision History

Date	Notes
2/25/2025	New program

Austedo (deutetrabenazine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-155386
<b>Guideline Name</b>	Austedo (deutetrabenazine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Austedo, Austedo XR	
Diagnosis	Moderate to Severe Tardive dyskinesia
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of moderate to severe tardive dyskinesia (TD) secondary to treatment with a centrally acting dopamine receptor blocking agent (DRBA)	

**AND**

**2** - Prescribed by or in consultation with a psychiatrist or neurologist

**AND**

**3** - Patient is 18 years of age or older

**AND**

**4** - Patient has an Abnormal Involuntary Movement Scale (AIMS) score of 3 or 4 on any one of the AIMS items 1 through 9

**AND**

**5** - Austedo is not prescribed concurrently with tetrabenazine or Ingrezza

**AND**

**6** - Dose does not exceed 48 mg per day

Notes	*NOTE: Patients will be approved for ONE strength of Austedo/Austedo XR ONLY: Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Austedo products. End-date/Retire all other active PA's for Austedo products.
-------	---

Product Name:Austedo, Austedo XR	
Diagnosis	Moderate to Severe Tardive dyskinesia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient is responding positively to therapy as evidenced by a reduction in the baseline score of any one of the AIMS items 1 through 9

**AND**

2 - Austedo is not prescribed concurrently with tetrabenazine or Ingrezza

**AND**

3 - Dose does not exceed 48 mg per day

Notes

\*NOTE: Patients will be approved for ONE strength of Austedo/Austedo XR ONLY: Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Austedo products. End-date/Retire all other active PA's for Austedo products.

Product Name:Austedo, Austedo XR

Diagnosis Chorea Associated with Huntington Disease

Approval Length 6 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Diagnosis of chorea associated with Huntington's Disease

**AND**

2 - Prescribed by or in consultation with a neurologist

**AND**

3 - Patient is 18 years of age or older

**AND**

**4** - Targeted mutation analysis demonstrates a cytosine-adenine-guanine (CAG) trinucleotide expansion of  $\geq 36$  repeats in the huntingtin (HTT) gene

**AND**

**5** - Patient has a Unified Huntington Disease Rating Scale (UHDRS) score ranging from 1 to 4 on any one of UHDRS chorea items 1 through 7

**AND**

**6** - Austedo is not prescribed concurrently with tetrabenazine or Ingrezza

**AND**

**7** - Dose does not exceed 48 mg per day

Notes

\*NOTE: Patients will be approved for ONE strength of Austedo/Austedo XR ONLY: Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Austedo products. End-date/Retire all other active PA's for Austedo products.

Product Name:Austedo, Austedo XR

Diagnosis | Chorea Associated with Huntington Disease

Approval Length | 12 month(s)

Therapy Stage | Reauthorization

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Patient is responding positively to therapy as evidenced by a reduction in the baseline score of any one of the UHDRS chorea items 1 through 7

**AND**

**2** - Austedo is not prescribed concurrently with tetrabenazine or Ingrezza

**AND**

**3** - Dose does not exceed 48 mg per day

Notes

\*NOTE: Patients will be approved for ONE strength of Austedo/Austedo XR ONLY: Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Austedo products. End-date/Retire all other active PA's for Austedo products.

## 2 . Revision History

Date	Notes
9/24/2024	Added note for PA Team, only 1 strength of Austedo can be approved

Azole Antifungals

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-143793
<b>Guideline Name</b>	Azole Antifungals
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	3/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Brand Sporanox capsules, generic itraconazole capsules	
Diagnosis	Systemic Fungal Infections
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following</p> <p>1.1 Diagnosis of ONE of the following:</p> <ul style="list-style-type: none"><li>Blastomycosis</li></ul>	

- Histoplasmosis
- Aspergillosis

**OR**

**1.2** Both of the following:

**1.2.1** Diagnosis of coccidioidomycosis

**AND**

**1.2.2** Patient has a history of failure, contraindication, intolerance, or resistance to fluconazole (generic Diflucan) as evidenced by submission of medical records or claims history

Product Name: Brand Sporanox capsules, generic itraconazole capsules	
Diagnosis	Onychomycosis Fingernails
Approval Length	2 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of fingernail onychomycosis confirmed by ONE of the following:</p> <ul style="list-style-type: none"> <li>• KOH (potassium hydroxide) test</li> <li>• Fungal culture</li> <li>• Nail biopsy</li> </ul> <p><b>AND</b></p> <p>2 - Patient has a history of at least a 6-week trial resulting in therapeutic failure, contraindication, intolerance, or resistance to Terbinafine as evidenced by submission of medical records or claims history</p>	

Product Name: Brand Sporanox capsules, generic itraconazole capsules	
Diagnosis	Onychomycosis Fingernails
Approval Length	2 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Both of the following:</p> <p>1.1 Three months have elapsed since completion of initial therapy for fingernail onychomycosis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Documentation of positive clinical response to therapy</p>	

Product Name: Brand Sporanox capsules, generic itraconazole capsules	
Diagnosis	Onychomycosis Toenails
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of toenail onychomycosis confirmed by ONE of the following:</p> <ul style="list-style-type: none"> <li>• KOH (potassium hydroxide) test</li> <li>• Fungal culture</li> <li>• Nail biopsy</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has a history of at least a 12-week trial resulting in therapeutic failure,</p>	

contraindication, intolerance, or resistance to Terbinafine as evidenced by submission of medical records or claims history.

Product Name:Brand Sporanox capsules, generic itraconazole capsules	
Diagnosis	Onychomycosis Toenails
Approval Length	3 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - BOTH of the following:</p> <p>1.1 Nine months have elapsed since completion of initial therapy for toenail onychomycosis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Documentation of positive clinical response to therapy</p>	

Product Name:Brand Sporanox Oral Solution, generic itraconazole oral solution	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following diagnoses:</p> <ul style="list-style-type: none"><li>• Oropharyngeal candidiasis</li><li>• Esophageal candidiasis</li></ul>	

Product Name:Brand Vfend tablets, generic voriconazole tablets	
Approval Length	12 month(s)

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - One of the following:</b></p> <p><b>1.1</b> Diagnosis of invasive aspergillosis including <i>Aspergillus fumigatus</i></p> <p style="text-align: center;"><b>OR</b></p> <p><b>1.2</b> ALL of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of Candidemia</li> <li>• Patient is non-neutropenic</li> <li>• Patient has a history of failure, contraindication, intolerance, or resistance to fluconazole (generic Diflucan) as evidenced by submission of medical records or claims history</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>1.3</b> Both of the following:</p> <p><b>1.3.1</b> ONE of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Candida infection in the abdomen</li> <li>• Candida infection in the kidney</li> <li>• Candida infection in the bladder wall</li> <li>• Candida infection in wounds</li> <li>• Disseminated Candida infections in skin</li> <li>• Esophageal candidiasis</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.3.2</b> Patient has a history of failure, contraindication, intolerance, or resistance to fluconazole (generic Diflucan) as evidenced by submission of medical records or claims history</p> <p style="text-align: center;"><b>OR</b></p>	

**1.4** Diagnosis of *Scedosporium apiospermum* infection (asexual form of *Pseudallescheria boydii*)

**OR**

**1.5** Diagnosis of *Fusarium* spp. infection including *Fusarium solani*

**OR**

**1.6** Diagnosis of *Exserohilum* species infection

Product Name: Brand Vfend Powder for Oral Suspension, generic voriconazole powder for oral suspension

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Both of the following:

**1.1** One of the following:

**1.1.1** Diagnosis of invasive aspergillosis including *Aspergillus fumigatus*

**OR**

**1.1.2** ALL of the following:

- Diagnosis of Candidemia
- Patient is non-neutropenic
- Patient has a history of failure, contraindication, intolerance, or resistance to fluconazole (generic Diflucan) as evidenced by submission of medical records or claims history

**OR**

**1.1.3** ONE of the following diagnoses:

- Candida infection in the abdomen
- Candida infection in the kidney
- Candida infection in the bladder wall
- Candida infection in wounds
- Disseminated Candida infections in skin
- Esophageal candidiasis

**OR**

**1.1.4** Diagnosis of *Scedosporium apiospermum* infection (asexual form of *Pseudallescheria boydii*)

**OR**

**1.1.5** Diagnosis of *Fusarium* spp. infection including *Fusarium solani*

**OR**

**1.1.6** Diagnosis of *Exserohilum* species infection

**AND**

**1.2** Physician has provided rationale for the patient needing to use voriconazole oral suspension instead of voriconazole tablets.

Product Name: Brand Noxafil tablets, generic posaconazole tablets	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - BOTH of the following:	

**1.1** Used as prophylaxis of invasive fungal infections caused by ONE of the following:

- Aspergillus
- Candida

**AND**

**1.2** One of the following conditions:

**1.2.1** Patient is at high risk of infections due to severe immunosuppression from ONE of the following conditions:

- Hematopoietic stem cell transplant (HSCT) with graft-versus-host disease (GVHD)
- Hematologic malignancies with prolonged neutropenia from chemotherapy [eg, acute myeloid leukemia (AML), myelodysplastic syndromes (MDS)]

**OR**

**1.2.2** Patient has a prior fungal infection requiring secondary prophylaxis

Product Name: Noxafil Suspension, Noxafil suspension packets

Diagnosis	Prophylaxis of Aspergillus or Candida Infections
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - BOTH of the following:

**1.1** Used as prophylaxis of invasive fungal infections caused by ONE of the following:

- Aspergillus
- Candida

**AND**

**1.2** One of the following conditions:

**1.2.1** Patient is at high risk of infections due to severe immunosuppression from ONE of the following conditions:

- Hematopoietic stem cell transplant (HSCT) with graft-versus-host disease (GVHD)
- Hematologic malignancies with prolonged neutropenia from chemotherapy [eg, acute myeloid leukemia (AML), myelodysplastic syndromes (MDS)]

**OR**

**1.2.2** Patient has a prior fungal infection requiring secondary prophylaxis

Product Name: Noxafil Suspension	
Diagnosis	Oropharyngeal Candidiasis (OPC)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - BOTH of the following:</p> <p><b>1.1</b> Diagnosis of oropharyngeal candidiasis (OPC)</p> <p><b>AND</b></p> <p><b>1.2</b> The patient has a history of failure, contraindication, intolerance, or resistance to TWO of the following as evidenced by submission of medical records or claims history:</p> <ul style="list-style-type: none"><li>• Fluconazole* (generic Diflucan)</li><li>• Itraconazole* (generic Sporanox)</li><li>• Clotrimazole Lozenges*</li></ul>	
Notes	*Drug may require PA

Product Name: Cresemba

Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Both of the following:</p> <p>1.1.1 Diagnosis of invasive aspergillosis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.2 Patient has a history of failure, contraindication, intolerance, or resistance to voriconazole* (generic Vfend) as evidenced by submission of medical records or claims history</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 Diagnosis of invasive mucormycosis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Both of the following:</p> <ul style="list-style-type: none"> <li>• Patient is 6 months of age or older</li> <li>• Patient weighs 16 kg or greater</li> </ul>	
Notes	*Drug may require PA

Product Name: Tolsura	
Approval Length	3 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Both of the following:

1.1 Diagnosis of ONE of the following fungal infections:

- Blastomycosis
- Histoplasmosis
- Aspergillosis

**AND**

1.2 Patient has a history of failure, contraindication, intolerance, or resistance to itraconazole\* capsules (generic Sporanox) as evidenced by submission of medical records or claims history

Notes	*Drug may require PA
-------	----------------------

Product Name: Brand Sporanox capsules, generic itraconazole capsules, Brand Sporanox oral solution, generic itraconazole oral solution, Brand Vfend tablets, generic voriconazole tablets, Brand Vfend powder for oral suspension, generic voriconazole powder for oral suspension, Brand Noxafil tablets, generic posaconazole tablets, Noxafil oral suspension, Noxafil suspension packets, Cresemba, Tolsura

Diagnosis	All Other Diagnoses
Guideline Type	Prior Authorization

**Approval Criteria**

1 - The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmaco-economic studies

- Other drug reference resources

**AND**

**2** - The medication is being prescribed by or in consultation with an infectious disease specialist

Notes	*Authorization duration based on provider recommended treatment durations, not to exceed 12 months
-------	--

## 2 . Revision History

Date	Notes
3/1/2024	Removed Likmez from PA, created new drug-specific guideline.

Baclofen products (Fleqsuvy, Lyvispah, Ozobax/Ozobax DS)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-211208
<b>Guideline Name</b>	Baclofen products (Fleqsuvy, Lyvispah, Ozobax/Ozobax DS)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Brand Fleqsuvy, generic baclofen suspension, Lyvispah, Ozobax/DS, Brand Baclofen oral solution, generic baclofen oral solution	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Trial and failure, or intolerance to baclofen tablets  <b>OR</b>	

2 - Patient is unable to swallow oral tablets

## 2 . Revision History

Date	Notes
3/26/2025	Added Brand/generic Fleqsuvy as targets.

Baxdela

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99516
<b>Guideline Name</b>	Baxdela
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Baxdela	
Diagnosis	Community-Acquired Bacterial Pneumonia
Approval Length	10 Days*
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - For continuation of therapy upon hospital discharge  <b>OR</b>	

**2** - As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

**3** - All of the following:

**3.1** Diagnosis of community-acquired bacterial pneumonia (CABP)

**AND**

**3.2** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Baxdela

**AND**

**3.3** History of failure, contraindication, or intolerance to **THREE** of the following antibiotics or antibiotic regimens:

- Amoxicillin\*\*
- A macrolide\*\*
- Doxycycline\*\*
- A fluoroquinolone\*\*
- Combination therapy with amoxicillin/clavulanate or cephalosporin **AND** a macrolide or doxycycline

Notes	*Note: Authorization will be issued for up to 10 days. **Drug may require PA
-------	---

Product Name: Baxdela	
Diagnosis	Acute Bacterial Skin and Skin Structure Infections
Approval Length	14 Days*
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - For continuation of therapy upon hospital discharge

**OR**

2 - As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

3 - All of the following:

3.1 One of the following diagnoses:

3.1.1 Both of the following

3.1.1.1 Acute bacterial skin and skin structure infections

**AND**

3.1.1.2 Infection caused by methicillin-resistant *Staphylococcus aureus* (MRSA) documented by culture and sensitivity report

**OR**

3.1.2 Both of the following:

3.1.2.1 Empirical treatment of patients with acute bacterial skin and skin structure infections

**AND**

3.1.2.2 Presence of MRSA infection is likely

**AND**

3.2 History of failure, contraindication, or intolerance to linezolid (generic Zyvox)

**AND**

**3.3** History of failure, contraindication, or intolerance to ONE of the following antibiotics:

- Sulfamethoxazole-trimethoprim (SMZ-TMP)\*\*
- A tetracycline\*\*
- Clindamycin\*\*

**OR**

**4** - All of the following:

**4.1** Diagnosis of acute bacterial skin and skin structure infections

**AND**

**4.2** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Baxdela

**AND**

**4.3** History of failure, contraindication, or intolerance to THREE of the following antibiotics:

- A penicillin\*\*
- A cephalosporin\*\*
- A tetracycline\*\*
- Sulfamethoxazole-trimethoprim (SMZ-TMP)\*\*
- Clindamycin\*\*

Notes	*Note: Authorization will be issued for up to 14 days. **Drug may require PA
-------	---

Product Name: Baxdela	
Diagnosis	Off-Label Uses*
Guideline Type	Prior Authorization

**Approval Criteria**

1 - For continuation of therapy upon hospital discharge

**OR**

2 - As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

Notes	*Note: Authorization duration based on provider recommended treatment durations, up to 6 months.
-------	--

**2 . Revision History**

Date	Notes
5/12/2021	Arizona Medicaid 7.1 Implementation

Belbuca, Butrans - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117161
<b>Guideline Name</b>	Belbuca, Butrans - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Brand Belbuca, Brand Butrans, generic buprenorphine patches *	
Diagnosis	Cancer/Hospice/End of Life related pain
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - The patient is being treated for cancer, hospice, or end of life related pain	

**AND**

**2** - If the request is for Belbuca or generic Butrans BOTH of the following:

**2.1** Prescriber attests the information provided is true and accurate to the best of their knowledge and they understand that a routine audit may be performed; and medical information necessary to verify the accuracy of the information provided may be requested

**AND**

**2.2** The patient has a history of failure, contraindication or intolerance to BRAND Butrans

Notes	* If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 60-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. If the member is currently taking the requested long-acting opioid for at least 30 days and has met the medical necessity authorization criteria requirements for treatment with an opioid, but has not tried brand buprenorphine patches a denial should be issued and a maximum 60-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. Additionally, a 12 month authorization should be entered for brand buprenorphine patches.
-------	---

Product Name: Brand Belbuca, Brand Butrans, generic buprenorphine patches	
Diagnosis	Cancer/Hospice/End of Life related pain
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The patient is being treated for cancer, hospice, or end of life related pain (Document diagnosis and date of diagnosis)	

**AND**

**2** - If the request is for Belbuca or generic Butrans ONLY: Prescriber attests the information provided is true and accurate to the best of their knowledge and they understand that a routine audit may be performed; and medical information necessary to verify the accuracy of the information provided may be requested

Product Name: Brand Belbuca, Brand Butrans, generic buprenorphine patches *	
Diagnosis	Non-cancer pain/Non-hospice/Non-end of life care pain
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Prescriber attests to ALL of the following:</p> <p><b>1.1</b> The information provided is true and accurate to the best of their knowledge and they understand that a routine audit may be performed; and medical information necessary to verify the accuracy of the information provided may be requested</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> Treatment goals are defined, including estimated duration of treatment</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.3</b> Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.4</b> Patient has been screened for substance abuse/opioid dependence</p>	

**AND**

**1.5** If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression

**AND**

**1.6** Pain is moderate to severe and expected to persist for an extended period of time

**AND**

**1.7** Pain is chronic

**AND**

**1.8** Pain is not postoperative (unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time)

**AND**

**1.9** Pain management is required around the clock with a long-acting opioid

**AND**

**2** - The patient has a history of failure, contraindication, or intolerance to a trial of tramadol IR (immediate release), unless the patient is already receiving chronic opioid therapy prior to surgery for postoperative pain, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time (Drug may require PA)

**AND**

**3** - If the request is for neuropathic pain (examples of neuropathic pain include neuralgias, neuropathies, fibromyalgia), BOTH of the following must be met:

**3.1** Unless it is contraindicated, the patient has not exhibited an adequate response to 8 weeks of treatment with gabapentin or pregabalin (Lyrica) titrated to a therapeutic dose (document date of trial)

**AND**

**3.2** Unless it is contraindicated, the patient has not exhibited an adequate response to at least 6 weeks of treatment with a tricyclic antidepressant titrated to the maximum tolerated dose (document drug and date of trial)

**AND**

**4** - If the request is for Belbuca or generic Butrans, the patient has a history of failure, contraindication or intolerance to BRAND Butrans

Notes	* If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 60-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. If the member is currently taking the requested long-acting opioid for at least 30 days and has met the medical necessity authorization criteria requirements for treatment with an opioid, but has not tried brand buprenorphine patches a denial should be issued and a maximum 60-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. Additionally, a 6 month authorization should be entered for brand buprenorphine patches.
-------	--

Product Name: Brand Belbuca, Brand Butrans, generic buprenorphine patches *	
Diagnosis	Non-cancer pain/Non-hospice/Non-end of life care pain
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Patient demonstrates meaningful improvement in pain and function (document improvement in function or pain score improvement)

**AND**

**2** - Identify rationale for not tapering and discontinuing opioid (document rationale)

**AND**

**3** - Prescriber attests to ALL of the following:

**3.1** The information provided is true and accurate to the best of their knowledge and they understand that a routine audit may be performed; and medical information necessary to verify the accuracy of the information provided may be requested

**AND**

**3.2** Treatment goals are defined, including estimated duration of treatment

**AND**

**3.3** Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention

**AND**

**3.4** Patient has been screened for substance abuse/opioid dependence

**AND**

**3.5** If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression

**AND**

**3.6** Pain is moderate to severe and expected to persist for an extended period of time

**AND**

**3.7** Pain is chronic

**AND**

**3.8** Pain is not postoperative (unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time)

**AND**

**3.9** Pain management is required around the clock with a long-acting opioid

**AND**

**4** - If the request is for Belbuca or generic Butrans, the patient has a history of failure, contraindication, or intolerance to BRAND Butrans

Notes

\* If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 60-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. If the member is currently taking the requested long-acting opioid for at least 30 days and has met the medical necessity authorization criteria requirements for treatment with an opioid, but has not tried brand buprenorphine patches a denial should be issued and a maximum 60-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. Additionally, a 6 month authorization should be entered for brand buprenorphine patches.

Product Name: Brand Belbuca, Brand Butrans, generic buprenorphine patches *	
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - The requested dose cannot be achieved by moving to a higher strength of the product</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - The requested dose is within the FDA (Food and Drug Administration) maximum dose per day, where an FDA maximum dose per day exists</p>	
Notes	*Approval durations: 12 months for cancer pain/hospice/end of life related pain; 6 months for non-cancer pain/non-hospice/non-end of life related pain

## 2 . Revision History

Date	Notes
11/21/2022	Added pregabalin as prerequisite option for neuropathic/nerve pain

Benlysta (belimumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114489
<b>Guideline Name</b>	Benlysta (belimumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Benlysta IV, Benlysta SQ	
Diagnosis	Systemic Lupus Erythematosus
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of systemic lupus erythematosus	

**AND**

**2** - Patient is 5 years of age or older

**AND**

**3** - Laboratory testing has documented the presence of autoantibodies [e.g., ANA, Anti-dsDNA, Anti-Sm, Anti-Ro/SSA, Anti-La/SSB]

**AND**

**4** - Patient is currently receiving standard immunosuppressive therapy [e.g., hydroxychloroquine, chloroquine, prednisone, azathioprine, methotrexate]

**AND**

**5** - Patient does NOT have severe active central nervous system lupus

**AND**

**6** - Patient is not receiving Benlysta in combination with a biologic DMARD (disease modifying anti-rheumatic drug) [e.g., Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Kineret (anakinra)]

Product Name: Benlysta IV, Benlysta SQ	
Diagnosis	Active Lupus Nephritis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of active lupus nephritis

**AND**

2 - Patient is 5 years of age or older

**AND**

3 - Patient is currently receiving standard immunosuppressive therapy for systemic lupus erythematosus [e.g., hydroxychloroquine, chloroquine, prednisone, azathioprine, methotrexate]

**AND**

4 - Patient does NOT have severe active central nervous system lupus

**AND**

5 - Patient is not receiving Benlysta in combination with a biologic DMARD (disease modifying anti-rheumatic drug) [e.g., Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Kineret (anakinra)]

Product Name: Benlysta IV, Benlysta SQ	
Diagnosis	Systemic Lupus Erythematosus, Active Lupus Nephritis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Benlysta therapy	

**AND**

**2** - Patient is not receiving Benlysta in combination with a biologic DMARD (disease modifying anti-rheumatic drug) [e.g., Enbrel (etanercept), Humira (adalimumab), Cimzia (certolizumab), Kineret (anakinra)]

## **2 . Revision History**

Date	Notes
9/26/2022	Updated age requirement. Added IV formulation as target.

Benznidazole

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99434
<b>Guideline Name</b>	Benznidazole
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name: Benznidazole	
Diagnosis	Chagas disease (American trypanosomiasis)
Approval Length	60 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Chagas disease (American trypanosomiasis) due to Trypanosoma cruzi	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copy guidelines starting with B and C from C&S Arizona to Arizona Medicaid

Bevacizumab Products

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325189
<b>Guideline Name</b>	Bevacizumab Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Preferred: Mvasi, Zirabev	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The drug is being used as indicated by National Comprehensive Cancer Network (NCCN) guidelines with a Category of Evidence and Consensus of 1, 2A, or 2B	

Product Name:Non-Preferred*: Alymsys, Avastin, Avzivi, Vegzelma, and newly launched bevacizumab products
--

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred products are Mvasi and Zirabev. **Please note: The drug(s) listed above may require additional review.</p>	
Notes	*Patients must use preferred bevacizumab biosimilar(s).

## 2 . Revision History

Date	Notes
7/16/2025	New program, updated NP section verbiage.

Bimzelx (bimekizumab-bkzx)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-335200
<b>Guideline Name</b>	Bimzelx (bimekizumab-bkzx)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Bimzelx	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** One of the following:

- At least 3% body surface area (BSA) involvement
- Severe scalp psoriasis
- Palmoplantar (i.e., palms, soles), facial, or genital involvement

**AND**

**1.3** Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies:

- corticosteroids (e.g., betamethasone, clobetasol)
- vitamin D analogs (e.g., calcitriol, calcipotriene)
- tazarotene
- calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies:

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab
- Otezla (apremilast)

- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a dermatologist

Product Name:Bimzelx	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by ONE of the following:</p> <ul style="list-style-type: none"> <li>• Reduction the body surface area (BSA) involvement from baseline</li> <li>• Improvement in symptoms (e.g., pruritus, inflammation) from baseline</li> </ul> <p><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a dermatologist</p>	

Product Name:Bimzelx	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

1.1 Diagnosis of active psoriatic arthritis (PsA)

**AND**

1.2 History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

1.3 History of failure, contraindication, or intolerance to ALL of the following:

- Enbrel (etanercept) or a preferred adalimumab biosimilar
- infliximab
- Orencia (abatacept)
- Otezla (apremilast)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

2 - Prescribed by or in consultation with one of the following:

- Dermatologist
- Rheumatologist

Product Name: Bimzelx	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by ONE of the following:

- Reduction in the total active (swollen and tender) joint count from baseline
- Improvement in symptoms (e.g., pain, stiffness, pruritus, inflammation) from baseline
- Reduction in the body surface area (BSA) involvement from baseline

**AND**

2 - Prescribed by or in consultation with one of the following:

- Dermatologist
- Rheumatologist

Product Name: Bimzelx	
Diagnosis	Non-radiographic Axial Spondyloarthritis (nr-AxSpA)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:	
1.1 Diagnosis of active non-radiographic axial spondyloarthritis	
<b>AND</b>	
1.2 Patient has objective signs of inflammation (e.g., C-reactive protein [CRP] levels above the upper limit of normal and/or sacroiliitis on magnetic resonance imaging [MRI], indicative of inflammatory disease, but without definitive radiographic evidence of structural damage on sacroiliac joints)	

**AND**

**1.3** History of a minimum duration of one month trial, contraindication, or intolerance to two different nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally tolerated doses

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name: Bimzelx

Diagnosis	Non-radiographic Axial Spondyloarthritis (nr-AxSpA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by improvement from baseline for at least one of the following:

- Disease activity (e.g., pain, fatigue, inflammation, stiffness)
- Function
- Lab values (erythrocyte sedimentation rate, C-reactive protein level)
- Axial status (e.g., lumbar spine motion, chest expansion)
- Total active (swollen and tender) joint count

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name: Bimzelx

Diagnosis	Ankylosing Spondylitis (AS)
-----------	-----------------------------

Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:</p> <p>1.1 Diagnosis of active ankylosing spondylitis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 History of a minimum duration of one month trial, contraindication, or intolerance to two different nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally tolerated doses</p> <p style="text-align: center;"><b>AND</b></p> <p>1.3 History of failure, contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"> <li>• A preferred adalimumab biosimilar or Enbrel (etanercept)</li> <li>• infliximab</li> <li>• Xeljanz (tofacitinib) oral tablet (IR or XR)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p>	

Product Name: Bimzelx	
Diagnosis	Ankylosing Spondylitis (AS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by improvement from baseline for at least one of the following:

- Disease activity (e.g., pain, fatigue, inflammation, stiffness)
- Function
- Lab values (erythrocyte sedimentation rate, C-reactive protein level)
- Axial status (e.g., lumbar spine motion, chest expansion)
- Total active (swollen and tender) joint count

**AND**

2 - Prescribed by or in consultation with a rheumatologist

Product Name: Bimzelx	
Diagnosis	Hidradenitis Suppurativa (HS)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderate to severe hidradenitis suppurativa (i.e., Hurley Stage II or III)	
<b>AND</b>	
2 - Paid claims or submission of medical records (e.g., chart notes) confirming trial and failure, contraindication, or intolerance to a preferred adalimumab biosimilar	
<b>AND</b>	

**3** - Prescribed by or in consultation with a dermatologist

Product Name: Bimzelx

Diagnosis | Hidradenitis Suppurativa (HS)

Approval Length | 12 month(s)

Therapy Stage | Reauthorization

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by at least one of the following:

- Reduction in the abscess and inflammatory nodule count from baseline
- Reduced formation of new sinus tracts and scarring
- Improvement in symptoms (e.g., pain, suppuration) from baseline

**AND**

**2** - Prescribed by or in consultation with a dermatologist

**2 . Revision History**

Date	Notes
7/17/2025	Updated preferred agents/embedded steps, updated criteria throughout. PsA and AS: Xeljanz to say IR or XR, removed old verbiage from embedded step. Updated submission of records verbiage throughout.

Blood Glucose Monitors

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-99566
<b>Guideline Name</b>	Blood Glucose Monitors
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Non-preferred Blood Glucose Monitors*	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Patient is visually impaired	
Notes	*Please reference background table for list of Non-preferred Blood Glucose Monitors **Approve Glucose Monitor at NDC Level

## 2 . Background

Benefit/Coverage/Program Information			
<b>Non-preferred Blood Glucose Monitors*</b>			
CONTOUR KIT NEXT LNK	EASY TOUCH KIT MONITOR	EASYMAX V KIT SYSTEM	
CONTOUR NXT KIT LINK 2.4	KROGER BGM KIT SYSTEM	EASYMAX NG KIT SYSTEM	
CONTOUR KIT NEXT EZ	ELEMENT AUTO KIT SYSTEM	MEIJER BGM KIT ESSENTIA	
CONTOUR KIT NEXT	SMARTEST KIT EJECT	MEIJER GLUCO KIT MONITOR	
CONTOUR KIT MONITOR	SMARTEST KIT PROTEGE	MEIJER BGM KIT PREMIUM	
RELION MICRO KIT	SMARTEST KIT PRONTO	FORA V30A KIT	
RELION KIT MONITOR	SMARTEST KIT PERSONA	FORA TN'G KIT VOICE	
BD LOGIC KIT MONITOR	GLUCOCOM KIT MONITOR	REFUAH PLUS KIT SYSTEM	
BD LATITUDE KIT	RIGHTEST SYS KIT GM300	KROGER BGM KIT	
BD LATITUDE KIT SYSTEM	RIGHTEST SYS KIT GM100	KROGER BGM KIT PREMIUM	
QUICKTEK KIT	RIGHTEST SYS KIT GM550	CONTOUR KIT LINK 2.4	
ADVANCE KIT INTUITIO	IGLUPOSE KIT	EASYMAX V KIT SYSTEM	
GLUCOCARD KIT SHNE CON	NOVA MAX KIT SYSTEM	EASYMAX NG KIT SYSTEM	
GLUCOCARD KIT SHNE EXP	WAVESENSE KIT KEYNOTE	MYGLUCOHEALT KIT SYSTEM	

GLUCOCARD KIT EXPRESSI	AGAMA JAZZ KIT WRLSS 2	MICRODOT KIT SYSTEM
POCKETCHEM KIT EZ	AGAMATRIX KIT PRESTO	ONE TOUCH KIT VERIO FL
GLUCOCARD 01 KIT SYSTEM	WAVESENSE KIT AMP	RELION TRUE KIT MET AIR
GLUCOCARD 01 KIT MINI	SOLUS V2 KIT SYSTEM	VERASENS KIT
GLUCOCARD KIT X-METER	COOL MONITOR KIT	INFINITY KIT VOICE
GLUCOCARD KIT VITAL	TRUERESULT KIT MONITOR	OPTIUM KIT BL GLUC
RELION PREMI KIT COMP SYS	TRUERESULT KIT SYSTEM	PRECISION KIT XTRA
SMART SENSE KIT GLUC SYS	MEIJER BGM KIT ESSENTIA	PRECISION KIT LINK
CVS GLUCOSE KIT METER	MEIJER GLUCO KIT MONITOR	BIOTEL CARE KIT SYSTEM
INFINITY KIT SYSTEM	MEIJER BGM KIT PREMIUM	BIOTEL CARE KIT
EASYPRO KIT MONITOR	FORA V30A KIT	FREESTYLE KIT SIDEKICK
EASYPRO PLUS KIT	FORA TN'G KIT VOICE	FREESTYLE KIT FREEDOM
PRODIGY PCKT KIT METER	REFUAH PLUS KIT SYSTEM	KROGER BGM KIT PREMIUM
PRODIGY AUTO KIT MONITOR	KROGER BGM KIT	CONTOUR KIT LINK 2.4
PRODIGY VOIC KIT METER		
PRODIGY KIT NO CODIN		

### 3 . Revision History

Date	Notes
7/12/2021	New Program

Bonjesta and Diclegis

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99436
<b>Guideline Name</b>	Bonjesta and Diclegis
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Bonjesta, Brand Diclegis, generic doxylamine/pyridoxine	
Diagnosis	Nausea and vomiting associated with pregnancy
Approval Length	9 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of nausea and vomiting associated with pregnancy  <b>AND</b>	

**2** - Documented failure or contraindication to lifestyle modifications (e.g., diet, avoidance of triggers)

**AND**

**3** - Documented trial and failure or contraindication to a five day trial of over-the-counter doxylamine taken together with pyridoxine (i.e., not a combined dosage form, but separate formulations taken concomitantly)

## **2 . Revision History**

Date	Notes
3/10/2021	Bulk Copy guidelines starting with B and C from C&S Arizona to Arizona Medicaid

Brand Over Generic Not Covered

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99590
<b>Guideline Name</b>	Brand Over Generic Not Covered
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Generic products on a brand* over generic program	
Guideline Type	Administrative
<b>Approval Criteria</b>  1 - Requests for a generic product on a brand over generic program (presence of Brand over generic-Not Covered clinical program in formulary lookup) shall be denied. The plan's preferred product is the brand medication.	
Notes	* Brand product may require prior authorization.

## 2 . Revision History

Date	Notes
10/29/2021	Changed effective date to 12/1/21

Breast Cancer - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99541
<b>Guideline Name</b>	Breast Cancer - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Arimidex, generic anastrozole	
Diagnosis	Breast Cancer
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 Adjuvant treatment of postmenopausal patients with hormone receptor-positive early breast cancer	

**OR**

**1.2** First-line treatment of postmenopausal patients with hormone receptor-positive or hormone receptor status unknown locally advanced or metastatic breast cancer

**OR**

**1.3** Postmenopausal patients with disease progression following tamoxifen therapy

Product Name: Brand Aromasin, generic exemestane	
Diagnosis	Breast Cancer
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following:  <b>1.1</b> Adjuvant treatment of postmenopausal patients with estrogen receptor-positive early breast cancer who have received 2 to 3 years of tamoxifen and are switched to exemestane for completion of a total of 5 consecutive years of adjuvant hormonal therapy  <b>OR</b>  <b>1.2</b> Treatment of advanced breast cancer in postmenopausal patients whose disease has progressed following tamoxifen therapy	

Product Name: Brand Fareston, generic toremifene	
Diagnosis	Breast Cancer
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Treatment of metastatic breast cancer in postmenopausal patients with estrogen receptor positive tumors or with tumors of unknown estrogen receptor status

Product Name: Brand Arimidex, generic anastrozole, Brand Aromasin, generic exemestane, Brand Fareston, generic toremifene

Diagnosis	National Comprehensive Cancer Network (NCCN) Recommended Regimens
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Use supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.

Product Name: Brand Arimidex, generic anastrozole, Brand Aromasin, generic exemestane, Brand Fareston, generic toremifene

Diagnosis	National Comprehensive Cancer Network (NCCN) Recommended Regimens
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**2 . Revision History**

Date	Notes
6/3/2021	Arizona Medicaid 7.1 Implementation

Breo Ellipta

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-126227
<b>Guideline Name</b>	Breo Ellipta
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Brand Breo Ellipta, generic fluticasone-vilanterol	
Diagnosis	Asthma, COPD
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - All of the following:  1.1 Diagnosis of asthma	

**AND**

**1.2** Patient is 5 years of age or older

**AND**

**1.3** The patient has a history of failure, contraindication, or intolerance to treatment with ALL of the following preferred products:

- Advair Diskus (brand) or Advair HFA
- Dulera
- Symbicort

**OR**

**2** - All of the following:

**2.1** Diagnosis of chronic obstructive pulmonary disease (COPD)

**AND**

**2.2** Patient is 18 years of age or older

**AND**

**2.3** One of the following:

**2.3.1** History of failure, contraindication, or intolerance to treatment with at least a 30 day trial of an orally inhaled anticholinergic agent (e.g. Spiriva, Atrovent, Combivent, Tudorza)

**OR**

**2.3.2** History of failure, contraindication, or intolerance to treatment with at least a 30 day trial of an orally inhaled anticholinergic agent/long-acting beta-agonist combination agent (e.g. Anoro Ellipta, Stiolto Respimat)

**AND**

**2.4** The patient has a history of failure, contraindication, or intolerance to treatment with ALL of the following preferred products:

- Advair Diskus (brand) or Advair HFA
- Dulera
- Symbicort

## **2 . Revision History**

Date	Notes
6/26/2023	Added generic as NP target. Added age requirement criteria.

Brexafemme (ibrexafungerp)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-120435
<b>Guideline Name</b>	Brexafemme (ibrexafungerp)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Brexafemme	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Requested drug is being used for a Food and Drug Administration (FDA)-approved indication  <b>AND</b>	

**2** - Trial and failure, contraindication, or intolerance to both of the following:

- One intravaginal product (e.g., clotrimazole, miconazole, tioconazole, terconazole, boric acid)
- Oral fluconazole for a minimum of 3 days duration

## **2 . Revision History**

Date	Notes
1/24/2023	New program

Breztri, Trelegy Ellipta - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-153489
<b>Guideline Name</b>	Breztri, Trelegy Ellipta - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Trelegy Ellipta	
Diagnosis	Asthma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of asthma	

**AND**

**2** - History of failure, contraindication, or intolerance to treatment with ALL of the following preferred products:

- Advair Diskus (brand) or Advair HFA
- Dulera
- Brand Symbicort

Product Name: Breztri, Trelegy Ellipta

Diagnosis	COPD
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and/or emphysema

**AND**

**2** - History of failure, contraindication, or intolerance to treatment with at least a 30 day trial of both of the following used in combination:

- Stiolto Respimat (tiotropium-olodaterol)
- Flovent HFA (fluticasone propionate)

Product Name: Breztri, Trelegy Ellipta

Diagnosis	Asthma, COPD
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
9/26/2024	Added Breztri as NP target

Brilinta and Effient

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269203
<b>Guideline Name</b>	Brilinta and Effient
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Brilinta, generic ticagrelor	
Diagnosis	Acute coronary syndrome (ACS)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Diagnosis of acute coronary syndrome (ACS) [e.g., unstable angina (UA), non-ST elevation myocardial infarction (NSTEMI) or ST-segment elevation myocardial infarction (STEMI)]	

**OR**

**1.2** The medication is being used to reduce the risk of a first myocardial infarction (MI) or stroke in a patient with coronary artery disease (CAD) at high risk for such events [e.g., type 2 diabetes mellitus, hypertension, dyslipidemia, multi-vessel CAD, obesity, heart failure, current smoker or chronic kidney disease]

**OR**

**1.3** The medication is being used to reduce the risk of stroke in patients with acute ischemic stroke (NIH Stroke Scale score less than or equal to 5) or high-risk transient ischemic attack (TIA)

**AND**

**2** - For generic ticagrelor requests ONLY: history of failure or intolerance to Brand Brilinta

Product Name: Brand Effient, Generic prasugrel	
Diagnosis	Acute coronary syndrome (ACS)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of acute coronary syndrome (ACS) [e.g., unstable angina (UA), non-ST elevation myocardial infarction (NSTEMI) or ST-segment elevation myocardial infarction (STEMI)]</p> <p><b>AND</b></p> <p><b>2</b> - The patient must be managed with percutaneous coronary intervention (PCI)</p>	

## 2 . Revision History

Date	Notes
5/29/2025	Added generic ticagrelor 60mg

Buprenorphine Sublingual Tablet



## Prior Authorization Guideline

<b>Guideline ID</b>	GL-121788
<b>Guideline Name</b>	Buprenorphine Sublingual Tablet
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Buprenorphine Sublingual Tablet	
Approval Length	6 Months*
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of opioid abuse/dependence.</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p>	

**2.1** Patient is pregnant or breastfeeding;\*

**OR**

**2.2** Both of the following:

**2.2.1** Patient had an intolerance or side effect to buprenorphine-naloxone sublingual tablet or film;

**AND**

**2.2.2** Side effects or intolerances to buprenorphine-naloxone sublingual tablet or films were not resolved with a trial of anti-emetics (e.g. ondansetron) or non-opioid analgesics.

**OR**

**2.3** Patient has a contraindication to naloxone.

**OR**

**2.4** Both of the following:

**2.4.1** Patient has a severe allergy to naloxone [e.g., Stevens-Johnson syndrome, DRESS (Drug Rash with Eosinophilia and Systemic Symptoms)];

**AND**

**2.4.2** Provider has submitted a copy of the MedWatch Form 3500 to the Food and Drug Administration documenting the adverse reaction

**AND**

**3** - Patient is not currently on ANY of the following:

- Benzodiazepines (e.g. Alprazolam, Diazepam, Lorazepam)
- Hypnotics (e.g. Temazepam, Rozerem, Zolpidem)

- Opioids (e.g. Oxycodone, Tramadol, Hydrocodone)

**AND**

**4** - Prescriber attests that the Arizona State Board of Pharmacy Controlled Substance Prescription Drug Monitoring Program database has been reviewed and that patient has been warned about the dangers of ingesting concurrent sedating medications

Notes	*Approve for 1 year if pregnant or breastfeeding
-------	--

## 2 . Revision History

Date	Notes
2/27/2023	Removed DATA 2000 criterion

Bylvay (odevixibat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-131952
<b>Guideline Name</b>	Bylvay (odevixibat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Bylvay	
Diagnosis	Progressive Familial Intrahepatic Cholestasis (PFIC)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of progressive familial intrahepatic cholestasis (PFIC) type 1, 2, or 3 confirmed by one of the following: <ul style="list-style-type: none"><li>Diagnostic test (e.g., liver function test, liver ultrasound and biopsy, bile analysis)</li></ul>	

- Genetic Testing

**AND**

**2** - Patient is experiencing both of the following:

- Moderate to severe pruritus
- Patient has a serum bile acid concentration above the upper limit of the normal reference for the reporting laboratory

**AND**

**3** - Patient is 3 months of age or older

**AND**

**4** - Patient has had an inadequate response to at least two of the following treatments used for the relief of pruritus:

- Ursodeoxycholic acid (e.g., Ursodiol)
- Antihistamines (e.g., diphenhydramine, hydroxyzine)
- Rifampin
- Bile acid sequestrants (e.g., Questran, Colestid, Welchol)

**AND**

**5** - Prescribed dose is consistent with FDA-approved package labeling and does not exceed a total daily dose of 6 mg

**AND**

**6** - Prescribed by or in consultation with a hepatologist or gastroenterologist

Product Name:Bylvay	
Diagnosis	Progressive Familial Intrahepatic Cholestasis (PFIC)
Approval Length	12 Months

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., reduced serum bile acids, improved pruritus)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed dose is consistent with FDA-approved package labeling and does not exceed a total daily dose of 6 mg</p>	

Product Name:Bylvay	
Diagnosis	Alagille Syndrome (ALGS)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming both of the following:</p> <p>1.1 Diagnosis of Alagille Syndrome (ALGS)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Molecular genetic testing confirms mutations in the JAG1 or NOTCH2 gene</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is experiencing both of the following:</p>	

- Moderate to severe pruritus
- Patient has a serum bile acid concentration above the upper limit of the normal reference for the reporting laboratory

**AND**

**3** - Patient is 12 months of age or older

**AND**

**4** - Patient has had an inadequate response to at least two of the following treatments used for the relief of pruritus:

- Ursodeoxycholic acid (e.g., Ursodiol)
- Antihistamines (e.g., diphenhydramine, hydroxyzine)
- Rifampin
- Bile acid sequestrants (e.g., Questran, Colestid, Welchol)

**AND**

**5** - Prescribed by or in consultation with a hepatologist or gastroenterologist

Product Name:Bylvay	
Diagnosis	Alagille Syndrome (ALGS)
Approval Length	12 Months
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., reduced bile acids, reduced pruritus severity score)</p>	

## 2 . Revision History

Date	Notes
8/29/2023	Added criteria for new indication Alagille Syndrome

Cablivi

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99601
<b>Guideline Name</b>	Cablivi
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Cablivi	
Diagnosis	Acquired thrombotic thrombocytopenic purpura (aTTP)
Approval Length	2 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP)	

**AND**

**2** - Cablivi was initiated in the inpatient setting in combination with plasma exchange therapy

**AND**

**3** - Cablivi will be used in combination with immunosuppressive therapy (e.g., corticosteroids)

**AND**

**4** - Total treatment duration will be limited to 58 days beyond the last therapeutic plasma exchange

Product Name: Cablivi	
Diagnosis	Acquired thrombotic thrombocytopenic purpura (aTTP)
Approval Length	2 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Request is for a new (different) episode requiring the re-initiation of plasma exchange for the treatment of acquired thrombotic thrombocytopenic purpura (aTTP) (Documentation of date of prior episode and documentation date of new episode required)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona SP to Medicaid Arizona SP for 7/1 eff

Cabotegravir Containing Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-161708
<b>Guideline Name</b>	Cabotegravir Containing Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	1/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Vocabria*, Cabenuva*	
Diagnosis	Treatment of HIV-1 Infection
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - All of the following:  1.1 Diagnosis of HIV-1 infection	

**AND**

**1.2** Patient is 12 years of age or older

**AND**

**1.3** Patient's weight is greater than or equal to 35 kg

**AND**

**1.4** Patient is currently virologically suppressed (HIV-1 RNA less than 50 copies/mL) on a stable, uninterrupted antiretroviral regimen for at least 6 months

**AND**

**1.5** Patient has no history of treatment failure or known/suspected resistance to either cabotegravir or rilpivirine

**AND**

**1.6** Provider attests that patient would benefit from long-acting injectable therapy over standard oral regimens

**AND**

**1.7** Prescribed by or in consultation with a clinician with HIV expertise

**OR**

**2** - For continuation of prior therapy

Notes

\*If patient meets criteria above, please approve both Vocabria and Ca benuva at GPI list "CABOTTEGRPA".

Product Name:Vocabria**, Aprelude**	
Diagnosis	HIV-1 Pre-Exposure Prophylaxis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Requested drug is being used for pre-exposure prophylaxis (PrEP) to reduce the risk of sexually acquired HIV-1 infection</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient's weight is greater than or equal to 35 kg</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Documentation of both of the following U.S. Food and Drug (FDA)-approved test prior to use of Vocabria or Aprelude:</p> <ul style="list-style-type: none"> <li>• Negative HIV-1 antigen/antibody test</li> <li>• Negative HIV-1 RNA assay</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - One of the following:</p> <p>4.1 Trial and failure, contraindication or intolerance to BOTH of the following:</p> <ul style="list-style-type: none"> <li>• emtricitabine-tenofovir (generic Truvada)</li> <li>• Descovy</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>4.2 Submission of medical records (e.g., chart notes) from provider documenting BOTH of the following:</p>	

<ul style="list-style-type: none"> <li>• Patient would benefit from long-acting injectable therapy over standard oral regimens</li> <li>• Patient would be adherent to testing and dosing schedule</li> </ul>	
Notes	**If patient meets criteria above, please approve both Vocabria and Apretude at GPI list "APRETUDEPA"

Product Name:Vocabria**, Apretude**	
Diagnosis	HIV-1 Pre-Exposure Prophylaxis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Provider attests that patient is adherent to the testing appointments and scheduled injections of Apretude</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Documentation of both of the following U.S. Food and Drug (FDA)-approved test prior to each maintenance injection of Apretude for HIV PrEP:</p> <ul style="list-style-type: none"> <li>• Negative HIV-1 antigen/antibody test</li> <li>• Negative HIV-1 RNA assay</li> </ul>	
Notes	**If patient meets criteria above, please approve both Vocabria and Apretude at GPI list "APRETUDEPA"

## 2 . Revision History

Date	Notes
12/6/2024	Changed step through Brand Truvada (now NP) to generic

Calcium/Vitamin D

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99531
<b>Guideline Name</b>	Calcium/Vitamin D
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Provider has submitted lab work documenting a Vitamin D deficiency	
Notes	Calcium carbonate and calcium lactate are covered without the need for prior authorization.

## 2 . Revision History

Date	Notes
5/18/2021	7/1 Implementation

Camzyos (mavacamten)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114157
<b>Guideline Name</b>	Camzyos (mavacamten)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Camzyos	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of obstructive hypertrophic cardiomyopathy (HCM)  <b>AND</b>	

**2** - Patient has New York Heart Association (NYHA) Class II or III symptoms (e.g., shortness of breath, chest pain)

**AND**

**3** - Patient has a left ventricular ejection fraction of greater than or equal to 55%

**AND**

**4** - Patient has valsalva left ventricular outflow tract (LVOT) peak gradient greater than or equal to 50 mmHg at rest or with provocation

**AND**

**5** - Trial and failure, contraindication, or intolerance to both of the following at a maximally tolerated dose:

- non-vasodilating beta blocker (e.g., bisoprolol, propranolol)
- calcium channel blocker (e.g., verapamil, diltiazem)

**AND**

**6** - Prescribed by or in consultation with a cardiologist

Product Name: Camzyos	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy (e.g., improved symptom relief)	

**AND**

**2** - Patient has a left ventricular ejection fraction of greater than or equal to 50%

**AND**

**3** - Prescribed by or in consultation with a cardiologist

## **2 . Revision History**

Date	Notes
9/20/2022	New Program

Carbaglu (carglumic acid)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-104872
<b>Guideline Name</b>	Carbaglu (carglumic acid)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Brand Carbaglu, Generic carglumic acid	
Diagnosis	Acute Hyperammonemia due to N-acetylglutamate Synthase (NAGS) Deficiency
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of acute hyperammonemia due to N-acetylglutamate synthase (NAGS) deficiency	

**AND**

**2** - Medication will be used as adjunctive therapy to other ammonia lowering therapies (e.g., protein restriction, ammonia scavengers, dialysis)

**AND**

**3** - Prescribed by or in consultation with a specialist focused in the treatment of metabolic disorders

Product Name: Brand Carbaglu, Generic carglumic acid

Diagnosis	Acute Hyperammonemia due to Propionic Acidemia (PA) or Methylmalonic Acidemia (MMA)
-----------	---

Approval Length	1 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA)

**AND**

**2** - Medication will be used as adjunctive therapy to other ammonia lowering therapies (e.g. intravenous glucose, insulin, protein restriction, dialysis)

**AND**

**3** - Patient's plasma ammonia level is greater than or equal to 50 micromol/L

**AND**

4 - Medication will be used for a maximum duration of 7 days

**AND**

5 - Prescribed by or in consultation with a specialist focused in the treatment of metabolic disorders

Product Name: Brand Carbaglu, Generic carglumic acid

Diagnosis	Chronic Hyperammonemia due to N-acetylglutamate Synthase (NAGS) Deficiency
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of chronic hyperammonemia due to N-acetylglutamate synthase (NAGS) deficiency

**AND**

2 - NAGS deficiency has been confirmed by genetic/mutational analysis

**AND**

3 - Medication will be used as maintenance therapy

**AND**

4 - Prescribed by or in consultation with a specialist focused in the treatment of metabolic disorders

Product Name: Brand Carbaglu, Generic carglumic acid	
Diagnosis	Chronic Hyperammonemia due to N-acetylglutamate Synthase (NAGS) Deficiency
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting a positive clinical response to therapy (e.g., plasma ammonia level within the normal range)</p>	

## 2 . Revision History

Date	Notes
3/31/2022	New program for Carbaglu, mirrors ORx LOB. Added submission of MR to each section.

Casgevy (exagamglogene autotemcel injection)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144825
<b>Guideline Name</b>	Casgevy (exagamglogene autotemcel injection)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Casgevy	
Diagnosis	Sickle Cell Disease
Approval Length	1 Time Authorization in Lifetime*
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of sickle cell disease (SCD)	

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming patient has genotype  $\beta S/\beta S$ ,  $\beta S/\beta 0$ , or  $\beta S/\beta +$

**AND**

**3** - Patient is 12 years of age or older

**AND**

**4** - Provider attests that patient is clinically stable and eligible to undergo hematopoietic stem cell transplant (HSCT)

**AND**

**5** - Submission of medical records (e.g., chart notes) documenting patient has a history of at least 4 vaso-occlusive events (VOEs) in the past 24 months as defined by one of the following scenarios:

- Acute pain event requiring a visit to a medical facility and administration of pain medications (opioids or intravenous [IV] non-steroidal anti-inflammatory drugs [NSAIDs]) or RBC transfusions
- Acute chest syndrome
- Priapism lasting > 2 hours and requiring a visit to a medical facility
- Splenic sequestration

**AND**

**6** - Submission of medical records (e.g., chart notes) confirming patient has obtained a negative test result for all of the following prior to cell collection:

- Hepatitis B virus (HBV)
- Hepatitis C virus (HCV)
- Human immunodeficiency virus (HIV)

**AND**

**7** - Patient is anticipated to provide an adequate number of cells to meet the minimum recommended dose of  $3 \times 10^6$  CD34+ cells/kg

**AND**

**8** - Patient will receive both of the following:

**8.1** Full myeloablative conditioning with busulfan prior to treatment with Casgevy

**AND**

**8.2** Anti-seizure prophylaxis with agents other than phenytoin prior to initiating busulfan conditioning

**AND**

**9** - Prescriber attests that patient will discontinue disease modifying therapies for sickle cell disease (e.g., hydroxyurea, crizanlizumab, voxelotor) 8 weeks before the planned start of mobilization and conditioning

**AND**

**10** - Both of the following:

- Patient has never received any previous sickle cell gene therapy treatment in their lifetime (i.e., Casgevy, Lyfgenia)
- Patient has never received prior allogeneic transplant

**AND**

**11** - Prescribed by a provider at a SCD Treatment center with expertise in gene therapy

**AND**

**12** - Prescribed by one of the following:

- Hematologist/Oncologist
- Specialist with expertise in the diagnosis and management of sickle cell disease

Notes

\*Per prescribing information, Casgevy is for one-time, single dose intravenous use only

Product Name: Casgevy

Diagnosis

Transfusion-dependent  $\beta$ -thalassemia (TDT)

Approval Length

1 Time Authorization in Lifetime\*

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting a diagnosis of transfusion-dependent  $\beta$ -thalassemia (TDT)

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming presence of a mutation at both alleles of the  $\beta$ -globin gene (i.e.,  $\beta^0/\beta^0$ ,  $\beta^0/\beta^+$ ,  $\beta^+/\beta^+$ ,  $\beta^0/\beta^E$ )

**AND**

**3** - Submission of medical records (e.g., chart notes) confirming ONE of the following:

- Patient has a history of requiring at least 100 mL/kg/year of RBC transfusions in the prior 2 years
- Patient requires 10 units/year of RBC transfusions in the prior 2 years

**AND**

**4** - Patient is 12 years of age or older

**AND**

**5** - Provider attests that patient is clinically stable and eligible to undergo hematopoietic stem cell transplant (HSCT)

**AND**

**6** - Submission of medical records (e.g., chart notes) confirming patient has obtained a negative test result for all of the following prior to cell collection:

- Hepatitis B virus (HBV)
- Hepatitis C virus (HCV)
- Human immunodeficiency virus (HIV)

**AND**

**7** - Patient is anticipated to provide an adequate number of cells to meet the minimum recommended dose of  $3 \times 10^6$  CD34+ cells/kg

**AND**

**8** - Patient does not have any of the following:

- Severely elevated iron in the heart (e.g., patients with cardiac T2\* less than 10 msec by MRI)
- Advanced liver disease
- MRI results of the liver demonstrating liver iron content greater than or equal to 15 mg/g (unless biopsy confirms absence of advanced disease)

**AND**

**9** - Both of the following:

- Iron chelation therapy (e.g., deferoxamine, deferasirox) will be discontinued for at least 7 days prior to initiating myeloablative conditioning therapy
- Hydroxyurea, Oxbrta (voxelator), and Adakveo (crizanlizumab) will be discontinued at least 8 weeks prior to start of mobilization and conditioning

**AND**

**10** - Patient will receive both of the following:

**10.1** Full myeloablative conditioning with busulfan prior to treatment with Casgevy

**AND**

**10.2** Anti-seizure prophylaxis with agents other than phenytoin prior to initiating busulfan conditioning

**AND**

**11** - Both of the following:

- Patient has never received any previous transfusion dependent beta-thalassemia gene therapy treatment in their lifetime (i.e., Casgevy, Zynteglo)
- Patient has never received prior allogeneic transplant

**AND**

**12** - Prescribed by a provider at a treatment center with expertise in gene therapy

**AND**

**13** - Prescribed by one of the following:

- Hematologist/Oncologist
- Stem transplant specialist

Notes

\*Per prescribing information, Casgevy is for one-time, single dose intravenous use only

## 2 . Revision History

Date	Notes
3/25/2024	Added criteria for new indication of beta thalassemia

Cayston

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99603
<b>Guideline Name</b>	Cayston
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Cayston	
Diagnosis	Cystic Fibrosis (CF)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystic fibrosis (CF)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona SP to Medicaid Arizona SP for 7/1 eff

CGRP Inhibitors - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157013
<b>Guideline Name</b>	CGRP Inhibitors - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/4/2024
-----------------	-----------

## 1 . Criteria

Product Name:Aimovig, Emgality 120 mg/ml	
Diagnosis	Preventive Treatment of Migraine
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following:  1.1 Both of the following:	

**1.1.1** Diagnosis of episodic migraines

**AND**

**1.1.2** Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month

**OR**

**1.2** All of the following:

**1.2.1** Diagnosis of chronic migraines

**AND**

**1.2.2** Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months

**AND**

**1.2.3** Medication overuse headache has been considered and potentially offending medication(s) have been discontinued

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Two of the following:

**3.1** One of the following:

- History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine)

- Patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine)

**OR**

**3.2** One of the following:

- History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate)
- Patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate)

**OR**

**3.3** One of the following:

- History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol
- Patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, metoprolol

**AND**

**4** - Prescribed by or in consultation with one of the following specialists:

- Neurologist
- Pain specialist
- Headache specialist\*

**AND**

**5** - Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines

Notes	*Headache specialists are physicians certified by the United Council of Neurologic Subspecialties (UCNS).
-------	---

Product Name:Aimovig, Emgality 120 mg/ml	
Diagnosis	Preventive Treatment of Migraine
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan, sumatriptan)] has decreased since the start of CGRP therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with one of the following specialists:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain specialist</li> <li>• Headache specialist*</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - For Chronic Migraine only: Patient continues to be monitored for medication overuse headache (MOH)</p> <p style="text-align: center;"><b>AND</b></p> <p>5 - Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines</p>	
Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

Product Name:Emgality 100 mg/mL

Diagnosis	Episodic Cluster Headaches
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of episodic cluster headache</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least three months</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is 18 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with one of the following specialists:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain specialist</li> <li>• Headache specialist*</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>5 - Medication will not be used in combination with another injectable CGRP inhibitor</p>	
Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

Product Name:Emgality 100 mg/mL	
Diagnosis	Episodic Cluster Headaches

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following specialists:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain specialist</li> <li>• Headache specialist*</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Medication will not be used in combination with another injectable CGRP inhibitor</p>	
Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

Product Name:Ajovy, Qulipta, Vyepti	
Diagnosis	Preventive Treatment of Migraine
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Both of the following:</p>	

**1.1.1** Diagnosis of episodic migraines

**AND**

**1.1.2** Patient has 4 to 14 migraine days per month, but no more than 14 headache days per month

**OR**

**1.2** All of the following:

**1.2.1** Diagnosis of chronic migraines

**AND**

**1.2.2** Patient has greater than or equal to 15 headache days per month, of which at least 8 must be migraine days for at least 3 months

**AND**

**1.2.3** Medication overuse headache has been considered and potentially offending medication(s) have been discontinued

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Two of the following:

**3.1** One of the following:

- History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine)

- Patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine)

**OR**

**3.2** One of the following:

- History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate)
- Patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate)

**OR**

**3.3** One of the following:

- History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol
- Patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, metoprolol

**AND**

**4** - Trial and failure, contraindication, or intolerance to ALL of the following:

- Aimovig
- Emgality

**AND**

**5** - Prescribed by or in consultation with one of the following specialists:

- Neurologist
- Pain specialist
- Headache specialist\*

**AND**

**6** - Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines

Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).
-------	---

Product Name:Ajovy, Qulipta, Vyepti	
-------------------------------------	--

Diagnosis	Preventive Treatment of Migraine
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity

**AND**

2 - Use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan, sumatriptan)] has decreased since the start of CGRP therapy

**AND**

3 - Trial and failure, contraindication, or intolerance to ALL of the following:

- Aimovig
- Emgality

**AND**

4 - Prescribed by or in consultation with one of the following specialists:

- Neurologist
- Pain specialist
- Headache specialist\*

**AND**

**5** - For Chronic Migraine only: Patient continues to be monitored for medication overuse headache (MOH)

**AND**

**6** - Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines

Notes

\*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

Product Name:Nurtec ODT

Diagnosis

Preventive Treatment of Episodic Migraine

Approval Length

6 month(s)

Therapy Stage

Initial Authorization

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Both of the following:

**1.1** Diagnosis of episodic migraines

**AND**

**1.2** Patient has 4 to 18 migraine days per month, but no more than 18 headache days per month

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Two of the following:

**3.1** One of the following:

- History of failure (after at least a two month trial) or intolerance to Elavil (amitriptyline) or Effexor (venlafaxine)
- Patient has a contraindication to both Elavil (amitriptyline) and Effexor (venlafaxine)

**OR**

**3.2** One of the following:

- History of failure (after at least a two month trial) or intolerance to Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate)
- Patient has a contraindication to both Depakote/Depakote ER (divalproex sodium) and Topamax (topiramate)

**OR**

**3.3** One of the following:

- History of failure (after at least a two month trial) or intolerance to one of the following beta blockers: atenolol, propranolol, nadolol, timolol, or metoprolol
- Patient has a contraindication to all of the following beta blockers: atenolol, propranolol, nadolol, timolol, metoprolol

**AND**

**4** - Trial and failure, contraindication, or intolerance to ALL of the following:

- Aimovig
- Emgality

**AND**

**5** - Prescribed by or in consultation with one of the following specialists:

- Neurologist
- Pain specialist
- Headache specialist\*

**AND**

**6** - Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines

Notes

\*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

Product Name:Nurtec ODT	
Diagnosis	Preventive Treatment of Episodic Migraine
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient has experienced a positive response to therapy, demonstrated by a reduction in headache frequency and/or intensity</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Use of acute migraine medications [e.g., nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen), triptans (e.g., eletriptan, rizatriptan, sumatriptan)] has decreased since the start of CGRP therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with one of the following specialists:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain specialist</li> </ul>	

- Headache specialist\*

**AND**

**4** - Medication will not be used in combination with another CGRP inhibitor for the preventive treatment of migraines

Notes

\*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

Product Name:Nurtec ODT, Zavzpret

Diagnosis	Acute Treatment of Migraine
-----------	-----------------------------

Approval Length	3 month(s)
-----------------	------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of migraine with or without aura

**AND**

**2** - Will be used for the acute treatment of migraine

**AND**

**3** - Patient has fewer than 15 headache days per month

**AND**

**4** - Patient is 18 years of age or older

**AND**

**5** - Patient has a history of a one-month trial resulting in therapeutic failure, contraindication, or intolerance to FOUR of the following as evidenced by submission of medical records or claims history:

- naratriptan tablets
- rizatriptan tablets/ODT (Oral Disintegrating Tablets)
- sumatriptan auto injection/cartridge
- Imitrex nasal spray (Brand only)
- zolmitriptan tablets/ODT
- Zomig nasal spray (Brand only)

**AND**

**6** - Patient has a history of a one-month trial resulting in therapeutic failure, contraindication, or intolerance to Ubrelyv as evidenced by submission of medical records or claims history\*\*

**AND**

**7** - If patient has 4 or more headache days per month, patient must meet one of the following:

**7.1** Currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications

**OR**

**7.2** Currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications

**OR**

**7.3** Currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications

**AND**

**8** - Prescribed by or in consultation with one of the following specialists:

- Neurologist

- Pain specialist
- Headache specialist\*

**AND**

**9** - Medication will not be used in combination with another oral CGRP inhibitor

Notes	*Headache specialists are physicians certified by the United Council of Neurologic Subspecialties (UCNS). **Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from the manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.
-------	--

Product Name:Nurtec ODT, Zavzpret	
Diagnosis	Acute Treatment of Migraine
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with one of the following specialists:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain specialist</li> <li>• Headache specialist*</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Medication will not be used in combination with another oral CGRP inhibitor</p>	

Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).
-------	---

Product Name:Ubrelvy	
Diagnosis	Acute Treatment of Migraine
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of migraine with or without aura</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Will be used for the acute treatment of migraine</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Will not be used for preventive treatment of migraine</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Patient has fewer than 15 headache days per month</p> <p style="text-align: center;"><b>AND</b></p> <p>5 - Patient is 18 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>6 - Patient has a history of a one-month trial resulting in therapeutic failure, contraindication,</p>
--

or intolerance to TWO of the following as evidenced by submission of medical records or claims history:

- naratriptan tablets
- rizatriptan tablets/ODT (Oral Disintegrating Tablets)
- sumatriptan auto injection/cartridge
- zolmitriptan tablets/ODT
- Zomig nasal spray (Brand only)
- Imitrex nasal spray (Brand only)

**AND**

**7** - If patient has 4 or more headache days per month, patient must meet one of the following:

**7.1** Currently being treated with Elavil (amitriptyline) or Effexor (venlafaxine) unless there is a contraindication or intolerance to these medications

**OR**

**7.2** Currently being treated with Depakote/Depakote ER (divalproex sodium) or Topamax (topiramate) unless there is a contraindication or intolerance to these medications

**OR**

**7.3** Currently being treated with a beta blocker (i.e., atenolol, propranolol, nadolol, timolol, or metoprolol) unless there is a contraindication or intolerance to these medications

**AND**

**8** - Prescribed by or in consultation with one of the following specialists:

- Neurologist
- Pain specialist
- Headache specialist\*

**AND**

**9** - Medication will not be used in combination with another oral CGRP inhibitor

Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).
-------	---

Product Name:Ubrelvy	
Diagnosis	Acute Treatment of Migraine
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has experienced a positive response to therapy (e.g., reduction in pain, photophobia, phonophobia, nausea)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Will not be used for preventive treatment of migraine</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with one of the following specialists:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain specialist</li> <li>• Headache specialist*</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Medication will not be used in combination with another oral CGRP inhibitor</p>	
Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS).

## 2 . Revision History

Date	Notes
10/3/2024	Added embedded step through preferred agents to Ajoyv reauth section.

Cholbam

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99700
<b>Guideline Name</b>	Cholbam
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Cholbam	
Diagnosis	Bile Acid Synthesis Disorder
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of a bile acid synthesis disorder	

**AND**

**2** - It is due to single enzyme defects

Product Name:Cholbam	
Diagnosis	Peroxisomal Disorders Including Zellweger Spectrum Disorders
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of peroxisomal disorders including Zellweger spectrum disorders	
<b>AND</b>	
2 - Patient exhibits manifestations of liver disease, steatorrhea, or complications from decreased fat soluble vitamin absorption	
<b>AND</b>	
3 - It is being used as adjunctive treatment	

Product Name:Cholbam	
Diagnosis	All Indications
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to Cholbam therapy

**2 . Revision History**

Date	Notes
4/10/2021	7/1 Implementation

Cialis for BPH - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-105174
<b>Guideline Name</b>	Cialis for BPH - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Brand Cialis 5mg, generic tadalafil 5mg	
Diagnosis	Benign Prostatic Hyperplasia (BPH)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - All of the following:  1.1 The patient has a diagnosis of benign prostatic hyperplasia (BPH)	

**AND**

**1.2** History of failure, intolerance, or contraindication to BOTH of the following:

- Alpha Blockers (e.g., tamsulosin, alfuzosin ER, doxazosin, or terazosin)
- 5-alpha reductase inhibitors (e.g., finasteride)

**AND**

**1.3** Dose does not exceed 5 milligrams once daily

**AND**

**2** - Provider attests that patient is not using any form of organic nitrate (for example, nitroglycerin, isosorbide dinitrate, isosorbide mononitrate or amyl nitrate) or Adempas

## 2 . Revision History

Date	Notes
3/24/2022	Added physician attestation re: patient not using nitrates

Cibinqo (abrocitinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-141167
<b>Guideline Name</b>	Cibinqo (abrocitinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/7/2024
-----------------	----------

## 1 . Criteria

Product Name:Cibinqo	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of moderate to severe atopic dermatitis	

**AND**

**2** - Submission of medical records documenting one of the following:

- Involvement of at least 10% body surface area (BSA)
- SCORing Atopic Dermatitis (SCORAD) index value of at least 25 [A]

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist/Immunologist

**AND**

**4** - Submission of medical records (e.g., chart notes, lab work, imaging) or paid claims history documenting ALL of the following\*\*:

**4.1** History of failure, contraindication, or intolerance to the following topical therapies: (document drug, date of trial, and/or contraindication to medication)\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole)

**AND**

**4.2** Submission of medical records (e.g., chart notes, lab work, imaging) or paid claims history documenting trial and failure of a minimum 12-week supply of Dupixent (dupilumab) \*\*

**AND**

**4.3** Submission of medical records (e.g., chart notes, lab work, imaging) or paid claims history documenting trial and failure of a minimum 12-week supply of Adbry (tralokinumab-ldrm) \*\*

**AND**

**5** - Not used in combination with biologic immunomodulators (e.g., Dupixent, Adbry) or other immunosuppressants (e.g., azathioprine, cyclosporine)

**AND**

**6** - Patient is 12 years of age or older

Notes

\*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication  
\*\*PA may be required

Product Name: Cibinqo

Approval Length | 12 month(s)

Therapy Stage | Reauthorization

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting a positive clinical response to therapy as evidenced by at least ONE of the following:

- Reduction in body surface area involvement from baseline
- Reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline [A]

**AND**

**2** - Not used in combination with biologic immunomodulators (e.g., Dupixent, Adbry) or other immunosuppressants (e.g., azathioprine, cyclosporine)

**2 . Background**

**Clinical Practice Guidelines**

**Table 1. Relative potencies of topical corticosteroids [2]**

<b>Class</b>	<b>Drug</b>	<b>Dosage Form</b>	<b>Strength (%)</b>
Very high potency	Augmented betamethasone dipropionate	Ointment, gel	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream, lotion	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05
	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment, lotion	0.05

	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream, lotion	0.1
	Triamcinolone acetonide	Cream, ointment, lotion	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

### 3 . Revision History

Date	Notes
2/6/2024	Updated criteria to include submission of records where applicable, added step through Adbry.

Cimzia (certolizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300282
<b>Guideline Name</b>	Cimzia (certolizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Cimzia	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**1.2** History of failure to a 3 month trial of one non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Oencia (abatacept)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred tocilizumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Cimzia	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of active polyarticular juvenile idiopathic arthritis (PJIA)	
<b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:

**2.1** Minimum duration of a 6-week trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:

- methotrexate
- leflunomide

**AND**

**2.2** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Orenzia (abatacept)
- Xeljanz (tofacitinib) oral tablet
- A preferred tocilizumab biosimilar

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Product Name:Cimzia	
Diagnosis	Ankylosing Spondylitis (AS) or Non-Radiographic Axial (nr-axSpA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:	
1.1 One of the following diagnoses:	
<ul style="list-style-type: none"><li>• Active ankylosing spondylitis (AS)</li><li>• Active non-radiographic axial spondyloarthritis (nr-axSpA)</li></ul>	

**AND**

**1.2** History of failure to two NSAIDs (non-steroidal anti-inflammatory drugs; e.g., ibuprofen, naproxen) at maximally indicated doses, each used for at least 4 weeks within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Cimzia	
Diagnosis	Rheumatoid Arthritis (RA), Polyarticular Juvenile Idiopathic Arthritis (PJIA), Ankylosing Spondylitis (AS) or Non-Radiographic Axial (nr-axSpA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a rheumatologist	

Product Name:Cimzia	
Diagnosis	Crohn's Disease (CD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:</p> <p><b>1.1</b> Diagnosis of moderately to severely active Crohn's disease (CD)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> History of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)</li> <li>• 6-mercaptopurine (Purinethol)</li> <li>• Azathioprine (Imuran)</li> <li>• Methotrexate (Rheumatrex, Trexall)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.3</b> History of failure, contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"> <li>• A preferred adalimumab biosimilar</li> <li>• infliximab</li> <li>• A preferred ustekinumab biosimilar</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a gastroenterologist</p>	

Product Name:Cimzia

Diagnosis	Crohn's Disease (CD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a gastroenterologist</p>	

Product Name: Cimzia	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:</p> <p>1.1 Diagnosis of active psoriatic arthritis (PsA)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced</p>	

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Otezla (apremilast)
- Orenzia (abatacept)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Product Name:Cimzia	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with ONE of the following:	
<ul style="list-style-type: none"><li>• Rheumatologist</li></ul>	

- Dermatologist

Product Name:Cimzia	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:</p> <p>1.1 Diagnosis of moderate to severe plaque psoriasis (PsO)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Greater than or equal to 3% body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.3 History of failure to one of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., betamethasone, clobetasol, desonide)</li> <li>• Vitamin D analogs (e.g., calcitriol, calcipotriene)</li> <li>• Tazarotene</li> <li>• Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>1.4 History of failure of a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced</p>	

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies:

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a dermatologist

Product Name:Cimzia	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a dermatologist	

## 2 . Revision History

Date	Notes
7/3/2025	Updated preferred agents/embedded steps, updated criteria through out.

CMV and Herpes Virus Agents- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99518
<b>Guideline Name</b>	CMV and Herpes Virus Agents- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name: Brand Valcyte tabs/oral soln, generic valganciclovir tabs/oral soln, Brand Cytovene inj, generic ganciclovir inj, Foscavir inj	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Medication is being used for ONE of the following:  1.1 Cytomegalovirus (CMV) disease prophylaxis	

**OR**

**1.2** Cytomegalovirus (CMV) retinitis

**OR**

**1.3** Cytomegalovirus (CMV) retinitis prophylaxis

**OR**

**1.4** BOTH of the following:

**1.4.1** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology

**AND**

**1.4.2** The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plan's program\*

Notes	*Note: Medications used solely for anti-obesity/weight loss, cosmetic ( e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs ( Cialis/Tadalafil) are covered for clinical diagnoses other than ED.
-------	--

Product Name:cidofovir inj	
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Medication is being used for ONE of the following:

1.1 Cytomegalovirus (CMV) retinitis

**OR**

1.2 Cytomegalovirus (CMV) retinitis prophylaxis

**OR**

1.3 BOTH of the following:

1.3.1 The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology

**AND**

1.3.2 The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plan's program\*

Notes	*Note: Medications used solely for anti-obesity/weight loss, cosmetic ( e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs ( Cialis/Tadalafil) are covered for clinical diagnoses other than ED.
-------	--

Product Name:famciclovir tabs	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1** - Medication is being used for ONE of the following:

**1.1** Herpes genitalis

**OR**

**1.2** Herpes genitalis prophylaxis

**OR**

**1.3** Herpes labialis

**OR**

**1.4** Herpes simplex virus infection

**OR**

**1.5** Herpes zoster (shingles) infection

**OR**

**1.6** BOTH of the following:

**1.6.1** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology

**AND**

**1.6.2** The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plan's program\*

Notes	*Note: Medications used solely for anti-obesity/weight loss, cosmetic (e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs (Cialis/Tadalafil) are covered for clinical diagnoses other than ED.
-------	--

Product Name: Brand Valtrex tabs, generic valacyclovir tabs

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Medication is being used for ONE of the following:

1.1 Herpes genitalis

**OR**

1.2 Herpes genitalis prophylaxis

**OR**

1.3 Herpes labialis

**OR**

1.4 Herpes simplex virus infection

**OR**

1.5 Herpes zoster (shingles) infection

**OR**

**1.6** Varicella (chicken pox) infection

**OR**

**1.7** BOTH of the following

**1.7.1** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology

**AND**

**1.7.2** The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plan's program\*

Notes	*Note: Medications used solely for anti-obesity/weight loss, cosmetic ( e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs ( Cialis/Tadalafil) are covered for clinical diagnoses other than ED.
-------	--

## 2 . Revision History

Date	Notes
5/13/2021	Arizona Medicaid 7.1 Implementation

Colony Stimulating Factors - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325190
<b>Guideline Name</b>	Colony Stimulating Factors - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Preferred: Fulphila, Fynetra, Nivestym, Releuko	
Diagnosis	Bone Marrow/Stem Cell Transplant
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following:  1.1 Patient has non-myeloid malignancies and is undergoing myeloablative chemotherapy followed by autologous or allogeneic bone marrow transplant (BMT)	

**OR**

**1.2** Used for mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis

**OR**

**1.3** Patient has had a peripheral stem cell transplant (PSCT) and has received myeloablative chemotherapy

**AND**

**2** - Prescribed by, or in consultation with, a hematologist or oncologist

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko

Diagnosis	AML Induction or Consolidation Therapy
-----------	--

Approval Length	3 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of acute myeloid leukemia (AML)

**AND**

**2** - Patient has completed either induction or consolidation chemotherapy

**AND**

**3** - Prescribed by, or in consultation with, a hematologist or oncologist

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko	
Diagnosis	Neutropenia Associated with Cancer Chemotherapy –Dose Dense Chemotherapy
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Patient is receiving National Cancer Institute’s Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 Patient is receiving a dose-dense chemotherapy regimen for which the incidence of febrile neutropenia (FN) is unknown</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by, or in consultation with, a hematologist or oncologist</p>	

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko	
Diagnosis	Primary Prophylaxis of Chemotherapy-Induced Febrile Neutropenia (FN)
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Patient is receiving chemotherapy regimen(s) associated with greater than 20 percent incidence of febrile neutropenia (FN)</p>	

**OR**

**1.2 BOTH** of the following:

- Patient is receiving chemotherapy regimen(s) associated with 10-20 percent incidence of FN
- Patient has one or more risk factors associated with chemotherapy-induced infection, FN, or neutropenia

**AND**

**2** - Prescribed by, or in consultation with, a hematologist or oncologist

Product Name:Preferred: Fulphila, Fylnetra, Nivestym, Releuko

Diagnosis	Secondary Prophylaxis of Febrile Neutropenia (FN)
-----------	---

Approval Length	3 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Patient is receiving myelosuppressive anti-cancer drugs associated with neutropenia (absolute neutrophil count [ANC] less than or equal to 500 cells per mm<sup>3</sup>)

**AND**

**2** - Patient has a history of febrile neutropenia (FN) during a previous course of chemotherapy

**AND**

**3** - Prescribed by, or in consultation with, a hematologist or oncologist

Product Name:Preferred: Fulphila, Fylnetra, Nivestym, Releuko

Diagnosis	Treatment of Febrile Neutropenia (FN) (off-label)
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is receiving myelosuppressive anti-cancer drugs associated with neutropenia (absolute neutrophil count [ANC] less than or equal to 500 cells per mm<sup>3</sup>)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Diagnosis of febrile neutropenia (FN) and patient is considered high risk for infection-associated complications</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by, or in consultation with, a hematologist or oncologist</p>	

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko	
Diagnosis	Severe Chronic Neutropenia (SCN)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of severe chronic neutropenia (SCN) (i.e., congenital, cyclic, and idiopathic neutropenias with chronic absolute neutrophil count [ANC] less than or equal to 500 cells per mm<sup>3</sup>)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by, or in consultation with, a hematologist or oncologist</p>	

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko	
Diagnosis	HIV-Related Neutropenia (off-label)
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of human immunodeficiency virus (HIV) infection</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has an absolute neutrophil count (ANC) less than or equal to 1,000 cells per mm<sup>3</sup></p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by, or in consultation with, ONE of the following:</p> <ul style="list-style-type: none"> <li>• Hematologist</li> <li>• Oncologist</li> <li>• Infectious disease specialist</li> </ul>	

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko	
Diagnosis	Hepatitis C Treatment Related Neutropenia (off-label)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 ALL of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of hepatitis C virus</li> </ul>	

- Patient is undergoing treatment with Peg-Intron (peginterferon alfa-2b) or Pegasys (peginterferon alfa-2a)
- Documentation of neutropenia (absolute neutrophil count [ANC] less than or equal to 500 cells per mm<sup>3</sup>) after dose reduction of Peg-Intron or Pegasys

**OR**

**1.2 BOTH** of the following:

**1.2.1** Documentation of interferon-induced neutropenia (ANC less than or equal to 500 cells per mm<sup>3</sup>) due to treatment with Peg-Intron (peginterferon alfa-2b) or Pegasys (peginterferon alfa-2a)

**AND**

**1.2.2 ONE** of the following:

- Diagnosis of human immunodeficiency virus (HIV) co-infection
- Status post liver transplant
- Diagnosis of established cirrhosis

**AND**

**2** - Prescribed by, or in consultation with, a hematologist, oncologist, gastroenterologist, hepatologist, or infectious disease specialist

Product Name: Preferred: Fulphila, Fylnetra, Nivestym, Releuko	
Diagnosis	Hematopoietic Syndrome of Acute Radiation Syndrome
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has been acutely exposed to myelosuppressive doses of radiation	

**AND**

**2** - Prescribed by, or in consultation with, a hematologist or oncologist

Product Name:Non-Preferred\*: Granix, Neulasta, Neulasta Onpro, Neupogen, Nypozi, Nyvepria, Rolvedon, Ryzneuta, Stimufend, Udenyca, Zarxio, Ziextenzo, and newly launched CSF/biosimilar products

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
-----------------	--

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred products are Fulphila, Fylnetra, Nivestym, and Releuko.  
\*\*Please note: The drug(s) listed above may require additional review.

Notes	*Patients must use preferred colony stimulating factors.
-------	--

**2 . Revision History**

Date	Notes
7/16/2025	Updated preferred agents, updated criteria throughout. Updated NP section verbiage.

Combination Basal Insulin/GLP-1 Receptor Agonist

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99510
<b>Guideline Name</b>	Combination Basal Insulin/GLP-1 Receptor Agonist
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Soliqua	
Approval Length	12 month(s)
Guideline Type	Step Therapy
<b>Approval Criteria</b>  1 - Inadequately controlled on BOTH of the following <ul style="list-style-type: none"><li>GLP-1 (glucagon-like peptide-1) receptor agonist [e.g. Adlyxin (lixisenatide), Trulicity (dulaglutide), Victoza (liraglutide), Bydureon (exenatide extended-release), Byetta (exenatide)]</li><li>Basal insulin (e.g. insulin glargine, insulin degludec, insulin detemir)</li></ul>	

Product Name:Xultophy	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of type 2 diabetes mellitus</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Inadequately controlled on BOTH of the following</p> <ul style="list-style-type: none"> <li>• GLP-1 (glucagon-like peptide-1) receptor agonist [e.g. Adlyxin (lixisenatide), Trulicity (dulaglutide), Victoza (liraglutide), Bydureon (exenatide extended-release), Byetta (exenatide)]</li> <li>• Basal insulin (e.g. insulin glargine, insulin degludec, insulin detemir)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - History of failure, intolerance, or contraindication to Soliqua</p>	

Product Name:Xultophy	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Xultophy therapy</p>	

## 2 . Revision History

Date	Notes
5/24/2021	Arizona Medicaid 7.1 Implementation

Compounds and Bulk Powders

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-139359
<b>Guideline Name</b>	Compounds and Bulk Powders
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Requests for Compounds or Bulk Powders	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Administrative
<b>Approval Criteria</b> 1 - One of the following:  1.1 The compound is an antibiotic.	

**OR**

**1.2** Each active ingredient in the compounded drug is a covered medication

**AND**

**2** - ONE of the following:

**2.1** Each active ingredient in the compounded drug is to be administered for an FDA (Food and Drug Administration)-approved indication

**OR**

**2.2** The use of each active ingredient in the compounded drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**AND**

**3** - If a drug included in the compound requires prior authorization and/or step therapy, all drug specific clinical criteria must also be met

**AND**

**4** - The compounded drug must not include any ingredient that has been withdrawn or removed from the market due to safety reasons.

**AND**

**5** - ONE of the following:

**5.1** A unique vehicle is required for topically administered compounds

**OR**

**5.2** A unique dosage form is required for a commercially available product due to patient's age, weight, or inability to take a solid dosage form

**OR**

**5.3** A unique formulation is required for a commercially available product due to an allergy or intolerance to an inactive ingredient in the commercially available product

**OR**

**5.4** There is a shortage of the commercially available product per the FDA Drug Shortage database or the ASHP Current Drug Shortages tracking log

**AND**

**6** - Coverage for compounds and bulk powders will NOT be approved for any of the following:

**6.1** For topical compound preparations (e.g. creams, ointments, lotions, or gels to be applied to the skin for transdermal, transcutaneous, or any other topical route), requested compound contains any FDA approved ingredient that is not FDA approved for TOPICAL use (see Table 1 in Background section)

**OR**

**6.2** If the requested compound contains topical fluticasone, topical fluticasone will NOT be approved unless both of the following are met:

**6.2.1** Topical fluticasone is intended to treat a dermatologic condition (scar treatments are considered cosmetic and will not be covered)

**AND**

**6.2.2** Patient has a contraindication to all commercially available topical fluticasone formulations

**OR**

**6.3** Requested compound contains any ingredients when used for cosmetic purposes (see Table 2 in Background section)

**OR**

**6.4** Requested compound contains any ingredient(s) which are on the FDA's Do Not Compound List (see Table 3 in Background section)

Product Name:Requests for Compounds or Bulk Powders	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Administrative
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	

## 2 . Background

Benefit/Coverage/Program Information
<b>Table 1: Example topical compound preparations that contain any FDA approved ingredient that are not FDA approved for TOPICAL use, including but NOT LIMITED TO the following:</b>
(1) Ketamine

- (2) Gabapentin
- (3) Flurbiprofen (topical ophthalmic use not included)
- (4) Ketoprofen
- (5) Morphine
- (6) Nabumetone
- (7) Oxycodone
- (8) Cyclobenzaprine
- (9) Baclofen
- (10) Tramadol
- (11) Hydrocodone
- (12) Meloxicam
- (13) Amitriptyline
- (14) Pentoxifylline
- (15) Orphenadrine
- (16) Piroxicam
- (17) Levocetirizine
- (18) Amantadine
- (19) Oxytocin
- (20) Sumatriptan
- (21) Chorionic gonadotropin (human)
- (22) Clomipramine
- (23) Dexamethasone

- (24) Hydromorphone
- (25) Methadone
- (26) Papaverine
- (27) Mefenamic acid
- (28) Promethazine
- (29) Succimer DMSA
- (30) Tizanidine
- (31) Apomorphine
- (32) Carbamazepine
- (33) Ketorolac
- (34) Dimercaptopropane-sulfonate
- (35) Dimercaptosuccinic acid
- (36) Duloxetine
- (37) Fluoxetine
- (38) Bromfenac (topical ophthalmic use not included)
- (39) Nepafenac (topical ophthalmic use not included)

**Table 2: Example compounds that contain ingredients for cosmetic purposes:**

- (1) Hydroquinone
- (2) Acetyl hexapeptide-8
- (3) Tocopheryl Acid Succinate
- (4) PracaSil TM-Plus

- (5) Chrysaderm Day Cream
- (6) Chrysaderm Night Cream
- (7) PCCA Spira-Wash
- (8) Lipopen Ultra
- (9) Versapro
- (10) Fluticasone
- (11) Mometasone
- (12) Halobetasol
- (13) Betamethasone
- (14) Clobetasol
- (15) Triamcinolone
- (16) Minoxidil
- (17) Tretinoin
- (18) Dexamethasone
- (19) Spironolactone
- (20) Cycloserine
- (21) Tamoxifen
- (22) Sermorelin
- (23) Mederma Cream
- (24) PCCA Cosmetic HRT Base
- (25) Sanare Scar Therapy Cream
- (26) Scarcin Cream

- (27) Apothederm
- (28) Stera Cream
- (29) Copasil
- (30) Collagenase
- (31) Arbutin Alpha
- (32) Nourisil
- (33) Freedom Cepapro
- (34) Freedom Silomac Andydrous
- (35) Retinaldehyde
- (36) Apothederm

**Table 3: Example ingredients on the FDA's Do Not Compound List:**

- (1) 3,3',4',5-tetrachlorosalicylanilide
- (2) Adenosine phosphate
- (3) Adrenal cortex
- (4) Alatrofloxacin mesylate
- (5) Aminopyrine
- (6) Astemizole
- (7) Azaribine
- (8) Benoxaprofen
- (9) Bithionol
- (10) Camphorated oil

- (11) Carbetapentane citrate
- (12) Casein, iodinated
- (13) Cerivastatin sodium
- (14) Chlormadinone acetate
- (15) Chloroform
- (16) Cisapride
- (17) Defenfluramine hydrochloride
- (18) Diamthazole dihydrochloride
- (19) Dibromsalan
- (20) Dihydrostreptomycin sulfate
- (21) Dipyrone
- (22) Encainide hydrochloride
- (23) Etreinate
- (24) Fenfluramine hydrochloride
- (25) Flosequinan
- (26) Glycerol, iodinated
- (27) Grepafloxacin
- (28) Mepazine
- (29) Metabromsalan
- (30) Methapyrilene
- (31) Methopholine
- (32) Methoxyflurane

- (33) Mibefradil dihydrochloride
- (34) Nomifensine maleate
- (35) Novobiocin sodium
- (36) Oxyphenisatin acetate
- (37) Oxyphenisatin
- (38) Pemoline
- (39) Pergolide mesylate
- (40) Phenacetin
- (41) Phenformin hydrochloride
- (42) Phenylpropanolamine
- (43) Pipamazine
- (44) Potassium arsenite
- (45) Propoxyphene
- (46) Rapacuronium bromide
- (47) Rofecoxib
- (48) Sibutramine hydrochloride
- (49) Sparteine sulfate
- (50) Sulfadimethoxine
- (51) Sweet spirits of nitre
- (52) Tegaserod maleate
- (53) Temafloxacin hydrochloride
- (54) Terfenadine

- (55) Ticrynafen
- (56) Tribromsalan
- (57) Trichloroethane
- (58) Troglitazone
- (59) Trovafloxacin mesylate:
- (60) Urethane
- (61) Valdecoxib
- (62) Zomepirac sodium

### 3 . Revision History

Date	Notes
1/23/2024	Changed initial approval duration to 6 months and added reauth with 12 month approval duration.

Constipation Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-163779
<b>Guideline Name</b>	Constipation Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Amitiza, generic lubiprostone	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following: 1.1 ONE of the following diagnoses: <ul style="list-style-type: none"><li>Opioid-induced constipation in an adult with chronic, non-cancer pain</li></ul>	

- Opioid-induced constipation in patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation
- Chronic idiopathic constipation

**OR**

**1.2** Both of the following:

- Diagnosis of irritable bowel syndrome with constipation
- Patient was female at birth

**AND**

**2 - BOTH** of the following:

**2.1** Trial and failure, contraindication, or intolerance to an osmotic laxative e.g., (lactulose, polyethylene glycol, sorbitol)

**AND**

**2.2** Trial and failure, contraindication, or intolerance to **ONE** of the following:

- Bulk Forming Laxatives (e.g., psyllium, fiber)
- Stimulant Laxatives (e.g., bisacodyl, senna)

Product Name:lbsrela	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of irritable bowel syndrome with constipation	

**AND**

**2** - History of failure, contraindication or intolerance to BOTH of the following:

- Lactulose
- Polyethylene glycol (Miralax)

**AND**

**3** - History of failure, contraindication or intolerance to ONE of the following:

- Lubiprostone
- Linzess

Product Name:Linzess	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - One of the following:</p> <p><b>1.1</b> Both of the following:</p> <p><b>1.1.1</b> One of the following diagnoses:</p> <ul style="list-style-type: none"><li>• Chronic idiopathic constipation</li><li>• Irritable bowel syndrome with constipation</li></ul> <p><b>AND</b></p> <p><b>1.1.2</b> Patient is greater than or equal to 18 years of age</p>	

**OR**

**1.2** Both of the following (Applies to Linzess 72mcg requests ONLY)

- Diagnosis of functional constipation
- Patient is 6-17 years of age

**AND**

**2** - Both of the following:

**2.1** Trial and failure, contraindication, or intolerance to an osmotic laxative e.g., (lactulose, polyethylene glycol, sorbitol)

**AND**

**2.2** Trial and failure, contraindication, or intolerance to ONE of the following:

- Bulk Forming Laxatives (e.g., psyllium, fiber)
- Stimulant Laxatives (e.g., bisacodyl, senna)

Product Name: Brand Motegrity, generic prucalopride	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic idiopathic constipation	
<b>AND</b>	
2 - Both of the following	

**2.1** History of failure, contraindication or intolerance to BOTH of the following:

- Lactulose
- Polyethylene glycol (Miralax)

**AND**

**2.2** History of failure, contraindication, or intolerance to BOTH of the following:

- Linzess
- Lubiprostone

Product Name: Movantik	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - ONE of the following diagnoses:	
<ul style="list-style-type: none"><li>• Opioid-induced constipation in patients being treated for chronic, non-cancer pain</li><li>• Opioid-induced constipation in patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation</li></ul>	

Product Name: Relistor tablet, Relistor injection, Symproic	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - ONE of the following diagnoses:

- Opioid-induced constipation in patients being treated for chronic, non-cancer pain
- Opioid-induced constipation in patients with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation

**AND**

2 - History of failure, contraindication or intolerance to BOTH of the following:

- Lactulose
- Polyethylene glycol (Miralax)

**AND**

3 - History of failure, contraindication or intolerance to Movantik

**AND**

4 - For Relistor Injection requests ONLY: The patient is not able to swallow oral medications

Product Name:Trulance	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - ONE of the following diagnoses:	
<ul style="list-style-type: none"><li>• Chronic idiopathic constipation</li><li>• Irritable bowel syndrome with constipation</li></ul>	
<b>AND</b>	

2 - Patient is greater than or equal to 18 years of age

Product Name:Zelnorm	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of irritable bowel syndrome with constipation</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient was female at birth</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - History of failure, contraindication or intolerance to BOTH of the following:</p> <ul style="list-style-type: none"><li>• Lactulose</li><li>• Polyethylene glycol (Miralax)</li></ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - History of failure, contraindication or intolerance to ONE of the following:</p> <ul style="list-style-type: none"><li>• Lubiprostone</li><li>• Linzess</li></ul>	

Product Name:Brand Amitiza, generic lubiprostone, Ibsrela, Linzess, Brand Motegrity, generic prucalopride, Movantik, Relistor tablet, Relistor injection, Symproic, Trulance, Zelnorm	
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
1/30/2025	Added generic prucalopride (Motegrity) as NP target

## Continuous Blood Glucose Monitoring Devices (CGM)

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-157624
<b>Guideline Name</b>	Continuous Blood Glucose Monitoring Devices (CGM)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

#### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

### 1 . Criteria

Product Name: PREFERRED Continuous Glucose Monitors, Sensors, and Transmitters: Freestyle Libre receiver, Freestyle Libre 14 receiver/sensor, Freestyle Libre 2 receiver/sensor, Freestyle Libre 2 Plus/2+ sensor/system, Freestyle Libre 3 sensor, Freestyle Libre 3 Plus/3+ sensor/system	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following:	

**1.1** Submission of medical records (e.g., chart notes) documenting member is already established on an integrated closed loop insulin pump system. The current CGM product will be approved\* (NOTE: Members starting on a closed loop insulin pump system will be required to obtain a new PA if they are changing CGM devices)

**OR**

**1.2** Member is insulin dependent as confirmed by paid claims for insulin within the past 60 days and the request is for a Freestyle Libre product (Freestyle Libre products will adjudicate without a prior authorization submission when the member is insulin dependent as confirmed by insulin paid claims in the members PBM profile)

**OR**

**1.3** Submission of medical records (e.g., chart notes, lab results) documenting all of the following:

**1.3.1** One of the following:

**1.3.1.1** Both of the following:

- Diagnosis of Type I or II Diabetes Mellitus
- Frequent insulin adjustments are required based on the results of blood glucose monitoring or CGM testing results and supporting documentation has been submitted by provider

**OR**

**1.3.1.2** One of the following diagnoses:

- Gestational Diabetes
- Hypoglycemia Unawareness (HU) (defined as the onset of neuroglycopenia, low blood glucose in the brain, before the appearance of autonomic warning symptoms, or the failure to sense a significant fall in blood glucose below normal levels) (submission of medical records/supporting documentation is required)
- Documented Postprandial Hyperglycemia (submission of medical records/supporting documentation is required)
- Documented Recurrent Diabetic Ketoacidosis (submission of medical records/supporting documentation is required)

**OR**

**1.3.1.3** Member requires short term use (72 hours) to determine baseline insulin levels prior to insulin pump initiation

**AND**

**1.3.2** One of the following:

- Hemoglobin A1c > 7.0%
- Frequent hypoglycemic episodes as evidenced by submitted chart documentation
- Member has a diagnosis that is not defined by elevated hemoglobin A1c or frequent hypoglycemia (e.g., Gestational Diabetes)

**AND**

**1.3.3** Provider attests member is enrolled or has completed a comprehensive diabetes education program

**AND**

**1.3.4** Member must meet the FDA approved age for the requested product (new products entering the market shall not be approved below the FDA approved age)

Notes	*NOTE: Members starting on a closed loop insulin pump system will be required to obtain a new PA if they are changing CGM devices.  **Third Party Exception Flag must be flipped to = Y for the claim to pay with the PA in place. Please run a trial claim to make sure claim pays with PA  ***Approve Freestyle Libre products at NDC Level – With NDC List AZ MFR3 (see background section for details)
-------	--

Product Name:NONPREFERRED Continuous Glucose Monitors, Sensors, and Transmitters: Dexcom G6 receiver/sensor/transmitter, Dexcom G7 receiver/sensor, Guardian receiver/sensor/transmitter, Enlite sensor, Eversense sensor/transmitter, Minilink transmitter, Minimed 630G Guardian transmitter, Paradigm transmitter	
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p><b>1.1</b> Submission of medical records (e.g., chart notes) documenting member is already established on an integrated closed loop insulin pump system. The current CGM product will be approved* (NOTE: Members starting on a closed loop insulin pump system will be required to obtain a new PA if they are changing CGM devices)</p> <p style="text-align: center;"><b>OR</b></p> <p><b>1.2</b> Requests for a CGM product other than Freestyle Libre requires submission of medical records (e.g., chart notes, lab results) documenting ALL of the following:</p> <p><b>1.2.1</b> Member has tried and failed the Freestyle Libre system (For other AHCCCS Contractors required steps, please refer to Preferred CGM Products table)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2.2</b> One of the following:</p> <p><b>1.2.2.1</b> All of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of Type I or II Diabetes Mellitus</li> <li>• Member is insulin dependent as demonstrated by paid claims within the past 60 days</li> <li>• Frequent insulin adjustments are required based on the results of blood glucose monitoring or CGM testing results and supporting documentation has been submitted by the provider</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>1.2.2.2</b> One of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Gestational Diabetes</li> <li>• Hypoglycemia Unawareness (HU) (defined as the onset of neuroglycopenia, low blood glucose in the brain, before the appearance of autonomic warning symptoms, or the</li> </ul>	

failure to sense a significant fall in blood glucose below normal levels) (submission of medical records/supporting documentation is required)

- Documented Postprandial Hyperglycemia (submission of medical records/supporting documentation is required)
- Documented Recurrent Diabetic Ketoacidosis (submission of medical records/supporting documentation is required)

**OR**

**1.2.2.3** Member requires short term use (72 hours) to determine baseline insulin levels prior to insulin pump initiation

**AND**

**1.2.3** Member must meet the FDA approved age for the requested product (new products entering the market shall not be approved below the FDA approved age)

**AND**

**1.2.4** One of the following:

- Hemoglobin A1c > 7.0%
- Frequent hypoglycemic episodes as evidenced by submitted chart documentation
- Member has a diagnosis that is not defined by elevated hemoglobin A1c or frequent hypoglycemia (e.g., Gestational Diabetes)

**AND**

**1.2.5** Provider attests member is enrolled or has completed a comprehensive diabetes education program

Notes

\*NOTE: Members starting on a closed loop insulin pump system will be required to obtain a new PA if they are changing CGM devices

\*\*Third Party Exception Flag must be flipped to = Y for the claim to pay with the PA in place. Please run a trial claim to make sure claim pays with PA

\*\*\*Approve all NonPreferred CGM products at GPI Level – With GPI List AZMCGMNP (see background section for details)

Product Name: ALL Continuous Glucose Monitors, Sensors, and Transmitters: Freestyle Libre receiver, Freestyle Libre 14 receiver/sensor, Freestyle Libre 2 receiver/sensor, Freestyle Libre 2 Plus/2+ sensor/system, Freestyle Libre 3 sensor, Freestyle Libre 3 Plus/3+ sensor/system, Dexcom G6 receiver/sensor/transmitter, Dexcom G7 receiver/sensor, Guardian receiver/sensor/transmitter, Enlite sensor, Eversense sensor/transmitter, Minilink transmitter, Minimed 630G Guardian transmitter, Paradigm transmitter

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Member is using the same continuous glucose monitoring device on a regular basis as evidenced through the Member's claims history and the providers chart notes

**AND**

2 - Member is adherent to using the device

**AND**

3 - Member has shared the device readings with physician or healthcare professional for review as part of overall diabetes management

Notes	<p>Third Party Exception Flag must be flipped to = Y for the claim to pay with the PA in place. Please run a trial claim to make sure claim pays with PA</p> <p>Approve all Preferred Freestyle Libre products at NDC Level - With NDC List AZMFR3</p> <p>Approve all NonPreferred CGM products at GPI Level – With GPI List AZMCGMNP</p> <p>(see background section for details)</p>
-------	---

Product Name: ALL Continuous Glucose Monitors, Sensors, and Transmitters: Freestyle Libre receiver, Freestyle Libre 14 receiver/sensor, Freestyle Libre 2 receiver/sensor, Freestyle Libre 2 Plus/2+ sensor/system, Freestyle Libre 3 sensor, Freestyle Libre 3 Plus/3+ sensor/system, Dexcom G6 receiver/sensor/transmitter, Dexcom G7 receiver/sensor, Guardian

receiver/sensor/transmitter, Enlite sensor, Eversense sensor/transmitter, Minilink transmitter, Minimed 630G Guardian transmitter, Paradigm transmitter	
Diagnosis	Requests Exceeding Quantity Limit
Approval Length	1 Time(s)
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - Request is for a vacation override</p> <p style="text-align: center;"><b>OR</b></p> <p>2 - If not for a vacation override, requests for additional transmitter/sensor quantities should be denied</p> <ul style="list-style-type: none"> <li>• Dexcom 6 or 7 sensors: The plan covers a maximum of 3 sensors for a 30 day supply. For defective products, please contact Dexcom CARE at 1-888-738-3646 for a replacement.</li> <li>• For FreeStyle Libre 2, 3, or 3 plus/3+ sensors – The plan covers a maximum of 2 sensors for a 28-day supply. For defective products, please contact FreeStyle Libre Customer Support at 1-844-330-5535 for a replacement.</li> <li>• Guardian Sensor 3 or 4 products – The plan covers a maximum of 5 sensors (1box) for a 35-day supply. For defective products, please contact the Guardian Customer Service Center at 1-800-646-4633 for a replacement.</li> </ul>	
Notes	<p>*Requests for additional quantities for purposes other than a vacation override are to be denied, utilize the product specific denial verbiage below. Third Party Exception Flag must be flipped to = Y for the claim to pay with the PA in place. Please run a trial claim to ensure the claim adjudicates with PA Approve at NDC/GPI Level. Denial language:</p> <ul style="list-style-type: none"> <li>• Dexcom 6 or 7 transmitters - The prior authorization request for more than 1 transmitter in 90 days are to be denied. The plan covers a maximum of 1 transmitter for a 90-day supply. If the member has a defective transmitter, please contact Dexcom CARE at 1- 888-738-3646 for a replacement.</li> <li>• Dexcom 6 or 7 sensors - The prior authorization request for more than 3 sensors in 30 days are to be denied. The plan covers a maximum of 3 sensors for a 30-day supply. If the member has a defective sensor, please contact Dexcom CARE at 1-888-738-3646 for a replacement.</li> <li>• Dexcom G6 Receiver - The prior authorization request for more than 1 receiver in 365 days are to be denied. The plan covers a maximum</li> </ul>

	<p>m of 1 transmitter for a 365-day supply. If the member has a defective receiver, please contact Dexcom CARE at 1- 888-738-3646 for a replacement.</p> <p>FreeStyle Libre &amp; FreeStyle Libre 2, 3, or 3 plus/3+ sensors- The prior authorization request for more than 2 sensors, for a 28-day supply, are to be denied.</p> <ul style="list-style-type: none"> <li>• The plan covers a maximum of 2 sensors for a 28-Day supply. If you have a defective a sensor, please contact Abbott’s FreeStyle Libre Customer Support at 1-844-330-5535 for a replacement.</li> </ul> <p>Guardian Sensor 3 or 4 Sensors – The plan covers a maximum of 5 sensors (1 box) for a 35-day supply. For defective products, please contact the Guardian Customer Service Center at 1-800-646-4633 for a replacement.</p>
--	--

**2 . Background**

Benefit/Coverage/Program Information	
<b>Preferred CGM Products</b>	
Health Plan	CGM Step Therapy Requirements
<b>Arizona Complete Health</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>Banner University Family Care</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>Care 1st Health Plan</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>DCS Comprehensive Health Plan</b>	<b>Dexcom G6 &amp; G7 Freestyle Libre 2 &amp; 3</b>
<b>Division of Developmental Disabilities</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>AHCCCS Fee-For-Service American Indian Health Plan</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>Health Choice Arizona</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>Mercy Care</b>	<b>Dexcom G6 &amp; G7 Freestyle Libre 2 &amp; 3</b>

<b>Molina Healthcare</b>	<b>Freestyle Libre 2 &amp; 3</b>
<b>United Community Plan</b>	<b>Dexcom G6 &amp; G7 Freestyle Libre 2 &amp; 3</b>

**NDC List for Preferred CGM Products**

<b>NDC List</b>	<b>NDC</b>	<b>Product Label</b>	<b>GPI</b>	<b>GPI-14 Description</b>
AZMFR3	57599080300	FREESTY LIBR MIS 2 READER	97202012026200	*CONTINUOUS BLOOD GLUCOSE SYSTEM RECEIVER***
AZMFR3	57599000021	FREESTYLE MI S READER	97202012026200	*CONTINUOUS BLOOD GLUCOSE SYSTEM RECEIVER***
AZMFR3	57599000200	FREESTYLE MI S READER	97202012026200	*CONTINUOUS BLOOD GLUCOSE SYSTEM RECEIVER***
AZMFR3	57599082000	FREESTY LIBR MIS 3 READER	97202012046300	*CONTINUOUS BLOOD GLUCOSE SYSTEM RECEIVER***
AZMFR3	57599081800	FREESTY LIBR KIT 3 SENSOR	97202012046300	*CONTINUOUS BLOOD GLUCOSE SYSTEM SENSOR***
AZMFR3	57599084400	FREE LIBRE3 KIT PLUS/SEN	97202012046300	*CONTINUOUS GLUCOSE SYSTEM SENSOR***
AZMFR3	57599083500	FREE LIBRE2 KIT PLUS/SEN	97202012046300	*CONTINUOUS BLOOD GLUCOSE SYSTEM SENSOR***

AZMFR3	57599080000	FREESTY LIBR KIT 2 SENSOR	97202012046300	*CONTINUOUS BLOOD GLUCOSE SYSTEM SENSOR***
AZMFR3	57599000101	FREESTYLE KI T SENSOR	97202012046300	*CONTINUOUS BLOOD GLUCOSE SYSTEM SENSOR***

**GPI Lists for NonPreferred CGM Products**

GPI List	GPI	GPI-14 Description
AZMCGMNP	97202012026200	*CONTINUOUS BLOOD GLUCOSE SYSTEM RECEIVER***
AZMCGMNP	97202012046300	*CONTINUOUS BLOOD GLUCOSE SYSTEM SENSOR***
AZMCGMNP	97202012066300	*CONTINUOUS BLOOD GLUCOSE SYSTEM TRANSMITTER***

Third Party Exception Flag must be flipped to = Y for the claim to pay with the PA in place. Please run a trial claim to make sure claim pays with PA

**Notes**

Third Party Exception Flag must be flipped to = Y for the claim to pay with the PA in place. Please run a trial claim to make sure claim pays with PA

**Coverage Notes:**

AHCCCS Rule R9-22-202 requires that services be cost effective. The corresponding federal regulations are found in 42 CFR Part 447

R9-22-202. General Requirements

B. In addition to other requirements and limitations specified in this Chapter, the following general requirements apply: Only medically necessary, cost effective, and federally reimbursable and state-reimbursable services are covered.

### 3 . Revision History

Date	Notes
10/25/2024	Added Freestyle Libre 2Plus/+ system GPs, updated background tables, updated age criterion verbiage for Preferred products.

Copper Chelating Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-135313
<b>Guideline Name</b>	Copper Chelating Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	11/1/2023
-----------------	-----------

### 1 . Criteria

Product Name:Brand Depen Titratub, generic penicillamine tablets	
Diagnosis	Severe active rheumatoid arthritis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of severe active rheumatoid arthritis	

Product Name:Brand Depen Titratab, generic penicillamine tablets	
Diagnosis	Severe active rheumatoid arthritis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Depen Titratabs therapy</p>	

Product Name:Brand Depen Titratab, generic penicillamine tablets	
Diagnosis	Wilson's disease (i.e., hepatolenticular degeneration), Cystinuria
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient has ONE of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration)</li> <li>• Diagnosis of Cystinuria</li> </ul>	

Product Name:Brand Cuprimine, generic penicillamine capsules	
Diagnosis	Wilson's disease (i.e., hepatolenticular degeneration), Cystinuria, Severe active rheumatoid arthritis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Patient has ONE of the following diagnoses:

- Wilson's disease (i.e., hepatolenticular degeneration)
- Cystinuria
- Severe active rheumatoid arthritis

**AND**

2 - History of failure or intolerance to Depen (penicillamine)

Product Name: Brand Cuprimine, generic penicillamine capsules

Diagnosis	Wilson's disease (i.e., hepatolenticular degeneration), Cystinuria, Severe active rheumatoid arthritis
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to Cuprimine (penicillamine) therapy

Product Name: Brand Syprine, generic trientine, generic Clovique

Diagnosis	Wilson's disease (i.e., hepatolenticular degeneration)
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration)

**AND**

**2** - History of failure, contraindication, or intolerance to Depen (penicillamine) or Cuprimine (penicillamine)

Product Name: Brand Syprine, generic trientine, generic Clovique	
Diagnosis	Wilson's disease (i.e., hepatolenticular degeneration)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Syprine (trientine) therapy	

## 2 . Revision History

Date	Notes
10/23/2023	Added new GPI for trientine

Corlanor

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99441
<b>Guideline Name</b>	Corlanor
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Corlanor	
Diagnosis	Chronic Heart Failure
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Worsening heart failure in a diagnosis of stable, symptomatic chronic (e.g. New York Heart Association (NYHA) class II, III or IV) heart failure	

**AND**

**2** - Patient has a left ventricular ejection fraction (EF) less than or equal to 35%

**AND**

**3** - The patient is in sinus rhythm

**AND**

**4** - Patient has a resting heart rate greater than or equal to 70 beats per minute

**AND**

**5** - ONE of the following:

**5.1** Patient is on maximum tolerated doses of beta blockers (e.g., carvedilol, metoprolol succinate, bisoprolol)

**OR**

**5.2** Patient has a contraindication or intolerance to beta-blocker therapy

Product Name: Corlanor	
Diagnosis	Heart Failure due to Dilated Cardiomyopathy (DCM)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of stable symptomatic heart failure due to dilated cardiomyopathy (DCM)

**AND**

2 - Patient is in sinus rhythm

**AND**

3 - Patient has an elevated heart rate

Product Name: Corlanor	
Diagnosis	Chronic Heart Failure, Heart Failure due to Dilated Cardiomyopathy (DCM)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Corlanor therapy	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copy guidelines starting with B and C from C&S Arizona to Arizona Medicaid

Cosentyx (secukinumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300284
<b>Guideline Name</b>	Cosentyx (secukinumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Cosentyx SC	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:  1.1 Diagnosis of moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3 percent body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** History of failure to ONE of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies:

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab (Janssen manufacturer)
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

**AND**

**2** - Patient is 6 years of age or older

**AND**

**3** - Prescribed by or in consultation with a dermatologist

Product Name:Cosentyx SC	
Diagnosis	Hidradenitis Suppurativa (HS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - BOTH of the following:</p> <p><b>1.1</b> Submission of medical records (e.g., chart notes) confirming a diagnosis of moderate to severe hidradenitis suppurativa (HS)</p> <p><b>AND</b></p> <p><b>1.2</b> Paid claims or submission of medical records (e.g., chart notes) confirming trial and failure, contraindication, or intolerance to a preferred adalimumab biosimilar</p> <p><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a dermatologist</p>	

Product Name:Cosentyx SC	
Diagnosis	Plaque Psoriasis (PsO), Hidradenitis Suppurativa (HS)

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a dermatologist</p>	

Product Name: Cosentyx SC	
Diagnosis	Enthesitis-Related Arthritis (ERA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting BOTH of the following:</p> <p>1.1 Diagnosis of active enthesitis-related arthritis (ERA)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Trial and failure, contraindication, or intolerance to TWO preferred non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is 4 years of age or older</p>	

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Product Name: Cosentyx SC

Diagnosis	Enthesitis-Related Arthritis (ERA)
-----------	------------------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Documentation of a positive clinical response to therapy as evidenced by at least one of the following:

- Reduction in the total active (swollen and tender) joint count from baseline
- Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name: Cosentyx SC, Cosentyx IV

Diagnosis	Psoriatic Arthritis (PsA)
-----------	---------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

**1.1** Diagnosis of active psoriatic arthritis (PsA)

**AND**

**1.2** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab (Janssen manufacturer)
- Orencia (abatacept)
- Otezla (apremilast)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Patient is 2 years of age or older

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Rheumatologist
- Dermatologist

**AND**

**4** - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to self-administered Cosentyx SC (APPLIES TO REQUESTS FOR COSENTYX IV ONLY):

Product Name: Cosentyx SC, Cosentyx IV	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with ONE of the following:</p> <ul style="list-style-type: none"> <li>• Rheumatologist</li> <li>• Dermatologist</li> </ul>	

Product Name: Cosentyx SC, Cosentyx IV	
Diagnosis	Ankylosing Spondylitis (AS), Non-radiographic axial spondyloarthritis (nr-axSpA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:</p> <p>1.1 One of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Active ankylosing spondylitis (AS)</li> </ul>	

- Active non-radiographic axial spondyloarthritis (nr-axSpA)

**AND**

**1.2** History of failure to two NSAIDs (non-steroidal anti-inflammatory drugs) (e.g., ibuprofen, naproxen) at maximally indicated doses, each used for at least 4 weeks within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab (Janssen manufacturer)
- Xeljanz (tofacitinib) oral tablet (IR or XR)

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to self-administered Cosentyx SC (APPLIES TO REQUESTS FOR COSENTYX IV ONLY)

Product Name: Cosentyx SC, Cosentyx IV	
Diagnosis	Ankylosing Spondylitis (AS), Non-radiographic axial spondyloarthritis (nr-axSpA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with a rheumatologist

## 2 . Revision History

Date	Notes
7/3/2025	Updated preferred drugs/embedded steps, updated criteria throughout.

## Cough and Cold Products

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-104889
<b>Guideline Name</b>	Cough and Cold Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/28/2022
-----------------	-----------

## 1 . Criteria

Product Name:Hydromet, generic Tussionex, Z-Tuss AC, Tuzistra XR, Tussicaps, generic Tussionex, M-END PE, Poly-Tussin AC, Capcof, Pro-Red AC, Histex-AC, Maxi-Tuss, generic promethazine w/codeine, generic promethazine-phenylephrine-codeine, Rydex, Mar-Cof BP/Mar-Cof GG, Ninjacof-XG, Coditussin AC/Coditussin DAC, generic guaifenesin-codeine, generic pseudoephedrine w/codeine-guaifenesin, Tuxarin ER	
Diagnosis	Under the Age of 18 Years for Cough and Cold Products
Approval Length	30 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Prescriber attests they are aware of Food and Drug Administration (FDA) labeled contraindications regarding use of opioid containing cough and cold products in patients less	

than 18 years of age and feels the treatment with the requested product is medically necessary (Document rationale for use)

**AND**

**2** - Patient does not have a comorbid condition that may impact respiratory depression (e.g., asthma or other chronic lung disease, sleep apnea, body mass index greater than 30)

**AND**

**3** - Patient has tried and failed at least one non-opioid containing cough and cold remedy

Product Name:Hydromet, generic Tussionex, Z-Tuss AC, Tuzistra XR, Tussicaps, generic Tussionex, M-END PE, Poly-Tussin AC, Capcof, Pro-Red AC, Histex-AC, Maxi-Tuss, generic promethazine w/codeine, generic promethazine-phenylephrine-codeine, Rydex, Mar-Cof BP/Mar-Cof GG, Ninjacof-XG, Coditussin AC/Coditussin DAC, generic guaifenesin-codeine, generic pseudoephedrine w/codeine-guaifenesin, Tuxarin ER

Diagnosis	Quantity Limit
Approval Length	30 Day(s)
Guideline Type	Quantity Limit*

**Approval Criteria**

**1** - Prescriber attests that a larger quantity is medically necessary

**AND**

**2** - The requested dose is within the Food and Drug Administration (FDA) maximum dose per day, where an FDA maximum dose per day exists (See table in background section)

Notes	*Authorization will be issued for up to 30 days. The authorization should be entered for the quantity requested.
-------	--

**2 . Background**

**Benefit/Coverage/Program Information**

**CDC Recommended Opioid Maximum Morphine Milligram Equivalents per Day\***

Active Ingredient	FDA Label Max Daily Doses
Morphine	None
Hydromorphone	None
Hydrocodone	None
Tapentadol	600mg IR products
Oxymorphone	None
Oxycodone	None
Codeine	360mg
Pentazocine	None
Tramadol	400mg IR products
Meperidine	600mg
Butorphanol nasal	None
Opium	4 suppositories/day  Deodorized tincture: 24mg/day Camphorated tincture: 16mg/day
Acetaminophen	4g/day
Aspirin	2080mg/day
Ibuprofen	3200mg/day
Benzhydrocodone**	None

**3 . Revision History**

Date	Notes
3/28/2022	Updated product list, no changes to criteria.



## Coverage of Off-Label Non-FDA Approved Indications

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164857
<b>Guideline Name</b>	Coverage of Off-Label Non-FDA Approved Indications
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2025
-----------------	----------

## 1 . Criteria

Product Name:A drug (non-anti-cancer chemotherapeutic regimen) used for an off-label indication or non-FDA approved indication	
Diagnosis	Off-label non-cancer indication
Approval Length	12 month(s)
Guideline Type	Administrative
<b>Approval Criteria</b>  1 - The use of this drug is supported by information from ONE of the following appropriate compendia of current literature: <ul style="list-style-type: none"><li>Published practice guidelines and treatment protocols</li></ul>	

- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmaco-economic studies
- Other drug reference resources

**AND**

**2** - Per the AHCCCS Medical Policy Manual- Policy 310-V - Prescription Medications / Pharmacy Services, Section III 1, The Contractor and FFS Programs shall approve the Preferred Drugs listed for the therapeutic classes contained on the AHCCCS Drug List, as appropriate, before approving a Non-Preferred Drug unless ONE of the following applies:

**2.1** Submission of medical records (e.g., chart notes) or paid claims confirming history of trial resulting in a therapeutic failure, contraindication, or intolerance to at least THREE preferred alternatives (if available) (Prior trials of formulary/preferred drug list (PDL) alternatives must sufficiently demonstrate that the formulary/PDL alternatives are either ineffective or inappropriate at the time of the request)

**OR**

**2.2** Submission of medical records (e.g., chart notes) from prescribing clinician supporting the medical necessity of the Non-Preferred Drug over the Preferred Drug for the patient

Notes	Off-label use may be reviewed for medical necessity and denied as such if the off-label criteria are not met. Please refer to drug specific PA guideline for off-label criteria if available.
-------	---

Product Name: A drug or biological in an anti-cancer chemotherapeutic regimen	
Diagnosis	Off-label cancer indication
Approval Length	12 month(s)
Guideline Type	Administrative

## **Approval Criteria**

**1** - One of the following:

**1.1** Diagnosis is supported as a use in AHFS DI

**OR**

**1.2** Diagnosis is supported as a use in the National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B (see NCCN Categories of Evidence and Consensus table in Background section)

**OR**

**1.3** Diagnosis is supported in the FDA Uses/Non-FDA Uses section in DRUGDEX Evaluation with a Strength of Recommendation rating of Class I, Class IIa, or Class IIb (see DRUGDEX Strength of Recommendation table in Background section)

**OR**

**1.4** Diagnosis is supported as an indication in Clinical Pharmacology

**OR**

**1.5** Off-label use is supported in one of the published, peer-reviewed medical literature listed below:

- American Journal of Medicine
- Annals of Internal Medicine
- Annals of Oncology
- Annals of Surgical Oncology
- Biology of Blood and Marrow Transplantation
- Blood
- Bone Marrow Transplantation
- British Journal of Cancer
- British Journal of Hematology
- British Medical Journal
- Cancer
- Clinical Cancer Research
- Drugs

- European Journal of Cancer (formerly the European Journal of Cancer and Clinical Oncology)
- Gynecologic Oncology
- International Journal of Radiation, Oncology, Biology, and Physics
- The Journal of the American Medical Association
- Journal of Clinical Oncology
- Journal of the National Cancer Institute
- Journal of the National Comprehensive Cancer Network (NCCN)
- Journal of Urology
- Lancet
- Lancet Oncology
- Leukemia
- The New England Journal of Medicine
- Radiation Oncology

**OR**

**1.6** Diagnosis is supported as a use in Wolters Kluwer Lexi-Drugs rated as "Evidence Level A" with a "Strong" recommendation. (see Lexi-Drugs Strength of Recommendation table in Background section)

**AND**

**2 -** Per the AHCCCS Medical Policy Manual- Policy 310-V - Prescription Medications / Pharmacy Services, Section III 1, The Contractor and FFS Programs shall approve the Preferred Drugs listed for the therapeutic classes contained on the AHCCCS Drug List, as appropriate, before approving a Non-Preferred Drug unless ONE of the following applies:

**2.1** Submission of medical records (e.g., chart notes) or paid claims confirming history of trial resulting in a therapeutic failure, contraindication, or intolerance to at least THREE preferred alternatives (if available) (Prior trials of formulary/preferred drug list (PDL) alternatives must sufficiently demonstrate that the formulary/PDL alternatives are either ineffective or inappropriate at the time of the request)

**OR**

**2.2** Submission of medical records (e.g., chart notes) from prescribing clinician supporting the medical necessity of the Non-Preferred Drug over the Preferred Drug for the patient

Notes	Off-label use may be reviewed for medical necessity and denied as such if the off-label criteria are not met. Please refer to drug specific PA guideline for off-label criteria if available.
-------	---

## 2 . Background

Clinical Practice Guidelines		
<b>DRUGDEX Strength of Recommendation</b>		
<b>Class</b>	<b>Recommendation</b>	<b>Description</b>
Class I	Recommended	The given test or treatment has been proven useful, and should be performed or administered.
Class IIa	Recommended, In Most Cases	The given test or treatment is generally considered to be useful, and is indicated in most cases.
Class IIb	Recommended, in Some Cases	The given test or treatment may be useful, and is indicated in some, but not most, cases.
Class III	Not Recommended	The given test or treatment is not useful, and should be avoided
Class Indeterminate	Evidence Inconclusive	
<b>NCCN Categories of Evidence and Consensus [A]</b>		
<b>Category</b>	<b>Level of Consensus</b>	
1	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.	
2A	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.	

2B	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

**Lexi-Drugs: Strength of Recommendation for Inclusion in Lexi-Drugs for Oncology Off-Label Use and Level of Evidence Scale for Oncology Off-Label Use**

**Strength of Recommendation for Inclusion**

<b>Strong (for proposed off-label use)</b>	The evidence persuasively supports the off-label use (ie, Level of Evidence A).
<b>Equivocal (for proposed off-label use)</b>	The evidence to support the off-label use is of uncertain clinical significance (ie, Level of Evidence B, C). Additional studies may be necessary to further define the role of this medication for the off-label use.

<p><b>Against proposed off-label use</b></p>	<p>The evidence either advocates against the off-label use or suggests a lack of support for the off-label use (independent of Level of Evidence). Additional studies are necessary to define the role of this medication for the off-label use.</p>	
--	--	--

**Level of Evidence Scale for Oncology Off-Label Use**

<p><b>A</b></p>	<p><b>Consistent evidence from well-performed randomized, controlled trials or overwhelming evidence of some other form (eg, results of the introduction of penicillin treatment) to support off-label use. Further research is unlikely to change confidence in the estimate of benefit.</b></p>
<p><b>B</b></p>	<p><b>Evidence from randomized, controlled trials with important limitations (eg, inconsistent results, methodologic flaws, indirect, imprecise); or very strong evidence of some other research design. Further research (if performed) is likely to have an impact on confidence in the estimate of benefit and risk and may change the estimate.</b></p>

<b>C</b>	<b>Evidence from observational studies (eg, retrospective case series/reports providing significant impact on patient care); unsystematic clinical experience; or potentially flawed randomized, controlled trials (eg, when limited options exist for condition). Any estimate of effect is uncertain.</b>
<b>G</b>	<b>Use has been substantiated by inclusion in at least one evidence-based or consensus-based clinical practice guideline.</b>

### 3 . Endnotes

- A. NCCN Categories of Evidence and Consensus. Category 1: The recommendation is based on high-level evidence (i.e., high-powered randomized clinical trials or meta-analyses), and the NCCN Guideline Panel has reached uniform consensus that the recommendation is indicated. In this context, uniform means near unanimous positive support with some possible neutral positions. Category 2A: The recommendation is based on lower level evidence, but despite the absence of higher level studies, there is uniform consensus that the recommendation is appropriate. Lower level evidence is interpreted broadly, and runs the gamut from phase II to large cohort studies to case series to individual practitioner experience. Importantly, in many instances, the retrospective studies are derived from clinical experience of treating large numbers of patients at a member institution, so NCCN Guideline Panel Members have first-hand knowledge of the data. Inevitably, some recommendations must address clinical situations for which limited or no data exist. In these instances the congruence of experience-based judgments provides an informed if not confirmed direction for optimizing patient care. These recommendations carry the implicit recognition that they may be superseded as higher level evidence becomes available or as outcomes-based information becomes more prevalent. Category 2B: The recommendation is based on lower level evidence, and there is nonuniform consensus that the recommendation should be made. In these instances, because the evidence is not conclusive, institutions take different approaches to the management of a particular clinical scenario. This nonuniform consensus does not represent a major disagreement, rather it recognizes that given imperfect information, institutions may adopt different approaches. A Category 2B designation should signal to the user that more than one approach can be inferred from the existing data. Category 3: Including the recommendation has engendered a major disagreement among the NCCN Guideline Panel Members. The level of evidence is not pertinent in this category, because experts can disagree about the significance of high level trials. Several circumstances can cause major disagreements. For example, if substantial data exist about two interventions but they have never been directly compared in a randomized trial, adherents to one set of data may not accept the

interpretation of the other side's results. Another situation resulting in a Category 3 designation is when experts disagree about how trial data can be generalized. An example of this is the recommendation for internal mammary node radiation in postmastectomy radiation therapy. One side believed that because the randomized studies included this modality, it must be included in the recommendation. The other side believed, based on the documented additional morbidity and the role of internal mammary radiation therapy in other studies, that this was not necessary. A Category 3 designation alerts users to a major interpretation issue in the data and directs them to the manuscript for an explanation of the controversy.

- B. Abstracts (including meeting abstracts) are excluded from consideration. When evaluating peer-reviewed medical literature, the following (among other things) should be considered: 1) Whether the clinical characteristics of the beneficiary and the cancer are adequately represented in the published evidence 2) Whether the administered chemotherapy regimen is adequately represented in the published evidence. 3) Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. 4) Whether the study is appropriate to address the clinical question. The following should be considered: a) Whether the experimental design, in light of the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover.); b) That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs; and c) That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.

#### 4 . Revision History

Date	Notes
2/25/2025	Added criteria directing patients to preferred drugs or clinical rationale from provider. Non-cancer section: removed option allowing for FDA approved indication compendia support

Crenessity (crinecerfont)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-193196
<b>Guideline Name</b>	Crenessity (crinecerfont)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Crenessity	
Diagnosis	Congenital Adrenal Hyperplasia (CAH)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of classic 21-hydroxylase deficiency congenital adrenal hyperplasia	

**AND**

**2** - Patient is 4 years of age or older

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims documenting patient is receiving chronic treatment with glucocorticoid (GC) replacement therapy (e.g., dexamethasone, hydrocortisone, methylprednisolone) for adrenal insufficiency as one of the following:

**3.1** Both of the following:

- Patient is 4 to 17 years old
- Daily GC dose is greater than 12 mg/m<sup>2</sup>/day in hydrocortisone dose equivalents

**OR**

**3.2** Both of the following:

- Patient is 18 years of age or older
- Daily GC dose is greater than 13 mg/m<sup>2</sup>/day in hydrocortisone dose equivalents

**AND**

**4** - Prescribed by an endocrinologist

Product Name:Crenessity	
Diagnosis	Congenital Adrenal Hyperplasia (CAH)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting that patient demonstrates positive clinical response to therapy (e.g., lowered androgen levels, reduced daily dose of steroids)

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims documenting patient continues to receive chronic treatment with glucocorticoid (GC) replacement therapy (e.g., dexamethasone, hydrocortisone, methylprednisolone)

**AND**

**3** - Prescribed by or in consultation with an endocrinologist

**2 . Revision History**

Date	Notes
2/25/2025	New program

Cuvrior (trientine hydrochloride)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-127083
<b>Guideline Name</b>	Cuvrior (trientine hydrochloride)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Cuvrior	
Diagnosis	Wilson's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Wilson's disease (i.e., hepatolenticular degeneration)	

**AND**

**2** - Documentation of one of the following:

- Presence of Kayser-Fleisher rings
- Serum ceruloplasmin (CPN) less than 20 mg/dL
- 24-hour urinary copper excretion greater than 100 mcg
- Liver biopsy with copper dry weight greater than 250 mcg/g
- ATP7B mutation via genetic testing

**AND**

**3** - Trial and failure, contraindication, or intolerance to generic penicillamine capsules

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Gastroenterologist
- Hepatologist

Product Name:Cuvrior	
Diagnosis	Wilson's disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of a positive clinical response to therapy	

## 2 . Revision History

Date	Notes
6/26/2023	New Program

Cystaran, Cystadrops

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99663
<b>Guideline Name</b>	Cystaran, Cystadrops
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Cystaran, Cystadrops	
Diagnosis	Cystinosis
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystinosis	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona Medicaid SP to Medicaid Arizona SP for eff 7 /1

Daliresp (roflumilast)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117633
<b>Guideline Name</b>	Daliresp (roflumilast)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Brand Daliresp, generic roflumilast	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of severe to very severe chronic obstructive pulmonary disease (COPD) (i.e., FEV1 less than or equal to 50% of predicted)	

**AND**

**2** - COPD is associated with chronic bronchitis

**AND**

**3** - History of COPD exacerbation(s)

Product Name:Brand Daliresp, generic roflumilast	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Daliresp therapy	

## 2 . Revision History

Date	Notes
12/4/2022	Added generic roflumilast as target

Daraprim

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99605
<b>Guideline Name</b>	Daraprim
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Daraprim, generic pyrimethamine	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Medical record documentation (e.g. chart notes) of one of the following:  1.1 Treatment of severe acquired toxoplasmosis, including toxoplasmic encephalitis  <b>OR</b>	

**1.2 Treatment of congenital toxoplasmosis**

**OR**

**1.3 Secondary prophylaxis of toxoplasmic encephalitis**

**OR**

**1.4 ALL of the following:**

**1.4.1** Primary Pneumocystis pneumonia (PCP) prophylaxis in human immunodeficiency virus (HIV)-infected patients or as secondary prophylaxis in HIV-infected patients who have been treated for an acute episode of Pneumocystis pneumonia

**AND**

**1.4.2** Patient has experienced intolerance to prior prophylaxis with trimethoprim-sulfamethoxazole (TMP-SMX)

**AND**

**1.4.3 ONE of the following:**

**1.4.3.1** Patient has been re-challenged with trimethoprim-sulfamethoxazole (TMP-SMX) using a desensitization protocol and is still unable to tolerate

**OR**

**1.4.3.2** Evidence of moderately severe or life threatening-reaction to trimethoprim-sulfamethoxazole (TMP-SMX) in the past (e.g. toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome)

**OR**

**1.5 ALL of the following:**

**1.5.1** Primary prophylaxis of toxoplasmic encephalitis

**AND**

**1.5.2** Toxoplasma immunoglobulin G (IgG) positive

**AND**

**1.5.3** CD4 (cluster of differentiation 4) less than or equal to 100 cells per mm<sup>3</sup> if initiating prophylaxis or CD4 100-200 cells per mm<sup>3</sup> if reinstating prophylaxis

**AND**

**1.5.4** Will be used in combination with dapsone or atovaquone

**AND**

**1.5.5** Patient has experienced intolerance to prior prophylaxis with trimethoprim-sulfamethoxazole (TMP-SMX)

**AND**

**1.5.6** ONE of the following:

**1.5.6.1** Patient has been re-challenged with trimethoprim-sulfamethoxazole (TMP-SMX) using a desensitization protocol and is still unable to tolerate

**OR**

**1.5.6.2** Evidence of moderately severe or life threatening-reaction to trimethoprim-sulfamethoxazole (TMP-SMX) in the past (e.g. toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome)

Notes

\*Consider discontinuation of primary prophylaxis if CD4 greater than 200 cells/mm<sup>3</sup> for greater than 3 months after institution of combination antiretroviral therapy.

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona SP to Medicaid Arizona SP for 7/1 eff

Daxxify (botulinum toxin type a injection)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135325
<b>Guideline Name</b>	Daxxify (botulinum toxin type a injection)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Daxxify	
Diagnosis	Cervical Dystonia
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of cervical dystonia	

Product Name:Daxxify	
Diagnosis	Cervical Dystonia
Approval Length	3 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - At least 3 months have or will have elapsed since the last treatment</p>	

Product Name:Daxxify	
Diagnosis	Cosmetic Use
Approval Length	N/A - requests for cosmetic use are excluded and will not be approved
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Requests for coverage of any Daxxify product for treating the appearance of facial lines are not authorized and will not be approved. These uses are considered cosmetic only and are excluded from coverage.</p>	
Notes	Requests for coverage of any Daxxify product for treating the appearance of facial lines are not authorized and will not be approved. These uses are considered cosmetic only and are excluded from coverage.

## 2 . Revision History

Date	Notes
------	-------

10/23/2023	New program
------------	-------------

Daybue (trofinetide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-125943
<b>Guideline Name</b>	Daybue (trofinetide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Daybue	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Rett syndrome  <b>AND</b>	

**2 - One of the following:**

**2.1** Submission of medical records (e.g., chart notes) confirming presence of ALL of the following clinical signs and symptoms:

- A pattern of development, regression, then recovery or stabilization
- Partial or complete loss of purposeful hand skills such as grasping with fingers, reaching for things, or touching things on purpose
- Partial or complete loss of spoken language
- Repetitive hand movements, such as wringing the hands, washing, squeezing, clapping, or rubbing
- Gait abnormalities, including walking on toes or with an unsteady, wide-based, stiff-legged gait

**OR**

**2.2** Submission of medical records (e.g., chart notes) documenting molecular genetic testing confirms mutations in the MECP2 gene

**AND**

**3 - Patient is 2 years of age or older**

**AND**

**4 - Prescribed by or in consultation with one of the following:**

- Geneticist
- Neurologist

Product Name:Daybue	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy

**2 . Revision History**

Date	Notes
5/22/2023	New program

DDAVP (desmopressin) tablets - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-105310
<b>Guideline Name</b>	DDAVP (desmopressin) tablets - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Brand DDAVP tablets, generic desmopressin acetate tablets	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Diagnosis of central diabetes insipidus</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** Diagnosis of polyuria and/or polydipsia following head trauma or surgery in the pituitary region

**OR**

**1.3** Diagnosis of primary nocturnal enuresis

**AND**

**2** - For Brand DDAVP ONLY: Trial and failure to generic desmopressin tablets (verified via paid pharmacy claims or submission of medical records)

Notes	NOTE: Plan setup requires use of generic desmopressin tablets before Brand DDAVP
-------	--

## 2 . Revision History

Date	Notes
3/29/2022	Added step through generic tablets for Brand.

Declomycin - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99559
<b>Guideline Name</b>	Declomycin - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:demeclocycline*	
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 Diagnosis of ONE of the following: <ul style="list-style-type: none"><li>Rocky Mountain spotted fever, typhus fever and the typhus group, Q fever, rickettsialpox and tick fevers caused by rickettsiae</li><li>Respiratory tract infections caused by Mycoplasma pneumoniae</li><li>Lymphogranuloma venereum due to Chlamydia trachomatis</li></ul>	

- Psittacosis (Ornithosis) due to *Chlamydia psittaci*
- Trachoma due to *Chlamydia trachomatis*
- Inclusion conjunctivitis caused by *Chlamydia trachomatis*
- Nongonococcal urethritis in adults caused by *Ureaplasma urealyticum* or *Chlamydia trachomatis*
- Relapsing fever due to *Borrelia recurrentis*
- Chancroid caused by *Haemophilus ducreyi*
- Plague due to *Yersinia pestis*
- Tularemia due to *Francisella tularensis*
- Cholera caused by *Vibrio cholerae*
- *Campylobacter fetus* infections caused by *Campylobacter fetus*
- Brucellosis due to *Brucella* species (in conjunction with streptomycin)
- Bartonellosis due to *Bartonella bacilliformis*
- Granuloma inguinale caused by *Calymmatobacterium granulomatis*
- Infection due to *Escherichia coli*
- Infection due to *Enterobacter aerogenes*
- Infection due to *Shigella* species
- Infection due to *Acinetobacter* species
- Respiratory tract infections caused by *Haemophilus influenza*
- Respiratory tract and urinary tract infections caused by *Klebsiella* species
- Upper respiratory infections caused by *Streptococcus pneumoniae*
- Skin and skin structure infections caused by *Staphylococcus aureus*.
- Uncomplicated urethritis in men due to *Neisseria gonorrhoeae*, and for the treatment of other uncomplicated gonococcal infections
- Infections in women caused by *Neisseria gonorrhoeae*
- Syphilis caused by *Treponema pallidum* subspecies *pallidum*
- Yaws caused by *Treponema pallidum* subspecies *pertenue*
- Listeriosis due to *Listeria monocytogenes*
- Anthrax due to *Bacillus anthracis*
- Vincent's infection caused by *Fusobacterium fusiforme*
- Actinomycosis caused by *Actinomyces israelii*
- Clostridial diseases caused by *Clostridium* species
- Acute intestinal amebiasis, as adjunctive therapy
- Severe acne, as adjunctive therapy

**OR**

**1.2** The medication is being prescribed by or in consultation with an Infectious Disease specialist

Notes	*Approval duration: 6 months
-------	------------------------------

## 2 . Revision History

Date	Notes
------	-------

6/23/2021	update program
-----------	----------------

Dificid

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99444
<b>Guideline Name</b>	Dificid
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Dificid	
Approval Length	10 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of Clostridioides difficile-associated diarrhea (CDAD) [previously known as Clostridium difficile- associated diarrhea]  <b>AND</b>	

**2 - ONE of the following:**

**2.1** History of failure, contraindication, or intolerance to Firvanq (vancomycin) oral solution

**OR**

**2.2** History of failure, contraindication, or intolerance to oral Vancocin (vancomycin) capsules or vancomycin oral solution (NOT Firvanq) if the prescriber provides a reason or special circumstance the patient cannot use Firvanq

**OR**

**2.3** For continuation of prior Difcid therapy

## **2 . Revision History**

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Dofetilide - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99445
<b>Guideline Name</b>	Dofetilide - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:: Brand Tikosyn, generic dofetilide	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of ONE of the following: <ul style="list-style-type: none"><li>Atrial fibrillation</li><li>Atrial flutter</li></ul>	

**AND**

**2** - Patient requires ONE of the following:

- Conversion to normal sinus rhythm
- Maintenance of normal sinus rhythm

**AND**

**3** - Verification that the patient has already started on dofetilide while in the hospital for a minimum of 3 days

**AND**

**4** - Patient does NOT have severe renal impairment [Creatinine Clearance (CrCl) less than 20 milliliters per minute]

**AND**

**5** - Patient does NOT have congenital or acquired long QT syndromes

**AND**

**6** - Patient is NOT concurrently using cimetidine, hydrochlorothiazide, ketoconazole, megestrol, prochlorperazine, trimethoprim, dolutegravir or verapamil

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Dojolvi (triheptanoin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116190
<b>Guideline Name</b>	Dojolvi (triheptanoin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Dojolvi	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of a long-chain fatty acid oxidation disorder (LC-FAOD) has been confirmed by at least two of the following: <ul style="list-style-type: none"><li>Disease specific elevation of acyl-carnitines on a newborn blood spot or in plasma</li><li>Low enzyme activity in cultured fibroblasts</li></ul>	

- One or more known pathogenic mutations in CPT2, ACADVL, HADHA, or HADHB

**AND**

**2** - Not used with any other medium-chain triglyceride (MCT) product

**AND**

**3** - Prescribed by or in consultation with a clinical specialist knowledgeable in appropriate disease-related dietary management (e.g., geneticist, cardiologist, gastroenterologist, etc.)

Product Name: Dojolvi	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Prescriber attests to continued need of therapy</p> <p><b>AND</b></p> <p><b>2</b> - Not used with any other medium-chain triglyceride (MCT) product</p> <p><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with a clinical specialist knowledgeable in appropriate disease-related dietary management (e.g., geneticist, cardiologist, gastroenterologist, etc.)</p>	

## 2 . Revision History

Date	Notes
------	-------

10/28/2022	New Program
------------	-------------

DPP-4 Inhibitors - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157632
<b>Guideline Name</b>	DPP-4 Inhibitors - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name: PREFERRED: generic alogliptin, generic alogliptin-metformin, generic alogliptin-pioglitazone, Janumet, Janumet XR, Januvia, Jentadueto, Jentadueto XR, Tradjenta	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The patient has a diagnosis of type 2 diabetes mellitus  <b>AND</b>	

**2 - ONE of the following:**

**2.1** History of failure to metformin at a minimum dose of 1500 milligrams daily for 90 days

**OR**

**2.2** Contraindication or intolerance to metformin

Product Name:NON-PREFERRED: Kombiglyze XR, generic saxagliptin-metformin ER, Brand Onglyza, generic saxagliptin, Brand Zituvimet, Brand Sitagliptin-metformin, Zituvimet XR, Brand Zituvio, Brand Sitagliptin

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1 -** The patient has a diagnosis of type 2 diabetes mellitus

**AND**

**2 - ONE of the following:**

**2.1** History of failure to metformin at a minimum dose of 1500 milligrams daily for 90 days

**OR**

**2.2** Contraindication or intolerance to metformin

**AND**

**3 - ONE of the following:**

**3.1** History of failure for 90 days to three of the following:

- generic alogliptin, generic alogliptin-metformin, or generic alogliptin-pioglitazone

- Januvia
- Janumet/Janumet XR
- Jentadueto/Jentadueto XR
- Tradjenta

**OR**

**3.2 Intolerance or contraindication to THREE of the following:**

- generic alogliptin, generic alogliptin-metformin, or generic alogliptin-pioglitazone
- Januvia
- Janumet/Janumet XR
- Jentadueto/Jentadueto XR
- Tradjenta

## 2 . Revision History

Date	Notes
10/25/2024	Added Zituvimet and Zituvimet XR as NP targets

Drugs for Compassionate Use

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-154534
<b>Guideline Name</b>	Drugs for Compassionate Use
<b>Formulary</b>	<ul style="list-style-type: none"> <li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li> <li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li> </ul>

**Guideline Note:**

Effective Date:	10/1/2024
-----------------	-----------

### 1 . Criteria

Product Name:Eulexin, Humatin, Leukeran, Myleran, Tabloid	
Approval Length	N/A- drugs not covered by Medicaid must be obtained through Waylis Therapeutics for Compassionate Use
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Provider attests that the requested medication will be obtained under compassionate use through Wayliss Pharmaceuticals</p>	
Notes	Approval Length: N/A- Requests for drugs not covered by Medicaid must be obtained through Waylis Therapeutics for Compassionate Use

## 2 . Revision History

Date	Notes
9/26/2024	New program

Dry Eye Disease - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144641
<b>Guideline Name</b>	Dry Eye Disease - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Preferred: Xiidra	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Tear deficiency associated with ocular inflammation due to one of the following: <ul style="list-style-type: none"><li>Moderate to severe keratoconjunctivitis sicca</li><li>Moderate to severe dry eye disease</li></ul>	

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming diagnosis by **ONE** of the following diagnostic tests:

- Schirmer test
- Ocular surface dye staining (e.g., rose bengal, fluorescein, lissamine green)
- Tear function index/fluorescein clearance test
- Tear break up time
- Tear film osmolarity
- Slit lamp lid evaluation
- Lacrimal gland function

**AND**

**3** - Medication is not being prescribed to manage dry eyes peri-operative elective eye surgery (e.g., LASIK)

**AND**

**4** - Submission of medical records (e.g., chart notes) or paid claims confirming trial and failure, contraindication, or intolerance to at least one OTC ocular lubricant (e.g., artificial tears, lubricating gels/ointments) in the past 60 days

**AND**

**5** - Prescribed by or in consultation with **ONE** of the following:

- Ophthalmologist
- Optometrist
- Rheumatologist

**AND**

**6** - Submission of medical records (e.g., chart notes) or paid claims confirming a minimum trial of 60 days of Brand Restasis single dose vials, unless contraindicated

Product Name: Non-Preferred: Cequa, generic cyclosporine emulsion, Miebo, Restasis MultiDose, Tyrvaya, Vevye

Approval Length | 12 month(s)

Therapy Stage | Initial Authorization

Guideline Type | Prior Authorization

### Approval Criteria

1 - Tear deficiency associated with ocular inflammation due to one of the following:

- Moderate to severe keratoconjunctivitis sicca
- Moderate to severe dry eye disease

**AND**

2 - Submission of medical records (e.g., chart notes) confirming diagnosis by ONE of the following diagnostic tests:

- Schirmer test
- Ocular surface dye staining (e.g., rose bengal, fluorescein, lissamine green)
- Tear function index/fluorescein clearance test
- Tear break up time
- Tear film osmolarity
- Slit lamp lid evaluation
- Lacrimal gland function

**AND**

3 - Medication is not being prescribed to manage dry eyes peri-operative elective eye surgery (e.g., LASIK)

**AND**

4 - Submission of medical records (e.g., chart notes) or paid claims confirming trial and failure, contraindication, or intolerance to at least one OTC ocular lubricant (e.g., artificial tears, lubricating gels/ointments) in the past 60 days

**AND**

**5** - Prescribed by or in consultation with ONE of the following:

- Ophthalmologist
- Optometrist
- Rheumatologist

**AND**

**6** - Submission of medical records (e.g., chart notes) or paid claims confirming a minimum trial of 60 days of BOTH of the following, unless contraindicated:

- Brand Restasis single dose vials
- Xiidra (PA may be required)

Product Name: Preferred: Xiidra; Non-Preferred: Cequa, generic cyclosporine emulsion, Miebo, Restasis MultiDose, Tyrvaya, Vevye	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy (e.g., increased tear production or improvement in dry eye symptoms)	

## 2 . Revision History

Date	Notes
3/27/2024	Updated criteria/preferred status from Jan P&T, Xiidra now preferred. Added Miebo, Tyrvaya and Restasis Multidose as NP targets.

Duexis and Vimovo - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99563
<b>Guideline Name</b>	Duexis and Vimovo - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Duexis	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following risk factors for NSAID (non-steroidal anti-inflammatory drug) induced adverse GI (gastrointestinal) events: <ul style="list-style-type: none"><li>Patient is greater than or equal to 65 years of age</li><li>Prior history of peptic, gastric, or duodenal ulcer</li><li>History of NSAID-related ulcer</li><li>History of clinically significant GI bleeding</li></ul>	

- Untreated or active H. Pylori gastritis
- Concurrent use of oral corticosteroids (eg, prednisone, prednisolone, dexamethasone)
- Concurrent use of anticoagulants (eg, warfarin, heparin)
- Concurrent use of antiplatelets (eg, aspirin including low-dose, clopidogrel)

**AND**

**2** - Documentation of history of failure, contraindication, or intolerance to THREE combinations of preferred NSAIDS taken with preferred H2 (histamine 2)-receptor antagonists. (Provide name and date preferred products were tried)\*

**AND**

**3** - Physician has provided rationale for needing to use fixed-dose combination therapy with Duexis instead of taking individual products in combination.

Notes	*Please reference background section for preferred products table
-------	---

Product Name: Brand Vimovo, generic naproxen-esomeprazole

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - ONE of the following risk factors for NSAID (non-steroidal anti-inflammatory drug) induced adverse GI (gastrointestinal) events:

- Patient is greater than or equal to 65 years of age
- Prior history of peptic, gastric, or duodenal ulcer
- History of NSAID-related ulcer
- History of clinically significant GI bleeding
- Untreated or active H. Pylori gastritis
- Concurrent use of oral corticosteroids (eg, prednisone, prednisolone, dexamethasone)
- Concurrent use of anticoagulants (eg, warfarin, heparin)
- Concurrent use of antiplatelets (eg, aspirin including low-dose, clopidogrel)

**AND**

**2** - Documentation of history of failure, contraindication, or intolerance to THREE combinations of preferred NSAIDS taken with preferred proton pump inhibitors (PPIs). (Provide name and date preferred products were tried)\*

**AND**

**3** - Physician has provided rationale for needing to use fixed-dose combination therapy with Vimovo instead of taking individual products in combination.

Notes

\*Please reference background section for preferred products table

## 2 . Background

Benefit/Coverage/Program Information		
Preferred Table		
NSAIDS	Proton Pump Inhibitors (PPIs)	H2 (histamine 2)-receptor antagonists
Diclofenac DR (Generic Voltaren)	esomeprazole (Generic Nexium)	Famotidine (Generic Pepcid)
Diclofenac ER (Generic Voltaren ER)	lansoprazole (Generic Prevacid)	Nizatidine (Generic Axid)
Etodolac (Generic Lodine)	omeprazole (Generic Prilosec)	Ranitidine (Generic Zantac)
Etodolac ER (Generic Lodine ER)	pantoprazole sodium (Generic Protonix)	
Fenoprofen (Generic Nalfon)		
Flurbiprofen (Generic Ansaid)		

Ibuprofen		
Indomethacin (Generic Indocin)		
Ketorolac (Generic Toradol)		
Mefenamic (Generic Ponstel)		
Meloxicam (Generic Mobic)		
Nabumetone (Generic Relafen)		
Nabumetone DS (Generic Relafen DS)		
Naproxen (Generic Anaprox)		
Naproxen DR (Generic Anaprox DR)		
Naproxen EC (Generic Anaprox EC)		
Oxaprozin (Generic Daypro)		

Piroxicam (Generic Feldene)		
Sulindac (Generic Clinoril)		

Dupixent (dupilumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164442
<b>Guideline Name</b>	Dupixent (dupilumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name: Dupixent	
Diagnosis	Atopic Dermatitis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of documentation (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderate to severe chronic atopic dermatitis	

**AND**

**1.2** One of the following:

**1.2.1** Both of the following:

**1.2.1.1** Patient is 12 years of age and older

**AND**

**1.2.1.2** History of failure, contraindication, or intolerance to ALL of the following: (document drug, date of trial, and/or contraindication to medication)\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole)
- Adbry (tralokinumab-ldrm)
- Opzelura (ruxolitinib)

**OR**

**1.2.2** Both of the following:

**1.2.2.1** Patient is less than 12 years of age

**AND**

**1.2.2.2** History of failure, contraindication, or intolerance to ALL of the following: (document drug, date of trial, and/or contraindication to medication)\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole)
- One topical corticosteroid [e.g., mometasone furoate, fluocinolone acetonide (generic Synalar), fluocinonide]

**AND**

**2** - Patient is NOT receiving Dupixent in combination with another biologic medication [e.g., Xolair (omalizumab), Rituxan (rituximab), Enbrel (etanercept), Remicade/Inflectra (infliximab)]

**AND**

**3** - Prescribed by one of the following:

- Dermatologist
- Allergist
- Immunologist

Notes

\*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication

Product Name: Dupixent

Diagnosis Atopic Dermatitis

Approval Length 6 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Submission of documentation (e.g., chart notes) or verification of paid claims confirming ALL of the following:

**1.1** Positive clinical response to therapy

**AND**

**1.2** One of the following:

**1.2.1** Both of the following:

**1.2.1.1** Patient is 12 years of age and older

**AND**

**1.2.1.2** History of failure, contraindication, or intolerance to ALL of the following: (document drug, date of trial, and/or contraindication to medication)\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole)
- Adbry (tralokinumab-ldrm)
- Opzelura (ruxolitinib)

**OR**

**1.2.2** Both of the following:

**1.2.2.1** Patient is less than 12 years of age

**AND**

**1.2.2.2** History of failure, contraindication, or intolerance to ALL of the following: (document drug, date of trial, and/or contraindication to medication)\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole)
- One topical corticosteroid [e.g., mometasone furoate, fluocinolone acetonide (generic Synalar), fluocinonide]

**AND**

**2** - Patient is NOT receiving Dupixent in combination with another biologic medication [e.g., Xolair (omalizumab), Rituxan (rituximab), Enbrel (etanercept), Remicade/Inflectra (infliximab)]

**AND**

**3** - Prescribed by one of the following:

- Dermatologist
- Allergist
- Immunologist

Notes

\*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication

Product Name: Dupixent	
Diagnosis	Asthma
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is 6 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of documentation (e.g., chart notes) confirming diagnosis of moderate to severe asthma</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - ONE of the following:</p> <p>3.1 ALL of the following:</p> <p>3.1.1 Classification of asthma as uncontrolled or inadequately controlled as defined by at least ONE of the following</p> <ul style="list-style-type: none"> <li>• Poor symptom control (e.g., Asthma Control Questionnaire [ACQ] score consistently greater than 1.5 or Asthma Control Test [ACT] score consistently less than 20)</li> <li>• Two or more bursts of systemic corticosteroids for at least 3 days each in the previous 12 months</li> <li>• Asthma-related emergency treatment (e.g., emergency room visit, hospital admission, or unscheduled physician's office visit for nebulizer or other urgent treatment)</li> <li>• Airflow limitation (e.g., after appropriate bronchodilator withhold forced expiratory volume in 1 second [FEV1] less than 80% predicted [in the face of reduced FEV1/forced vital capacity [FVC] defined as less than the lower limit of normal])</li> <li>• Patient is currently dependent on oral corticosteroids for the treatment of asthma</li> </ul> <p style="text-align: center;"><b>AND</b></p>	

**3.1.2** Dupixent will be used in combination with one of the following:

**3.1.2.1** ONE high-dose (appropriately adjusted for age) combination inhaled corticosteroid (ICS)/long-acting beta2 agonist (LABA) [e.g., Advair/AirDuo Respiclick (fluticasone propionate/salmeterol), Symbicort (budesonide/formoterol), Breo Ellipta (fluticasone furoate/vilanterol)] (see Table 2 in Background section)

**OR**

**3.1.2.2** Combination therapy including BOTH of the following:

**3.1.2.2.1** ONE high-dose (appropriately adjusted for age) ICS product [e.g., ciclesonide (Alvesco), mometasone furoate (Asmanex), beclomethasone dipropionate (QVAR)] (see Table 2 in Background section)

**AND**

**3.1.2.2.2** ONE additional asthma controller medication [e.g., LABA - olodaterol (Striverdi) or indacaterol (Arcapta); leukotriene receptor antagonist – montelukast (Singulair); theophylline]

**AND**

**3.1.3** ONE of the following:

**3.1.3.1** Submission of medical records (e.g., chart notes, laboratory values, etc.) documenting that asthma is an eosinophilic phenotype as defined by a baseline (pre-dupilumab treatment) peripheral blood eosinophil level greater than or equal to 150 cells/microliter within the past 6 weeks

**OR**

**3.1.3.2** Patient is currently dependent on oral corticosteroids for the treatment of asthma

**OR**

**3.2** Patient is currently on Dupixent therapy

**AND**

**4** - Patient is NOT receiving Dupixent in combination with ONE of the following:

- Anti-interleukin-5 therapy [e.g. Nucala (mepolizumab), Cinqair (reslizumab), Fasenra (benralizumab)]
- Anti-IgE (immunoglobulin E) therapy [e.g. Xolair (omalizumab)]

**AND**

**5** - Prescribed by one of the following:

- Pulmonologist
- Allergist
- Immunologist

Product Name: Dupixent	
Diagnosis	Asthma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of documentation (e.g., chart notes) confirming positive clinical response to Dupixent therapy as demonstrated by at least ONE of the following:	
<ul style="list-style-type: none"><li>• Reduction in the frequency of exacerbations</li><li>• Decreased utilization of rescue medications</li><li>• Increase in percent predicted forced expiratory volume in 1 second (FEV1) from pretreatment baseline</li><li>• Reduction in severity or frequency of asthma-related symptoms (e.g., wheezing, shortness of breath, coughing, etc.)</li><li>• Reduction in oral corticosteroid requirements</li></ul>	

**AND**

**2** - Dupixent is being used in combination with an inhaled corticosteroid (ICS)-containing controller medication (see Table 2 in Background section)

**AND**

**3** - Patient is NOT receiving Dupixent in combination with ONE of the following:

- Anti-interleukin-5 therapy [e.g. Nucala (mepolizumab), Cinqair (reslizumab), Fasenna (benralizumab)]
- Anti-IgE (immunoglobulin E) therapy [e.g. Xolair (omalizumab)]

**AND**

**4** - Prescribed by one of the following:

- Pulmonologist
- Allergist
- Immunologist

Product Name: Dupixent	
Diagnosis	Chronic Rhinosinusitis with Nasal Polyposis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient is 12 years of age or older	
<b>AND</b>	

**2** - Submission of documentation (e.g., chart notes) confirming ONE of the following:

**2.1** ALL of the following:

**2.1.1** Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) defined by ALL of the following:

**2.1.1.1** TWO or more of the following symptoms for greater than or equal to 12 weeks duration:

- Mucopurulent discharge
- Nasal obstruction and congestion
- Decreased or absent sense of smell
- Facial pressure or pain

**AND**

**2.1.1.2** ONE of the following:

- Evidence of inflammation on paranasal sinus examination or computed tomography (CT)
- Evidence of purulence coming from paranasal sinuses or ostiomeatal complex

**AND**

**2.1.1.3** The presence of nasal polyps

**AND**

**2.1.2** ONE of the following:

- Patient has required prior sino-nasal surgery
- Patient has required systemic corticosteroids in the previous 2 years

**AND**

**2.1.3** Patient has been unable to obtain symptom relief after trial of ALL of the following agents/classes of agents:

- Nasal saline irrigations

- Intranasal corticosteroids (e.g. fluticasone, mometasone, triamcinolone, etc.)
- Antileukotriene agents (e.g. montelukast, zafirlukast, zileuton)

**OR**

**2.2** ALL of the following:

**2.2.1** Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP)

**AND**

**2.2.2** Patient is currently on Dupixent therapy

**AND**

**3** - Patient will receive Dupixent as add-on maintenance therapy in combination with intranasal corticosteroids

**AND**

**4** - Patient is NOT receiving Dupixent in combination with another biologic medication [e.g., Xolair (omalizumab), Nucala (mepolizumab), Cinqair (reslizumab), Fasentra (benralizumab)]

**AND**

**5** - Prescribed by one of the following:

- Otolaryngologist
- Allergist
- Immunologist

Product Name: Dupixent	
Diagnosis	Chronic Rhinosinusitis with Nasal Polyposis
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of documentation (e.g., chart notes) confirming positive clinical response to Dupixent therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient will continue to receive Dupixent as add-on maintenance therapy in combination with intranasal corticosteroids</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is NOT receiving Dupixent in combination with another biologic medication [e.g., Xolair (omalizumab), Nucala (mepolizumab), Cinqair (reslizumab), Fasenra (benralizumab)]</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by one of the following:</p> <ul style="list-style-type: none"> <li>• Otolaryngologist</li> <li>• Allergist</li> <li>• Immunologist</li> </ul>	

Product Name: Dupixent	
Diagnosis	Eosinophilic Esophagitis (EoE)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

**1** - Both of the following:

- Patient is at least 1 year of age
- Patient weighs at least 15 kg

**AND**

**2** - Submission of documentation (e.g., chart notes) confirming diagnosis of eosinophilic esophagitis (EoE)

**AND**

**3** - Patient has symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, gastroesophageal reflux disease [GERD]/heartburn symptoms, chest pain, abdominal pain)

**AND**

**4** - Submission of documentation (e.g., chart notes, lab values) confirming patient has at least 15 intraepithelial eosinophils per high power field (HPF)

**AND**

**5** - Other causes of esophageal eosinophilia have been excluded

**AND**

**6** - Paid claims or submission of documentation (e.g., chart notes) confirming trial and failure, contraindication, or intolerance to at least an 8-week trial of one of the following:

- Proton pump inhibitors (e.g., pantoprazole, omeprazole)
- Topical (esophageal) corticosteroids (e.g., budesonide, fluticasone)

**AND**

**7** - Prescribed by one of the following:

- Gastroenterologist
- Allergist
- Immunologist

Product Name: Dupixent	
Diagnosis	Eosinophilic Esophagitis (EoE)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of documentation (e.g., chart notes) confirming positive clinical response to therapy as evidenced by improvement of at least one of the following from baseline:</p> <ul style="list-style-type: none"> <li>• Symptoms (e.g., dysphagia, food impaction, heartburn, chest pain)</li> <li>• Histologic measures (e.g., esophageal intraepithelial eosinophil count)</li> <li>• Endoscopic measures (e.g., edema, furrows, exudates, rings, strictures)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by one of the following:</p> <ul style="list-style-type: none"> <li>• Gastroenterologist</li> <li>• Allergist</li> <li>• Immunologist</li> </ul>	

Product Name: Dupixent	
Diagnosis	Prurigo Nodularis (PN)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient is 18 years of age or older

**AND**

2 - Submission of documentation (e.g., chart notes) confirming diagnosis of prurigo nodularis (PN)

**AND**

3 - Patient has at least 20 nodular lesions

**AND**

4 - Trial and failure, contraindication, or intolerance to one previous PN treatment (e.g., topical corticosteroids, topical calcineurin inhibitors [pimecrolimus, tacrolimus], topical capsaicin)

**AND**

5 - Prescribed by one of the following:

- Dermatologist
- Allergist
- Immunologist

Product Name: Dupixent	
Diagnosis	Prurigo Nodularis (PN)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of documentation (e.g., chart notes) confirming positive clinical response to therapy as evidenced by improvement of at least one of the following:

- Reduction in the number of nodular lesions from baseline
- Improvement in symptoms (e.g., pruritus, inflammation) from baseline

**AND**

2 - Prescribed by one of the following:

- Dermatologist
- Allergist
- Immunologist

Product Name: Dupixent	
Diagnosis	Chronic obstructive pulmonary disease (COPD)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is 18 years of age or older</p> <p><b>AND</b></p> <p>2 - Submission of medical records (e.g., chart notes) confirming diagnosis of chronic obstructive pulmonary disease (COPD)</p> <p><b>AND</b></p>	

**3** - Submission of medical records (e.g., chart notes) confirming the presence of Type 2 inflammation evidenced by blood eosinophils greater than or equal to 300 cells per microliter at baseline

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) confirming the patient is receiving one of the following therapies at maximally tolerated doses:

- Triple therapy (i.e., an inhaled corticosteroid (ICS), a long-acting muscarinic antagonist (LAMA) and a long-acting beta agonist (LABA)
- If ICS are contraindicated, a LAMA and a LABA

**AND**

**5** - Patient must have post-bronchodilator forced expiratory volume [FEV1] / forced vital capacity [FVC] ratio less than 0.70

**AND**

**6** - Patient has had BOTH of the following within the past 12 months:

- At least two exacerbations where systemic corticosteroids [intramuscular, intravenous, or oral (e.g., prednisone)] were required at least once
- A COPD-related hospitalization

**AND**

**7** - Submission of medical records (e.g., chart notes) confirming the patient experiences dyspnea during everyday activities (e.g., needs to stop for breath when walking on level ground)

Product Name: Dupixent	
Diagnosis	Chronic obstructive pulmonary disease (COPD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a positive clinical response to therapy (e.g., improved lung function, a reduction in COPD exacerbations)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Paid claims or submission of medical records (e.g., chart notes) confirming the patient continues to receive one of the following therapies:</p> <ul style="list-style-type: none"> <li>• Triple therapy (i.e., an inhaled corticosteroid (ICS), a long-acting muscarinic antagonist (LAMA) and a long-acting beta agonist (LABA)</li> <li>• If ICS are contraindicated, a LAMA and a LABA</li> </ul>	

## 2 . Background

Benefit/Coverage/Program Information			
Table 1: Relative potencies of topical corticosteroids			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment, gel	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream, lotion	0.05

	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05
	tridifloronide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment, lotion	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream, lotion	0.1
	Triamcinolone acetonide	Cream, ointment, lotion	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetamide	Cream, solution	0.01
	Dexamethasone	Cream	0.1

Lowest potency	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

**Table 2: Low, medium and high daily doses of inhaled corticosteroids Adults and adolescents (12 years of age and older)**

Drug	Daily dose (mcg)		
	Low	Medium	High
Beclometasone dipropionate (CFC)	200-500	>500-1000	>1000
Beclometasone dipropionate (HFA)	100-200	>200-400	>400
Budesonide DPI	200-400	>400-800	>800
Ciclesonide (HFA)	80-160	>160-320	>320
Fluticasone furoate (DPI)	100	N/A	200
Fluticasone propionate (DPI)	100-250	>250-500	>500
Fluticasone propionate (HFA)	100-250	>250-500	>500
Mometasone furoate	110-220	>220-440	>440
Triamcinolone acetonide	400-1000	>1000-2000	>2000

### 3 . Revision History

Date	Notes
1/30/2025	Added age criterion to initial auth for PN and COPD indications

Durezol

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99567
<b>Guideline Name</b>	Durezol
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Durezol	
Approval Length	2 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, contraindication, or intolerance to BOTH of the following: <ul style="list-style-type: none"><li>prednisolone 1%</li><li>dexamethasone ophthalmic drops and/or ointment.</li></ul>	

## 2 . Revision History

Date	Notes
7/8/2021	Changed approval length to 2 months

Duvyzat (givinostat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157794
<b>Guideline Name</b>	Duvyzat (givinostat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Duvyzat	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of Duchenne muscular dystrophy (DMD)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting one of the following:

**2.1** Patient has a confirmed mutation of the dystrophin gene

**OR**

**2.2** Muscle biopsy confirmed an absence of dystrophin protein

**AND**

**3** - Patient is 6 years of age or older

**AND**

**4** - Submission of medical records (e.g., chart notes) confirming patient is ambulatory without needing an assistive device (e.g., without side-by-side assist, cane, walker, wheelchair, etc.) prior to initiating Duvyzat

**AND**

**5** - Submission of medical records (e.g., chart notes) or paid claims confirming requested drug will be used concomitantly with a corticosteroid regimen (e.g., prednisone/prednisolone, Emflaza [deflazacort], Agamree)

**AND**

**6** - Prescribed by or in consultation with a pediatric neurologist with expertise in treating DMD

Product Name:Duvyzat	
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming patient has experienced a benefit from therapy (e.g., improvement in preservation of muscle strength)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Submission of medical records (e.g., chart notes) confirming patient is maintaining ambulatory status without needing an assistive device (e.g., without side-by-side assist, cane, walker, wheelchair, etc.)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Submission of medical records (e.g., chart notes) or paid claims confirming patient continues to receive concomitant corticosteroid regimen (e.g., prednisone/prednisolone, Emflaza [deflazacort], Agamree)</p>	

## 2 . Revision History

Date	Notes
10/25/2024	New program

Ebglyss (lebrikizumab-lbkz)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-161671
<b>Guideline Name</b>	Ebglyss (lebrikizumab-lbkz)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Ebglyss	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderate to severe atopic dermatitis	

**AND**

**1.2** One of the following:

- Involvement of at least 10% body surface area (BSA)
- SCORing Atopic Dermatitis (SCORAD) index value of at least 25

**AND**

**1.3** Both of the following:

- Patient is 12 years of age or older
- Patient weighs at least 40 kg

**AND**

**1.4** History of failure, contraindication, or intolerance to ALL of the following: (document drug, date of trial, and/or contraindication to medication)\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole) ointment
- Adbry (tralokizumab-ldrm)
- Dupixent (dupilumab)

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist
- Immunologist

Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication
-------	--

Product Name: Ebglyss	
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of documentation (e.g., chart notes) demonstrating positive clinical response to therapy as evidenced by at least ONE of the following:</p> <ul style="list-style-type: none"> <li>• Reduction in body surface area involvement from baseline</li> <li>• Reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Dermatologist</li> <li>• Allergist</li> <li>• Immunologist</li> </ul>	

## 2 . Background

Benefit/Coverage/Program Information			
Table 1. Relative potencies of topical corticosteroids [2]			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
	Amcinonide	Cream, lotion, ointment	0.1

High Potency	Augmented betamethasone dipropionate	Cream	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05
	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream	0.1
Triamcinolone acetonide	Cream, ointment	0.1	
Lower- medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05

	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

**3 . Revision History**

Date	Notes
12/6/2024	New program

Ecoza (econazole)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99550
<b>Guideline Name</b>	Ecoza (econazole)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name:Ecoza, Generic econazole	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, contraindication, or intolerance to ALL of the following: <ul style="list-style-type: none"><li>butenafine</li><li>ciclopirox</li></ul>	

- clotrimazole
- clotrimazole w/ betamethasone
- ketoconazole
- miconazole
- nystatin
- terbinafine
- tolnaftate

## 2 . Revision History

Date	Notes
6/10/2021	Update guideline

Eculizumab (Soliris, Bkembv, Epysqli)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325191
<b>Guideline Name</b>	Eculizumab (Soliris, Bkembv, Epysqli)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Epysqli	
Diagnosis	Paroxysmal nocturnal hemoglobinuria (PNH)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of paroxysmal nocturnal hemoglobinuria (PNH)	

**AND**

**2 - BOTH** of the following:

- Flow cytometry analysis confirming presence of PNH clones
- Laboratory results, signs, and/or symptoms attributed to PNH (e.g., abdominal pain, anemia, dyspnea, extreme fatigue, smooth muscle dystonia, unexplained/unusual thrombosis, hemolysis/hemoglobinuria, kidney disease, pulmonary hypertension, etc.)

**AND**

**3 - Patient is treatment naïve** with eculizumab therapy

**AND**

**4 - Requested medication is dosed according to the United States Food and Drug Administration (FDA) labeled dosing for PNH**

**AND**

**5 - Prescribed by, or in consultation with, ONE** of the following:

- Hematologist
- Oncologist

<b>Product Name:Epysqli</b>	
Diagnosis	Paroxysmal nocturnal hemoglobinuria (PNH)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) demonstrating a positive clinical response from baseline (e.g., increased or stabilization of hemoglobin levels, reduction in transfusions, improvement in hemolysis, decrease in lactate dehydrogenase [LDH], increased reticulocyte count, etc.)

**AND**

2 - Requested medication is dosed according to the United States Food and Drug Administration (FDA) labeled dosing for paroxysmal nocturnal hemoglobinuria (PNH)

**AND**

3 - Prescribed by, or in consultation with, ONE of the following:

- Hematologist
- Oncologist

Product Name:Epysqli	
Diagnosis	Atypical hemolytic uremic syndrome (aHUS)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) supporting a diagnosis of atypical hemolytic uremic syndrome (aHUS) by ruling out BOTH of the following:	
<ul style="list-style-type: none"><li>• Shiga toxin E. coli-related hemolytic uremic syndrome (STEC-HUS)*</li><li>• Thrombotic thrombocytopenia purpura (TTP) (e.g., rule out ADAMTS13 deficiency)</li></ul>	
<b>AND</b>	
2 - Laboratory results, signs, and/or symptoms attributed to aHUS (e.g., thrombocytopenia, microangiopathic hemolysis, thrombotic microangiopathy, acute renal failure, etc.)	

**AND**

**3** - Patient is treatment naïve with eculizumab therapy

**AND**

**4** - Requested medication is dosed according to the Food and Drug Administration (FDA) labeled dosing for aHUS

**AND**

**5** - Prescribed by, or in consultation with, a hematologist or nephrologist

Product Name:Epysqli	
Diagnosis	Atypical hemolytic uremic syndrome (aHUS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes) demonstrating a positive clinical response from baseline (e.g., reduction of plasma exchanges, reduction of dialysis, increased platelet count, reduction of hemolysis)	
<b>AND</b>	
<b>2</b> - Requested medication is dosed according to the United States Food and Drug Administration (FDA) labeled dosing for atypical hemolytic uremic syndrome (aHUS)	
<b>AND</b>	

3 - Prescribed by, or in consultation with, a hematologist or nephrologist

Product Name:Epysqli

Diagnosis	Generalized myasthenia gravis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming ALL of the following:

1.1 Diagnosis of generalized myasthenia gravis (gMG)

**AND**

1.2 Patient has not failed a previous course of eculizumab therapy

**AND**

1.3 Positive serologic test for anti-acetylcholine receptor (AChR) antibodies

**AND**

1.4 ONE of the following:

- History of abnormal neuromuscular transmission test demonstrated by single-fiber electromyography (SFEMG) or repetitive nerve stimulation
- History of positive anticholinesterase test, e.g., edrophonium chloride test
- Patient has demonstrated improvement in myasthenia gravis (MG) signs on oral cholinesterase inhibitors, as assessed by the treating neurologist

**AND**

**1.5** Patient has a Myasthenia Gravis Foundation of America (MGFA) Clinical Classification of class II, III, or IV at initiation of therapy

**AND**

**1.6** Patient has a Myasthenia Gravis-specific Activities of Daily Living scale (MG-ADL) total score greater than or equal to 6 at initiation of therapy

**AND**

**2** - Patient is 6 years of age or older

**AND**

**3** - ONE of the following:

**3.1** For patients between 6 and 17 years of age, trial and failure, contraindication, or intolerance to one of the following:

- Immunosuppressive therapy (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus)
- Chronic plasmapheresis or plasma exchange (PE)
- Intravenous immunoglobulin (IVIG)

**AND**

**3.2** For patients 18 years of age or older, BOTH of the following:

- History of failure of at least TWO immunosuppressive agents over the course of at least 12 months [e.g., azathioprine, methotrexate, cyclosporine, mycophenolate, etc.]
- Patient has required TWO or more courses of plasmapheresis/plasma exchanges and/or intravenous immune globulin for at least the previous 12 months without symptom control

**AND**

**4** - Patient is currently on a stable therapeutic dose (at least 3 to 6 months) of immunosuppressive therapy

**AND**

**5** - Requested medication is initiated and titrated according to the United States Food and Drug Administration (FDA) labeled dosing for gMG: up to a maximum of 1200 milligrams every 2 weeks

**AND**

**6** - Prescribed by, or in consultation, with a neurologist

Product Name:Epysqli	
Diagnosis	Generalized myasthenia gravis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) demonstrating a positive clinical response from baseline as evidenced by ALL of the following:

- Improvement and/or maintenance of at least a 3 point improvement (reduction in score) in the Myasthenia Gravis Activities of Daily Living (MG-ADL) score from pre-treatment baseline
- Reduction in signs and symptoms of myasthenia gravis
- Maintenance, reduction, or discontinuation of dose(s) of baseline immunosuppressive therapy (IST) prior to starting eculizumab therapy (Note: Add on, dose escalation of IST, or additional rescue therapy from baseline to treat myasthenia gravis or exacerbation of symptoms while on eculizumab therapy will be considered as treatment failure)

**AND**

**2** - Requested medication is dosed according to the United States Food and Drug Administration (FDA) labeled dosing for generalized myasthenia gravis (gMG): up to a maximum of 1200 milligrams every 2 weeks

**AND**

**3** - Prescribed by, or in consultation, with a neurologist

Product Name:Epysqli	
Diagnosis	Neuromyelitis optica spectrum disorder (NMOSD)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming ALL of the following:</p> <p>1.1 Diagnosis of neuromyelitis optica spectrum disorder (NMOSD)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Past medical history of ONE of the following:</p> <ul style="list-style-type: none"><li>• Optic neuritis</li><li>• Acute myelitis</li><li>• Area postrema syndrome: Episode of otherwise unexplained hiccups or nausea and vomiting</li><li>• Acute brainstem syndrome</li><li>• Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions</li><li>• Symptomatic cerebral syndrome with NMOSD-typical brain lesions</li></ul> <p style="text-align: center;"><b>AND</b></p> <p>1.3 Positive serologic test for anti-aquaporin-4 immunoglobulin G (AQP4-IgG)/NMO-IgG antibodies</p>	

**AND**

**1.4** Diagnosis of multiple sclerosis or other diagnoses have been ruled out

**AND**

**2** - Patient has not failed a previous course of eculizumab therapy

**AND**

**3** - History of failure of, contraindication, or intolerance to rituximab (Ruxience, Truxima) therapy

**AND**

**4** - One of the following:

**4.1** History of at least two relapses during the previous 12 months prior to initiating eculizumab therapy

**OR**

**4.2** History of at least three relapses during the previous 24 months, at least one relapse occurring within the past 12 months prior to initiating eculizumab therapy

**AND**

**5** - Requested medication is initiated and titrated according to the U.S. FDA labeled dosing for NMOSD, up to a maximum of 1200 mg every 2 weeks

**AND**

**6** - Prescribed by, or in consultation with, a neurologist

Product Name:Epysqli	
Diagnosis	Neuromyelitis optica spectrum disorder (NMOSD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) demonstrating a positive clinical response from baseline as evidenced by BOTH of the following:</p> <p><b>1.1</b> Reduction in the number and/or severity of relapses or signs and symptoms of neuromyelitis optica spectrum disorder (NMOSD)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> Maintenance, reduction, or discontinuation of dose(s) of any baseline immunosuppressive therapy (IST) prior to starting eculizumab therapy. (Note: Add on, dose escalation of IST, or additional rescue therapy from baseline to treat NMOSD or exacerbation of symptoms while on eculizumab therapy will be considered as treatment failure)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Requested medication is dosed according to the U.S. FDA (Food and Drug Administration) labeled dosing for NMOSD: up to a maximum of 1200 mg every 2 weeks</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by, or in consultation with, a neurologist</p>	

Product Name:Non-Preferred*: Soliris, Bkemy, and newly launched eculizumab products	
Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred product is Epysqli (eculizumab-aagh). \*\*Please note: The drug(s) listed above may require additional review.

Notes

\*Patients must use preferred eculizumab biosimilar.

**2 . Revision History**

Date	Notes
7/16/2025	Updated preferred/npd status for all targets, updated NPD section. Updated NP section verbiage.

Egrifta

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99606
<b>Guideline Name</b>	Egrifta
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Egrifta SV	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of human immunodeficiency virus (HIV)-associated lipodystrophy	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona SP to Medicaid Arizona SP for 7/1 eff

Elaprase - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99607
<b>Guideline Name</b>	Elaprase - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Elaprase	
Diagnosis	Hunter syndrome
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Hunter syndrome (Mucopolysaccharidosis II, MPS II)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona SP to Medicaid Arizona SP for 7/1 eff

Elevidys (delandistrogene moxeparvovec-rokl)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269207
<b>Guideline Name</b>	Elevidys (delandistrogene moxeparvovec-rokl)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Elevidys	
Diagnosis	Duchenne Muscular Dystrophy (DMD)
Approval Length	1 Time Authorization in Lifetime
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of Duchenne Muscular Dystrophy (DMD)	

**AND**

**2** - Mutation in the DMD gene has been confirmed

**AND**

**3** - No deletion in exon 8 or exon 9 in the DMD gene is present

**AND**

**4** - Patient is 4 years of age or older

**AND**

**5** - Patient is ambulatory without needing an assistive device (e.g., without side-by-side assist, cane, walker, wheelchair, etc.)

**AND**

**6** - Anti-AAVrh74 total binding antibody titers are less than 1:400

**AND**

**7** - Patient will receive a corticosteroid regimen prior to and following the administration of Elevidys in line with the FDA-approved recommendations in the labeling

**AND**

**8** - Provider attests that all of the following laboratory values have been checked prior to therapy and will be monitored after administration according to the FDA-approved recommendations in the labeling:

- Liver function (i.e., clinical exam, GGT, total bilirubin)

- Platelet counts
- Troponin-I

**AND**

**9** - Patient has a left ventricular ejection fraction of greater than or equal to 40 percent (%)

**AND**

**10** - Patient does not have clinical signs or symptoms of infection

**AND**

**11** - Patient will not receive exon-skipping therapies for DMD [e.g., Amondys (casimersen), Exondys 51 (eteplirsen), Viltepso (viltolarsen), Vyondys 53 (golodirsen)] concomitantly or following Elevidys treatment

**AND**

**12** - Prescribed by a neurologist with expertise in the treatment of DMD at an authorized treatment center with expertise in gene therapy

**AND**

**13** - Patient has never received Elevidys treatment in their lifetime

## 2 . Revision History

Date	Notes
5/29/2025	Added GPI 74600030406470, no change to clinical criteria.

Elidel (pimecrolimus) cream, tacrolimus ointment

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152812
<b>Guideline Name</b>	Elidel (pimecrolimus) cream, tacrolimus ointment
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

### 1 . Criteria

Product Name:Brand Elidel, generic pimecrolimus, generic tacrolimus 0.03%, generic tacrolimus 0.1%	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The patient is 2 years of age or older  <b>AND</b>	

**2 - ONE of the following:**

**2.1** History of failure, contraindication, or intolerance to ONE topical corticosteroid in the past 90 days

**OR**

**2.2** Drug is being prescribed for the facial or groin area

## **2 . Revision History**

Date	Notes
9/24/2024	Updated guideline name. Removed Brand Protopic as target, all other drugs on PA require age/step through topical corticosteroid

Elmiron

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99448
<b>Guideline Name</b>	Elmiron
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Elmiron	
Diagnosis	Bladder pain or discomfort associated with interstitial cystitis
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Patient has a documented diagnosis of bladder pain or discomfort associated with interstitial cystitis	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Emflaza (deflazacort)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144631
<b>Guideline Name</b>	Emflaza (deflazacort)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Brand Emflaza, generic deflazacort	
Diagnosis	Duchenne Muscular Dystrophy
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Duchenne muscular dystrophy	

**AND**

**2** - Patient is 2 years of age or older

**AND**

**3** - History of failure, contraindication, or intolerance to ONE of the following for the treatment of Duchenne muscular dystrophy:

- Prednisone
- Prednisolone

**AND**

**4** - Prescribed by or in consultation with a neurologist

**AND**

**5** - If the request is for generic deflazacort, patient must have tried and failed Brand Emflaza

Product Name:Brand Emflaza, generic deflazacort	
Diagnosis	Duchenne Muscular Dystrophy
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Physician attestation that the patient has had a positive clinical response to therapy	
<b>AND</b>	

2 - If the request is for generic deflazacort, patient must have tried and failed Brand Emflaza

## 2 . Revision History

Date	Notes
3/19/2024	Updated guideline name, added step through preferred Brand Emflaza for generic deflazacort

Emrosi (minocycline)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164415
<b>Guideline Name</b>	Emrosi (minocycline)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Emrosi	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Medication will be used for treatment of inflammatory lesions (papules and pustules) of rosacea  <b>AND</b>	

2 - History of failure to generic minocycline 50mg capsules

## 2 . Revision History

Date	Notes
1/30/2025	New program

Enbrel (etanercept)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-139348
<b>Guideline Name</b>	Enbrel (etanercept)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Enbrel	
Diagnosis	Moderately to Severely Active Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active Rheumatoid Arthritis (RA)	

**AND**

**2** - History of failure to a 3 month trial of ONE non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial)\*

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Notes

\*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial

Product Name:Enbrel

Diagnosis Moderately to Severely Active Rheumatoid Arthritis (RA)

Approval Length 12 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to Enbrel therapy

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Enbrel

Diagnosis Moderately to Severely Active Polyarticular Juvenile Idiopathic Arthritis

Approval Length 12 month(s)

Therapy Stage Initial Authorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is 2 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a rheumatologist</p>	

Product Name:Enbrel	
Diagnosis	Moderately to Severely Active Polyarticular Juvenile Idiopathic Arthritis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Enbrel therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p>	

Product Name:Enbrel	
Diagnosis	Active Psoriatic Arthritis

Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of active psoriatic arthritis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial)*</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with ONE of the following:</p> <ul style="list-style-type: none"> <li>• Rheumatologist</li> <li>• Dermatologist</li> </ul>	
Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial

Product Name:Enbrel	
Diagnosis	Active Psoriatic Arthritis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Enbrel therapy</p>	

**AND**

**2** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Product Name:Enbrel	
Diagnosis	Moderate to Severe Chronic Plaque Psoriasis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of moderate to severe chronic plaque psoriasis</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Greater than or equal to 3% body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Both of the following:</p> <p><b>3.1</b> History of failure to one of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial):*</p> <ul style="list-style-type: none"><li>• Corticosteroids (e.g., betamethasone, clobetasol, desonide)</li><li>• Vitamin D analogs (e.g., calcitriol, calcipotriene)</li><li>• Tazarotene</li><li>• Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)</li><li>• Anthralin</li></ul>	

- Coal tar

**AND**

**3.2** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial)\*

**AND**

**4** - Prescribed by or in consultation with a dermatologist

Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial
-------	--

Product Name:Enbrel	
Diagnosis	Moderate to Severe Chronic Plaque Psoriasis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Enbrel therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a dermatologist</p>	

Product Name:Enbrel	
Diagnosis	Ankylosing spondylitis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of active ankylosing spondylitis

**AND**

2 - History of failure to two non-steroidal anti-inflammatory drugs (NSAIDs: e.g., ibuprofen, naproxen) at maximally indicated doses, each used for at least 4 weeks within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trials)\*

**AND**

3 - Prescribed by or in consultation with a rheumatologist

Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial
-------	--

Product Name:Enbrel	
Diagnosis	Ankylosing Spondylitis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Enbrel therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a rheumatologist	

## 2 . Revision History

Date	Notes
1/26/2024	Added age criteria for PJIA indication, removed concomitant use safety criterion from all sections

Encelto (revakinagene taroretcel-lwey)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-274200
<b>Guideline Name</b>	Encelto (revakinagene taroretcel-lwey)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Encelto	
Approval Length	1 Time Authorization per Eye in Lifetime
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of macular telangiectasia type 2 (MacTel)  <b>AND</b>	

**2** - Provider attests patient has not received prior treatment with Encelto in the affected eye in their lifetime

**AND**

**3** - Prescribed by or in consultation with an ophthalmologist experienced in the treatment of retinal diseases

## **2 . Revision History**

Date	Notes
5/29/2025	New program

Endari

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99450
<b>Guideline Name</b>	Endari
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Endari	
Diagnosis	Sickle cell disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - BOTH of the following: <ul style="list-style-type: none"><li>Diagnosis of sickle cell disease</li></ul>	

- Used to reduce acute complications of sickle cell disease

**AND**

**2 - ONE of the following:**

- Patient is using Endari with concurrent hydroxyurea therapy
- Patient is unable to take hydroxyurea due to a contraindication or intolerance

**AND**

**3 - Patient has had 2 or more painful sickle cell crises within the past 12 months**

Product Name:Endari	
Diagnosis	Sickle cell disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Endari therapy</p>	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Enjaymo (sutimlimab-jome)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-123730
<b>Guideline Name</b>	Enjaymo (sutimlimab-jome)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Enjaymo	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of cold agglutinin disease (CAD) based on ALL of the following:	

- Presence of chronic hemolysis (e.g., bilirubin level above the normal reference range, elevated lactated dehydrogenase [LDH], decreased haptoglobin, increased reticulocyte count)
- Positive polyspecific direct antiglobulin test (DAT)
- Monospecific DAT strongly positive for C3d
- Cold agglutinin titer greater than or equal to 64 measured at 4 degree celsius
- Direct antiglobulin test (DAT) result for Immunoglobulin G (IgG) of 1 plus or less

**AND**

**2** - Patient does not have cold agglutinin syndrome secondary to other factors (e.g., overt hematologic malignancy, primary immunodeficiency, infection, rheumatologic disease, systemic lupus erythematosus or other autoimmune disorders)

**AND**

**3** - Baseline hemoglobin level less than or equal to 10.0 gram per deciliter (g/dL)

**AND**

**4** - One of the following:

- Prescribed dose will not exceed 6,500 mg on day 0, 7, and every 14 days thereafter for patients weighing between 39 kg to less than 75 kg
- Prescribed dose will not exceed 7,500 mg on day 0, 7, and every 14 days thereafter for patients for patients weighing 75 kg or greater

**AND**

**5** - Prescribed by or in consultation with a hematologist

Product Name:Enjaymo	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Submission of medical records (e.g., chart notes) demonstrating positive clinical response to therapy as evidenced by ALL of the following:

- The patient has not required any blood transfusions after the first 5 weeks of therapy with Enjaymo
- Hemoglobin level greater than or equal to 12 gram per deciliter (g/dL) or increased greater than or equal to 2 g/dL from baseline

**AND**

2 - One of the following:

- Prescribed dose will not exceed 6,500 mg on day 0, 7, and every 14 days thereafter for patients weighing between 39 kg to less than 75 kg
- Prescribed dose will not exceed 7,500 mg on day 0, 7, and every 14 days thereafter for patients for patients weighing 75 kg or greater

**AND**

3 - Prescribed by or in consultation with a hematologist

## 2 . Background

Clinical Practice Guidelines	
<b>Weight-Based Dosing</b>	
The dosing is 6,500mg or 7,500mg Enjaymo (based on body weight) intravenously over approximately 60 minutes on Day 0, Day 7, and every 14 days thereafter	
Body Weight Range	Dose
39kg to less than 75kg	6,500 mg

75kg or greater	7,500 mg
-----------------	----------

**3 . Revision History**

Date	Notes
3/23/2023	New program

Entocort EC

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99451
<b>Guideline Name</b>	Entocort EC
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Entocort EC, generic budesonide	
Diagnosis	Chrohn's Disease
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Entocort EC is being used for the treatment of Crohn's disease	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Entresto

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157622
<b>Guideline Name</b>	Entresto
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

### 1 . Criteria

Product Name:Entresto tablet, Entresto sprinkle capsule	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - As continuation of therapy initiated during an inpatient stay  <b>OR</b>	

**2 - Both of the following:**

**2.1** Diagnosis of pediatric heart failure with systemic left ventricular systolic dysfunction which is symptomatic

**AND**

**2.2** Prescribed by or in consultation with a cardiologist

**OR**

**3 - ALL of the following:**

**3.1** Diagnosis of heart failure (with or without hypertension)

**AND**

**3.2** Ejection fraction is less than or equal to 40 percent

**AND**

**3.3** Heart failure is classified as ONE of the following:

- New York Heart Association Class II
- New York Heart Association Class III
- New York Heart Association Class IV

**AND**

**3.4** ONE of the following:

**3.4.1** Patient is on a stabilized dose and receiving concomitant therapy with ONE of the following beta-blockers:

- bisoprolol
- carvedilol
- metoprolol

**OR**

**3.4.2** Patient has a contraindication or intolerance to beta-blocker therapy

**AND**

**3.5** Patient does not have a history of angioedema

**AND**

**3.6** Patient will discontinue any use of concomitant ACE (angiotensin converting enzyme) Inhibitor or ARB (angiotensin II receptor blocker) before initiating treatment with Entresto

**AND**

**3.7** Patient is not concomitantly on aliskiren therapy

**AND**

**3.8** Entresto is prescribed by, or in consultation with, a cardiologist

Product Name:Entresto tablet, Entresto sprinkle capsule	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - The Entresto dose has been titrated to a dose of 97 mg (milligrams) /103 mg twice daily, or to a maximum dose as tolerated by the patient	

**AND**

**2** - Documentation of positive clinical response to therapy

## **2 . Revision History**

Date	Notes
10/25/2024	Removed ops note regarding ACEI discontinuation

Entyvio (vedolizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300286
<b>Guideline Name</b>	Entyvio (vedolizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Entyvio IV	
Diagnosis	Crohn's Disease (CD)
Approval Length	14 Weeks
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's disease	

**AND**

**2** - One of the following:

- Frequent diarrhea and abdominal pain
- At least 10% weight loss
- Complications such as obstruction, fever, abdominal mass
- Abnormal lab values (e.g., C-reactive protein [CRP])
- CD Activity Index (CAI) greater than 220

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

- 6-mercaptopurine
- azathioprine
- corticosteroids (e.g., prednisone)
- methotrexate

**AND**

**4** - One of the following:

**4.1** Paid claims or submission of medical records (e.g., chart notes) confirming history of failure, contraindication, or intolerance to ALL of the following :

- A preferred adalimumab biosimilar
- infliximab
- A preferred ustekinumab biosimilar

**OR**

**4.2** Paid claims or submission of medical records (e.g., chart notes) confirming continuation of prior Entyvio therapy, defined as no more than a 45-day gap in therapy

**AND**

5 - Prescribed by or in consultation with a gastroenterologist

Product Name:Entyvio IV

Diagnosis	Crohn's Disease (CD)
-----------	----------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:

- Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline
- Reversal of high fecal output state

**AND**

2 - Prescribed by or in consultation with a gastroenterologist

Product Name:Entyvio IV

Diagnosis	Ulcerative Colitis (UC)
-----------	-------------------------

Approval Length	4 Week(s)
-----------------	-----------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis

**AND**

**2** - One of the following:

- Greater than 6 stools per day
- Frequent blood in the stools
- Frequent urgency
- Presence of ulcers
- Abnormal lab values (e.g., hemoglobin, ESR, CRP)
- Dependent on, or refractory to, corticosteroids

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to **ONE** of the following conventional therapies:

- 6-mercaptopurine
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)
- Azathioprine
- Corticosteroids (e.g., prednisone)

**AND**

**4** - One of the following:

**4.1** Paid claims or submission of medical records (e.g., chart notes) confirming history of failure, contraindication, or intolerance to **ALL** of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**OR**

**4.2** Paid claims or submission of medical records (e.g., chart notes) confirming continuation of prior Entyvio therapy, defined as no more than a 45-day gap in therapy

**AND**

**5** - Prescribed by or in consultation with a gastroenterologist

**AND**

**6** - Entyvio IV formulation will be used for induction purposes only and patient will be switched to the Entyvio SC (subcutaneous) formulation for week 6 dose

Product Name:Entyvio IV	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	N/A - Requests for Entyvio IV should be denied
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Requests for continuing Entyvio IV therapy should be denied. The plan's preferred product is Entyvio SC (SC will require PA)	
Notes	Requests for continuing Entyvio IV therapy should be denied. The plan's preferred product for UC indication is Entyvio SC (SC will require PA)

Product Name:Entyvio SC	
Diagnosis	Crohn's Disease (CD), Ulcerative Colitis (UC)
Approval Length	14 Week(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1 - One of the following:**

**1.1 All of the following:**

**1.1.1** Submission of medical records (e.g., chart notes) confirming a diagnosis of one of the following:

- Moderately to severely active Crohn's disease
- Moderately to severely active ulcerative colitis

**AND**

**1.1.2** Paid claims or submission of medical records (e.g., chart notes) confirming ONE of the following:

**1.1.2.1** Will be used as a maintenance dose following two doses of Entyvio IV\* for induction

**OR**

**1.1.2.2** Patient is currently established on Entyvio IV\*

**AND**

**1.1.3** Prescribed by or in consultation with a gastroenterologist

**OR**

**1.2** Patient has received 2 doses of Entyvio IV\* for induction

Notes

\* This product will require prior authorization

Product Name:Entyvio SC	
Diagnosis	Crohn's Disease (CD), Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:

- Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline
- Reversal of high fecal output state

## 2 . Revision History

Date	Notes
7/3/2025	Updated preferred agents/embedded steps, updated criteria throughout.

Eohilia (budesonide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-146018
<b>Guideline Name</b>	Eohilia (budesonide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Eohilia	
Diagnosis	Eosinophilic Esophagitis (EoE)
Approval Length	12 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of eosinophilic esophagitis (EoE)  <b>AND</b>	

**2** - Patient has symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, heartburn, abdominal pain)

**AND**

**3** - Patient has at least 15 intraepithelial eosinophils per high power field (HPF)

**AND**

**4** - Other causes of esophageal eosinophilia have been excluded

**AND**

**5** - Patient is 11 years of age or older

**AND**

**6** - Paid claims or submission of medical records (e.g., chart notes) confirming trial and failure (of a minimum 8-week duration), contraindication, or intolerance to a proton pump inhibitor (e.g., pantoprazole, omeprazole)

**AND**

**7** - Paid claims or submission of medical records (e.g., chart notes) confirming trial and failure (of a minimum 8-week duration), or intolerance to a topical (esophageal) corticosteroid (e.g., budesonide, fluticasone)

**AND**

**8** - Prescribed by or in consultation with one of the following:

- Allergist/Immunologist
- Gastroenterologist

## 2 . Revision History

Date	Notes
4/23/2024	New program

Epaned

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99453
<b>Guideline Name</b>	Epaned
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Epaned	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Patient is less than 8 years of age</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2 BOTH of the following:**

**1.2.1 ONE of the following diagnoses:**

- Hypertension
- Heart failure
- Asymptomatic left ventricular dysfunction, defined as left ventricular ejection fraction less than or equal to 35%

**AND**

**1.2.2 ONE of the following:**

**1.2.2.1** History of failure, contraindication, or intolerance to TWO formulary oral anti-hypertensives (e.g., angiotensin-converting enzyme (ACE) inhibitor, ACE inhibitor combination, angiotensin-receptor blockers (ARB), ARB combination, thiazide diuretic)

**OR**

**1.2.2.2** Patient is unable to ingest a solid dosage form (e.g. an oral tablet or capsule) due to ONE of the following:

- Oral/motor difficulties
- Dysphagia

## **2 . Revision History**

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Epinephrine Products

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-269193
<b>Guideline Name</b>	Epinephrine Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	6/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Epinephrine Pens (Non-Mylan Manufacturer)*, Neffy	
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - There is a shortage on Epinephrine Pens manufactured by Mylan.	
Notes	*Only approve other rebatable epinephrine autoinjectors if both the branded EpiPen and authorized generic are on the FDA shortage list.

Product Name:Epinephrine Pens (Mylan Manufacturer)
--

Approval Length	6 month(s)
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - Medication has been used or lost or the member is going on vacation.*</p>	
Notes	*Only approve other rebatable epinephrine autoinjectors if both the branded EpiPen and authorized generic are on the FDA shortage list

## 2 . Revision History

Date	Notes
5/29/2025	Added new Neffy GPI

Eplerenone- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99454
<b>Guideline Name</b>	Eplerenone- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Inspra, generic eplerenone	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of one of the following:  1.1 Symptomatic heart failure with reduced ejection fraction (HFrEF) after an acute myocardial infarction	

OR

1.2 Hypertension

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Epsolay (benzoyl peroxide) cream

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-108675
<b>Guideline Name</b>	Epsolay (benzoyl peroxide) cream
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2022
-----------------	----------

### 1 . Criteria

Product Name:Epsolay	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of rosacea  <b>AND</b>  2 - Patient has inflammatory lesions	

**AND**

**3** - Trial and failure (of a minimum 30-day supply), contraindication or intolerance to one preferred topical product for rosacea (e.g., metronidazole cream/gel/lotion) (verified via paid pharmacy claims)

## **2 . Revision History**

Date	Notes
6/24/2022	New Program

Erythropoietic Agents - AZM

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-325192
<b>Guideline Name</b>	Erythropoietic Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	8/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Preferred: Retacrit	
Diagnosis	Anemia Due to Chronic Kidney Disease (CKD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of chronic kidney disease (CKD)	

**AND**

**2** - Hematocrit is less than 30% at initiation of therapy

**AND**

**3** - ONE of the following:

**3.1** Patient is on dialysis

**OR**

**3.2** ALL of the following:

**3.2.1** Patient is NOT on dialysis

**AND**

**3.2.2** The rate of hematocrit decline indicates the likelihood of requiring a red blood cell (RBC) transfusion

**AND**

**3.2.3** Reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal

Product Name:Preferred: Retacrit	
Diagnosis	Anemia Due to Chronic Kidney Disease (CKD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

## **Approval Criteria**

1 - Diagnosis of chronic kidney disease (CKD)

**AND**

2 - ONE of the following:

2.1 Both of the following:

- Patient is on dialysis
- Most recent or average Hct over 3 months is 33% or less (Hgb 11 g/dL or less)

**OR**

2.2 All of the following:

- Patient is NOT on dialysis
- Most recent or average (avg) Hct over 3 mo is 30% or less (Hgb 10 g/dL or less)
- Reducing the risk of alloimmunization and/or other RBC transfusion-related risks is a goal

**OR**

2.3 Both of the following:

- Request is for a pediatric patient
- Most recent or average Hct over 3 mo is 36% or less (Hgb 12 g/dL or less)

**AND**

3 - One of the following:

- Decrease in the need for blood transfusion
- Hemoglobin (Hgb) increased greater than or equal to 1g/dL from pre-treatment level

Product Name:Preferred: Retacrit	
Diagnosis	Anemia Associated with Zidovudine Treatment in HIV-Infected Patients
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is receiving zidovudine administered at less than or equal to 4200 milligrams per week</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Endogenous serum erythropoietin level is less than or equal to 500 milliunits per milliliter</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Hematocrit is less than 30% at initiation of therapy</p>	

Product Name:Preferred: Retacrit	
Diagnosis	Anemia Due to Cancer Chemotherapy
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Hematocrit less than 30% at initiation of therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - There is a minimum of two additional months of planned chemotherapy</p>	

Product Name:Preferred: Retacrit	
Diagnosis	Preoperative Use for Reduction of Allogeneic Blood Transfusions in Surgery Patients
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Perioperative hematocrit is greater than 30% and less than or equal to 39%</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is at high risk for blood loss during surgery</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is unable or unwilling to donate autologous blood</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Surgery procedure is elective, non-cardiac, and non-vascular</p>	

Product Name:Preferred: Retacrit	
Diagnosis	Anemia Associated with Myelodysplastic Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of myelodysplastic disease (MDS)</p>	

**AND**

**2 - ONE of the following:**

- Serum erythropoietin level less than or equal to 500 milliunits per milliliter
- Hematocrit is less than or equal to 30% at the initiation of therapy

Product Name:Preferred: Retacrit	
Diagnosis	Anemia Associated with Myelodysplastic Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - One of the following:	
1.1 Hematocrit remains less than 36%	
<b>OR</b>	
1.2 Patient has demonstrated a response to therapy	

Product Name:Preferred: Retacrit	
Diagnosis	Anemia in Patients with Hepatitis C with Ribavirin and Interferon Therapy
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of hepatitis C virus (HCV) infection

**AND**

2 - Patient is receiving ribavirin and interferon therapy

**AND**

3 - Hematocrit is less than or equal to 30% at initiation of therapy

Product Name:Preferred: Retacrit\*

Diagnosis	Anemia in Patients with Hepatitis C with Ribavirin and Interferon Therapy
-----------	---

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - One of the following:

1.1 Hematocrit remains less than 36%

**OR**

1.2 Patient has demonstrated a response to therapy

Notes	*NOTE: Authorization will be issued for 12 months or if patient has demonstrated response to therapy, authorization will be issued for the full course of ribavirin therapy.
-------	--

Product Name:Preferred: Retacrit\*

Diagnosis	Erythropoietin Stimulating Agents –Off-Label Uses
-----------	---

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Off-label requests will be evaluated on a case-by-case basis by a clinical pharmacist

**AND**

2 - Requests for coverage in patients with hemoglobin (Hgb) greater than 10 grams per deciliter or hematocrit (Hct) greater than 30% will not be approved

Notes	*If the request is deemed medically necessary, the authorization will be issued for requested length of therapy.
-------	--

Product Name: Non-Preferred\*: Aranesp, Epogen, Mircera, Procrit, and newly launched erythropoietin stimulating agents

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
-----------------	--

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred product is Retacrit. \*\*Please note: The drug(s) listed above may require additional review.

Notes	*Patients must use preferred erythropoietin stimulating agent (Retacrit).
-------	---

**2 . Revision History**

Date	Notes
7/16/2025	Updated preferred/NPD agents, updated criteria throughout. Updated NP section verbiage.

Esbriet, Ofev

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116139
<b>Guideline Name</b>	Esbriet, Ofev
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Brand Esbriet, generic pirfenidone, Brand Pirfenidone 534 mg tablets, Ofev	
Diagnosis	Idiopathic Pulmonary Fibrosis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of idiopathic pulmonary fibrosis (IPF) as documented by ALL of the following criteria:  1.1 Exclusion of other known causes of interstitial lung disease (e.g. domestic and	

occupational environmental exposures, connective tissue disease, and drug toxicity), as documented by the following:

- ICD-10 Code J84.112 (Idiopathic pulmonary fibrosis)

**AND**

**1.2 ONE** of the following:

**1.2.1** In patients NOT subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF

**OR**

**1.2.2** In patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern reveal IPF or probable IPF

**AND**

**2** - The agent is not being used in combination with Esbriet or Ofev

**AND**

**3** - The prescriber is a pulmonologist

**AND**

**4** - If requesting Brand or generic pirfenidone ONLY: patient has tried and failed, or has intolerance to Brand Esbriet

Product Name: Brand Esbriet, generic pirfenidone, Brand Pirfenidone 534 mg tablets, Ofev	
Diagnosis	Idiopathic Pulmonary Fibrosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - The agent is not being used in combination with Esbriet or Ofev</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - The prescriber is a pulmonologist</p>	

Product Name:Ofev	
Diagnosis	Systemic Sclerosis-Associated Interstitial Lung Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of systemic sclerosis (SSc) - associated interstitial lung disease as documented by ALL of the following:</p> <p>1.1 ONE of the following:</p> <p>1.1.1 Skin thickening of the fingers of both hands extending proximal to the metacarpophalangeal joints</p> <p style="text-align: center;"><b>OR</b></p> <p>1.1.2 TWO of the following:</p>	

- Skin thickening of the fingers (e.g., puffy fingers, sclerodactyly of the fingers)
- Fingertip lesions (e.g., digital tip ulcers, fingertip pitting scars)
- Telangiectasia
- Abnormal nailfold capillaries
- Pulmonary arterial hypertension
- Raynaud’s phenomenon
- SSc-related autoantibodies (e.g., anticentromere, anti-topoisomerase I, anti-RNA polymerase III)

**AND**

**1.2** Presence of interstitial lung disease as determined by finding evidence of pulmonary fibrosis on high-resolution computed tomography (HRCT), involving at least 10 percent of the lungs

**AND**

**2** - The agent is not being used in combination with Esbriet

**AND**

**3** - The prescriber is a pulmonologist

Product Name:Ofev	
Diagnosis	Chronic fibrosing interstitial lung disease with a progressive phenotype
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype as documented by BOTH of the following criteria:</p>	

**1.1** Presence of fibrotic ILD as determined by finding evidence of pulmonary fibrosis on HRCT (high-resolution computed tomography), involving at least 10 percent of the lungs

**AND**

**1.2** Patient is presenting with clinical signs of progression as defined by ONE of the following in the previous 24 months:

**1.2.1** Forced vital capacity (FVC) decline of greater than 10 percent

**OR**

**1.2.2** TWO of the following:

- FVC decline of greater than or equal to 5 percent, but less than 10 percent
- Patient is experiencing worsening respiratory symptoms
- Patient is exhibiting increasing extent of fibrotic changes on chest imaging

**AND**

**2** - The agent is not being used in combination with Esbriet

**AND**

**3** - The prescriber is a pulmonologist

Product Name:Ofev	
Diagnosis	Systemic Sclerosis-Associated Interstitial Lung Disease, Chronic fibrosing interstitial lung disease with a progressive phenotype
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to therapy

**AND**

2 - Ofev is not being used in combination with Esbriet

**AND**

3 - The prescriber is a pulmonologist

## 2 . Revision History

Date	Notes
10/27/2022	Added pirfenidone as NP target

Estrogens- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99455
<b>Guideline Name</b>	Estrogens- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Femring	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of moderate to severe vasomotor symptoms due to menopause  <b>OR</b>  2 - Diagnosis of moderate to severe vulvar and vaginal atrophy due to menopause	

Product Name: Premarin	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of atrophic vaginitis and kraurosis vulvae</p>	

## 2 . Revision History

Date	Notes
3/10/2021	Bulk Copied C&S Arizona standard to Arizona Medicaid for 7/1 effective

Eucrisa

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117636
<b>Guideline Name</b>	Eucrisa
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Eucrisa	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - BOTH of the following:  1.1 History of failure, contraindication, or intolerance to ONE topical corticosteroid [e.g., mometasone furoate, flucinolone acetonide (generic Synalar), fluocinonide]	

**AND**

**1.2** ONE of the following:

**1.2.1** Patient is less than 2 years of age

**OR**

**1.2.2** Patient is greater than or equal to 2 years of age and has history of failure, contraindication, or intolerance to ONE topical calcineurin inhibitor [e.g., pimecrolimus (generic Elidel), tacrolimus (generic Protopic)]

## **2 . Revision History**

Date	Notes
12/4/2022	Changed from ST to PA

Evkeeza (evinacumab-dgnb)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-124825
<b>Guideline Name</b>	Evkeeza (evinacumab-dgnb)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Evkeeza	
Diagnosis	Homozygous Familial Hypercholesterolemia [HoFH]
Approval Length	6 Months [A]
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient is 5 years of age or older	

**AND**

**2** - Submission of medical records (e.g. chart notes) documenting diagnosis of homozygous familial hypercholesterolemia (HoFH) as confirmed by one of the following:

**2.1** Genetic confirmation of two mutant alleles at the LDLR, APOB, PCSK9, or LDLRAP1 gene locus

**OR**

**2.2** Both of the following:

**2.2.1** One of the following:

- Untreated/pre-treatment LDL-C greater than 500 mg/dL
- Treated LDL-C greater than 300 mg/dL

**AND**

**2.2.2** One of the following:

- Xanthoma before 10 years of age
- Evidence of heterozygous familial hypercholesterolemia in both parents

**AND**

**3** - Submission of medical records (e.g., chart notes) demonstrating that patient has failed to achieve a low-density lipoprotein-cholesterol (LDL-C) goal of less than 100 mg/dL despite use of both of the following: \*Paid pharmacy claims may be used to confirm trial requirements

**3.1** One of the following:

**3.1.1** Patient is currently treated with maximally tolerated statin therapy plus ezetimibe

**OR**

**3.1.2** Patient is unable to tolerate statin therapy as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms: [B]

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

**OR**

**3.1.3** Patient has a labeled contraindication to all statins

**OR**

**3.1.4** Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations greater than 10 times ULN

**AND**

**3.2** One of the following:

- Patient has been treated with PCSK9 therapy or did not respond to PCSK9 therapy
- Physician attests that the patient is known to have two LDL-receptor negative alleles (little to no residual function) and therefore would not respond to PCSK9 therapy
- Patient has a history of intolerance or contraindication to PCSK9 therapy
- Patient has previously been treated with Juxtapid (lomitapide)
- Patient has previously been treated with lipoprotein apheresis

**AND**

**4** - Patient will continue other traditional lipid-lowering therapies (e.g., maximally tolerated statins, ezetimibe) in combination with Evkeeza

**AND**

**5** - Dose will not exceed 15 milligrams per kilogram of bodyweight infused once every 4 weeks

**AND**

**6** - Prescribed by one of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

Product Name:Evkeeza	
Diagnosis	Homozygous Familial Hypercholesterolemia [HoFH]
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting LDL-C reduction from baseline while on Evkeeza therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient will continue other traditional lipid-lowering therapies (e.g., maximally tolerated statins, ezetimibe) in combination with Evkeeza</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Dose will not exceed 15 milligrams per kilogram of bodyweight infused once every 4 weeks</p> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Prescribed by one of the following:</p> <ul style="list-style-type: none"><li>• Cardiologist</li><li>• Endocrinologist</li></ul>	

- Lipid specialist

## 2 . Endnotes

- A. Per the 2018 ACC/AHA national treatment guidelines, adherence, response to therapy, and adverse effects should be monitored within 4 -12 weeks following LDL-C lowering medication initiation or dose adjustment, repeated every 3 to 12 months as needed. Additionally, in the Evkeeza pivotal trial the primary outcome of change in LDL-C was evaluated at 24 weeks. [1,2,6]
- B. In patients treated with statins, it is recommended to measure creatine kinase levels in individuals with severe statin-associated muscle symptoms. [6]

## 3 . Revision History

Date	Notes
4/20/2023	New program

Evrysdi (risdiplam)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-224200
<b>Guideline Name</b>	Evrysdi (risdiplam)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Evrysdi	
Diagnosis	Spinal Muscular Atrophy (SMA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of spinal muscular atrophy (SMA)	

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) confirming the mutation or deletion of genes in chromosome 5q resulting in ONE of the following:

**2.1** Homozygous gene deletion or mutation of SMN1 gene (e.g., homozygous deletion of exon 7 at locus 5q13)

**OR**

**2.2** Compound heterozygous mutation of SMN1 gene [e.g., deletion of SMN1 exon 7 (allele 1) and mutation of SMN1 (allele 2)]

**AND**

**3** - Patient is not dependent on invasive ventilation or tracheostomy

**AND**

**4** - Patient is not dependent on the use of non-invasive ventilation beyond use for naps and nighttime sleep

**AND**

**5** - Patient is not receiving concomitant chronic survival motor neuron (SMN)-modifying therapy [e.g., Spinraza (nusinersen)]

**AND**

**6** - Patient has not previously received gene replacement therapy for the treatment of SMA [e.g., Zolgensma (onasemnogene abeparvovec-xioi)]

**AND**

**7** - Submission of medical records (e.g., chart notes, laboratory values) documenting the baseline assessment of at least ONE of the following exams (based on patient age and motor ability) to establish baseline motor ability (baseline motor function analysis could include assessments evaluated prior to receipt of previous chronic SMN-modifying therapy if transitioning therapy)\*:

- Children’s Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND)
- Hammersmith Infant Neurological Exam Part 2 (HINE-2)
- Hammersmith Functional Motor Scale Expanded (HF MSE)
- Upper Limb Module (ULM) Test
- Motor Function Measure 32 (MFM-32) Scale

**AND**

**8** - Prescribed by a neurologist with expertise in the treatment of SMA

Notes	*Baseline assessments for patients less than 2 months of age requesting Evrysdi are not necessary in order not to delay access to initial therapy in recently diagnosed infants. Initial assessments shortly post-therapy can serve as baseline with respect to efficacy reauthorization assessment.
-------	--

Product Name: Evrysdi	
Diagnosis	Spinal Muscular Atrophy (SMA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes, laboratory values) with the most recent results documenting a positive clinical response to Evrysdi compared to pretreatment baseline status [inclusive of baseline assessments prior to receipt of previous chronic survival motor neuron (SMN)-modifying therapy] as demonstrated by at least ONE of the following exams:</p> <p><b>1.1</b> Children’s Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND) with ONE of the following:</p> <p><b>1.1.1</b> Improvement or maintenance of previous improvement of at least a 4-point increase in score from pretreatment baseline</p>	

**OR**

**1.1.2** Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**OR**

**1.2** Hammersmith Infant Neurological Exam Part 2 (HINE-2) with ONE of the following:

**1.2.1** Improvement or maintenance of previous improvement of at least a 2-point (or maximal score) increase in ability to kick

**OR**

**1.2.2** Improvement or maintenance of previous improvement of at least a 1-point increase in any other HINE-2 milestone (e.g., head control, rolling, sitting, crawling, etc.), excluding voluntary grasp

**OR**

**1.2.3** The patient exhibited improvement, or maintenance of previous improvement, in more HINE motor milestones than worsening, from pretreatment baseline (net positive improvement)

**OR**

**1.2.4** Patient has achieved and maintained any new motor milestones when they would otherwise be unexpected to do so

**OR**

**1.3** Hammersmith Functional Motor Scale Expanded (HFMSE) with ONE of the following:

**1.3.1** Improvement or maintenance of previous improvement of at least a 3-point increase in score from pretreatment baseline

**OR**

**1.3.2** Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**OR**

**1.4** Upper Limb Module (ULM) with ONE of the following:

**1.4.1** Improvement or maintenance of previous improvement of at least a 2-point increase in score from pretreatment baseline

**OR**

**1.4.2** Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**OR**

**1.5** Motor Function Measure 32 (MFM-32) with ONE of the following:

**1.5.1** Improvement or maintenance of previous improvement of at least a 3-point increase in score from pretreatment baseline

**OR**

**1.5.2** Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**AND**

**2** - Patient is not dependent on invasive ventilation or tracheostomy

**AND**

**3** - Patient is not dependent on the use of non-invasive ventilation beyond use for naps and nighttime sleep

**AND**

**4** - Patient is not receiving concomitant chronic SMN-modifying therapy [e.g., Spinraza (nusinersen)]

**AND**

**5** - Patient has not previously received gene replacement therapy for the treatment of spinal muscular atrophy (SMA) [e.g., Zolgensma (onasemnogene abeparvovec-xioi)]

**AND**

**6** - Prescribed by a neurologist with expertise in the treatment of SMA

## 2 . Revision History

Date	Notes
3/26/2025	Added new GPI for Evrysdi tablet

Exondys - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116192
<b>Guideline Name</b>	Exondys - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Exondys	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Duchenne muscular dystrophy (DMD)  <b>AND</b>	

**2** - Documentation of a confirmed mutation of the dystrophin gene amenable to exon 51 skipping

**AND**

**3** - Prescribed by or in consultation with a neurologist who has experience treating Duchenne Muscular Dystrophy

**AND**

**4** - Dose will not exceed 30 milligrams per kilogram of body weight once weekly

**AND**

**5** - If ambulatory, patient's condition has been evaluated via the 6-minute walk test (6MWT) or North Star ambulatory assessment (NSAA) [documentation of the patient's most recent results must be provided]

Product Name: Exondys

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - One of the following:

**1.1** Patient has been on therapy for less than 12 months and all of the following:

**1.1.1** Patient is tolerating therapy

**AND**

**1.1.2** Dose will not exceed 30 milligrams per kilogram of body weight once weekly

**AND**

**1.1.3** Prescribed by or in consultation with a neurologist who has experience treating Duchenne Muscular Dystrophy

**AND**

**1.1.4** If ambulatory, patient's condition has been evaluated via the 6-minute walk test (6MWT) or North Star ambulatory assessment (NSAA) [documentation of the patient's most recent results must be provided]

**OR**

**1.2** Patient has been on therapy for 12 months or more and all of the following:

**1.2.1** Patient has experienced a benefit from therapy (e.g., disease amelioration compared to untreated patients)

**AND**

**1.2.2** Patient is tolerating therapy

**AND**

**1.2.3** Dose will not exceed 30 milligrams per kilogram of body weight once weekly

**AND**

**1.2.4** Prescribed by or in consultation with a neurologist who has experience treating Duchenne Muscular Dystrophy

**AND**

**1.2.5** If ambulatory, patient's condition has been evaluated via the 6-minute walk test

(6MWT) or North Star ambulatory assessment (NSAA) [documentation of the patient's most recent results must be provided]

## 2 . Revision History

Date	Notes
10/28/2022	Removed age and ambulatory requirements

Ezallor Sprinkle (rosuvastatin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-131964
<b>Guideline Name</b>	Ezallor Sprinkle (rosuvastatin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Ezallor	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Both of the following:  1.1.1 Patient is less than 10 years of age	

**AND**

**1.1.2** Prescribed by or in consultation with a cardiologist

**OR**

**1.2** Both of the following:

**1.2.1** Medication is being used for one of the following:

**1.2.1.1** To reduce the risk of one of the following:

- Myocardial infarction (MI), stroke, revascularization procedures, and angina in adults with multiple risk factors for coronary heart disease (CHD) but without clinically evident CHD
- MI and stroke in adults with type 2 diabetes mellitus with multiple risk factors for CHD but without clinically evident CHD
- Non-fatal MI, fatal and non-fatal stroke, revascularization procedures, hospitalization for congestive heart failure, and angina in adults with clinically evident CHD

**OR**

**1.2.1.2** As an adjunct to diet to reduce low-density lipoprotein cholesterol (LDL-C) in one of the following:

- Adults with primary hyperlipidemia
- Adults and pediatric patients aged 10 years and older with heterozygous familial hypercholesterolemia (HeFH)

**OR**

**1.2.1.3** As an adjunct to other LDL-C-lowering therapies, or alone if such treatments are unavailable, to reduce LDL-C in adults and pediatric patients aged 7 years and older with homozygous familial hypercholesterolemia (HoFH)

**OR**

**1.2.1.4** As an adjunct to diet for the treatment of adults with one of the following:

- Primary dysbetalipoproteinemia
- Hypertriglyceridemia

**AND**

**1.2.2** One of the following:

**1.2.2.1** Trial and failure, contraindication, or intolerance to generic rosuvastatin tablets (verified via paid pharmacy claims or submitted chart notes)

**OR**

**1.2.2.2** Patient is unable to swallow oral tablets

## 2 . Revision History

Date	Notes
8/29/2023	New program

Fabry Disease Agents

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-150054
<b>Guideline Name</b>	Fabry Disease Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Fabrazyme	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Fabry disease  <b>AND</b>	

2 - Patient is 2 years of age or older

**AND**

3 - Submission of medical records (e.g., chart notes) confirming ONE of the following:

- Detection of pathogenic mutations in the GLA gene by molecular genetic testing
- Deficiency in  $\alpha$ -galactosidase A ( $\alpha$ -Gal A) enzyme activity in plasma, isolated leukocytes, or dried blood spots (DBS)
- Significant clinical manifestations (e.g., neuropathic pain, cardiomyopathy, renal insufficiency, angiokeratomas, cornea verticillata)

**AND**

4 - Will not be used in combination with Galafold (migalastat)

Product Name:Elfabrio	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Fabry disease	
<b>AND</b>	
2 - Submission of medical records (e.g., chart notes) confirming ONE of the following:	
<ul style="list-style-type: none"><li>• Detection of pathogenic mutations in the GLA gene by molecular genetic testing</li><li>• Deficiency in <math>\alpha</math>-galactosidase A (<math>\alpha</math>-Gal A) enzyme activity in plasma, isolated leukocytes, or dried blood spots (DBS)</li><li>• Significant clinical manifestations (e.g., neuropathic pain, cardiomyopathy, renal insufficiency, angiokeratomas, cornea verticillata)</li></ul>	

**AND**

**3** - Will not be used in combination with Galafold (migalastat)

Product Name: Fabrazyme, Elfabrio

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Documentation of positive clinical response to therapy

## 2 . Revision History

Date	Notes
7/20/2024	Added new GPI for Elfabrio

Fasenra (benralizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164441
<b>Guideline Name</b>	Fasenra (benralizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Fasenra	
Diagnosis	Severe Asthma
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of documentation (e.g., chart notes) confirming diagnosis of severe asthma	

**AND**

**2** - Submission of documentation (e.g., chart notes, lab values) confirming asthma is an eosinophilic phenotype as defined by a baseline (pre-treatment) peripheral blood eosinophil level greater than or equal to 150 cells per microliter

**AND**

**3** - One of the following:

**3.1** Patient has had at least two or more asthma exacerbations requiring systemic corticosteroids (e.g., prednisone) within the past 12 months

**OR**

**3.2** Prior asthma-related hospitalization within the past 12 months

**AND**

**4** - One of the following:

**4.1** Both of the following:

**4.1.1** Patient is 6 years of age or older but less than 12 years of age

**AND**

**4.1.2** Paid claims or submission of medical records (e.g., chart notes) confirming patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications:

**4.1.2.1** Both of the following:

- Medium-dose inhaled corticosteroid (e.g., greater than 100 – 200 mcg fluticasone propionate equivalent/day)

- Additional asthma controller medication (e.g., leukotriene receptor antagonist [LTRA] [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], long-acting muscarinic antagonist [LAMA] [e.g., tiotropium])

**OR**

**4.1.2.2** One of the following preferred medium-dose combination ICS/LABA product:

- Brand Advair Diskus (fluticasone propionate 100mcg/ salmeterol 50mcg)
- Advair HFA (fluticasone propionate 115mcg/ salmeterol 21mcg)
- Brand Symbicort (budesonide 80mcg/ formoterol 4.5mcg)
- Dulera (mometasone 100mcg/ formoterol 5mcg)

**OR**

**4.2** Both of the following:

**4.2.1** Patient is 12 years of age or older

**AND**

**4.2.2** Paid claims or submission of medical records (e.g., chart notes) confirming patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications:

**4.2.2.1** Both of the following:

- High-dose inhaled corticosteroid (ICS) (e.g., greater than 500 mcg fluticasone propionate equivalent/day)
- Additional asthma controller medication (e.g., leukotriene receptor antagonist [LTRA] [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], long-acting muscarinic antagonist [LAMA] [e.g., tiotropium])

**OR**

**4.2.2.2** One of the following maximally-dosed combination ICS/LABA product:

- Brand Advair (fluticasone propionate 500mcg/ salmeterol 50mcg)
- Advair HFA (fluticasone propionate 230mcg/ salmeterol 21mcg)
- Brand Symbicort (budesonide 160mcg/ formoterol 4.5mcg)

- Dulera (mometasone 200mcg/ formoterol 5mcg)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Allergist/Immunologist

Product Name:Fasenra	
Diagnosis	Severe Asthma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming a positive clinical response to therapy as evidenced by one of the following:</p> <ul style="list-style-type: none"> <li>• A reduction in asthma exacerbations</li> <li>• Improvement in forced expiratory volume in 1 second [FEV1] from baseline</li> </ul> <p><b>AND</b></p> <p><b>2</b> - Paid claims or submission of documentation (e.g., chart notes) confirming patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [LTRA] [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], long-acting muscarinic antagonist [LAMA] [e.g., tiotropium]) unless there is a contraindication or intolerance to these medications</p> <p><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with one of the following:</p>	

- Pulmonologist
- Allergist/Immunologist

Product Name:Fasenra	
Diagnosis	Eosinophilic Granulomatosis with Polyangiitis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records (e.g., chart notes) confirming patient's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Submission of medical records (e.g., chart notes) confirming patient is currently receiving corticosteroid therapy (e.g., prednisolone, prednisone) unless there is a contraindication or intolerance to corticosteroid therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Pulmonologist</li> <li>• Rheumatologist</li> <li>• Allergist/Immunologist</li> </ul>	

Product Name:Fasenra	
Diagnosis	Eosinophilic Granulomatosis with Polyangiitis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy (e.g., increase in remission time)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Pulmonologist</li> <li>• Rheumatologist</li> <li>• Allergist/Immunologist</li> </ul>	

## 2 . Background

<b>Clinical Practice Guidelines</b>			
<p>The Global Initiative for Asthma Global Strategy for Asthma Management and Prevention: Table 1. Low, medium and high daily doses of inhaled corticosteroids in adolescents and adults 12 years and older [4]</p>			
Inhaled corticosteroid	Total Daily ICS Dose (mcg)		
	Low	Medium	High
Beclometasone dipropionate (pMDI, standard particle, HFA)	200-500	> 500-1000	> 1000

Beclometasone dipropionate (DPI or pMDI, extrafine particle*, HFA)	100-200	> 200-400	> 400
Budesonide (DPI, or pMDI, standard particle, HFA)	200-400	> 400-800	> 800
Ciclesonide (pMDI, extrafine particle*, HFA)	80-160	> 160-320	> 320
Fluticasone furoate (DPI)	100		200
Fluticasone propionate (DPI)	100-250	> 250-500	> 500
Fluticasone propionate (pMDI, standard particle, HFA)	100-250	> 250-500	> 500
Mometasone furoate (DPI)	Depends on DPI device – see product information		
Mometasone furoate (pMDI, standard particle, HFA)	200-400		> 400
<p>DPI: dry powder inhaler; HFA: hydrofluoroalkane propellant; ICS: inhaled corticosteroid; N/A: not applicable; pMDI: pressurized metered dose inhaler (non-chlorofluorocarbon formulations); ICS by pMDI should be preferably used with a spacer *See product information.</p> <p><b><i>This is not a table of equivalence</i></b>, but instead, suggested total daily doses for the 'low', 'medium' and 'high' dose ICS options for adults/adolescents, based on available studies and product information. Data on comparative potency are not readily available and therefore this table does NOT imply potency equivalence. Doses may be country -specific depending on local availability, regulatory labelling and clinical guidelines.</p> <p>For new preparations, including generic ICS, the manufacturer's information should be reviewed carefully; products containing the same molecule may not be clinically equivalent.</p>			

**The Global Initiative for Asthma Global Strategy for Asthma Management and Prevention: Table 2. Low, medium and high daily doses of inhaled corticosteroids in children 6 – 11 years [4]**

Inhaled corticosteroid	Total Daily ICS Dose (mcg)		
	Low	Medium	High

Beclometasone dipropionate (pMDI, standard particle, HFA)	100-200	> 200-400	> 400
Beclometasone dipropionate (pMDI, extrafine particle, HFA)	50-100	> 100-200	> 200
Budesonide (DPI, or pMDI, standard particle, HFA)	100-200	> 200-400	> 400
Budesonide (nebulas)	250-500	>500-1000	>1000
Ciclesonide (pMDI, extrafine particle*, HFA)	80	>80-160	>160
Fluticasone furoate (DPI)	50		n.a.
Fluticasone propionate (DPI)	50-100	> 100-200	> 200
Fluticasone propionate (pMDI, standard particle, HFA)	50-100	> 100-200	> 200
Mometasone furoate (pMDI, standard particle, HFA)	100		200
<p>DPI: dry powder inhaler; HFA: hydrofluoroalkane propellant; ICS: inhaled corticosteroid; N/A: not applicable; pMDI: pressurized metered dose inhaler (non-chlorofluorocarbon formulations); ICS by pMDI should be preferably used with a spacer *See product information.</p> <p><b><i>This is not a table of equivalence</i></b>, but instead, suggested total daily doses for the 'low', 'medium' and 'high' dose ICS options for adults/adolescents, based on available studies and product information. Data on comparative potency are not readily available and therefore this table does NOT imply potency equivalence. Doses may be country -specific depending on local availability, regulatory labelling and clinical guidelines.</p> <p>For new preparations, including generic ICS, the manufacturer's information should be reviewed carefully; products containing the same molecule may not be clinically equivalent.</p>			

### 3 . Revision History

Date	Notes

1/30/2025	Asthma initial auth: updated embedded step for ICS/LABA directing pts to preferred agents.
-----------	--

Fecal Microbiota Agents

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-128980
<b>Guideline Name</b>	Fecal Microbiota Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Rebyota	
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of documentation (e.g., chart notes) confirming diagnosis of recurrent clostridioides difficile infection (CDI) as defined by both of the following:</p> <ul style="list-style-type: none"><li>Presence of diarrhea defined as a passage of 3 or more loose bowel movements within a 24-hour period for 2 consecutive days</li><li>A positive stool test for C.difficile toxin or toxigenic C.difficile</li></ul>	

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Patient has a history of one or more recurrent episodes of CDI

**AND**

**4** - Submission of medical records (e.g., chart notes) confirming BOTH of the following:

**4.1** Patient has completed at least 10 consecutive days of one of the following antibiotic therapies between 24 to 72 hours prior to initiating Rebyota\*:

- oral vancomycin
- Dificid (fidaxomicin)

**AND**

**4.2** Previous episode of CDI is under control (e.g., less than 3 unformed/loose [i.e., Bristol Stool Scale type 6-7] stools/day for 2 consecutive days)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Gastroenterologist
- Infectious disease specialist

Notes	*Trial requirements may be verified via paid pharmacy claims or submission of medical records/chart notes
-------	---

Product Name:Vowst	
Approval Length	14 Day(s)
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Submission of documentation (e.g., chart notes) confirming diagnosis of recurrent clostridioides difficile infection (CDI) as defined by both of the following:

- Presence of diarrhea defined as a passage of 3 or more loose bowel movements within a 24-hour period for 2 consecutive days
- A positive stool test for C.difficile toxin or toxigenic C.difficile

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Patient has a history of two or more recurrent episodes of CDI within 12 months

**AND**

**4** - Submission of medical records (e.g., chart notes) confirming ALL of the following:

**4.1** Patient has completed at least 10 consecutive days of one of the following antibiotic therapies 2-4 days prior to initiating Vowst\*:

- oral vancomycin
- Dificid (fidaxomicin)

**AND**

**4.2** Patient has completed the recommended course of magnesium citrate the day before and at least 8 hours prior to initiating Vowst [A]

**AND**

**4.3** Previous episode of CDI is under control (e.g., less than 3 unformed/loose [i.e., Bristol Stool Scale type 6-7] stools/day for 2 consecutive days)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Gastroenterologist
- Infectious disease specialist

Notes

\*Trial requirements may be verified via paid pharmacy claims or submission of medical records/chart notes

## **2 . Endnotes**

- A. Patients are required to take magnesium citrate 24 hours prior to the first dose of Vowst per the prescribing information. There is currently no efficacy data regarding the use of Vowst without magnesium citrate and the thought is that it helps to clear the antibiotics prior to administration of Vowst. [2,3]

## **3 . Revision History**

Date	Notes
7/26/2023	New program

Fentanyl IR

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99519
<b>Guideline Name</b>	Fentanyl IR
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Fentanyl citrate lozenges (generic Actiq)	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records demonstrating use is for the management of breakthrough pain associated with a cancer diagnosis (cancer diagnosis must be documented)  <b>AND</b>	

**2** - Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids (Document drug and date of trial):

- Morphine sulfate at a doses of greater than or equal to 60 milligrams per day
- Fentanyl transdermal patch at a dose of greater than or equal to 25 micrograms per hour
- Oxycodone at a dose of greater than or equal to 30 milligrams per day
- Oral hydromorphone at a dose of greater than or equal to 8 milligrams per day
- Oral oxymorphone at a dose of greater than or equal to 25 milligrams per day
- An alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 milligrams per day)

**AND**

**3** - The patient is currently taking a long-acting opioid around the clock for cancer pain (Document drug)

**AND**

**4** - ONE of the following:

**4.1** The patient is not concurrently receiving an alternative fentanyl transmucosal product

**OR**

**4.2** BOTH of the following:

**4.2.1** The patient is currently receiving an alternative transmucosal fentanyl product

**AND**

**4.2.2** The prescriber is requesting the termination of all current authorizations for alternative transmucosal fentanyl products in order to begin treatment with the requested medication (Only one transmucosal fentanyl product will be approved at a time. If previous authorizations cannot be terminated, the PA request will be denied)

Product Name: Abstral, Brand Actiq, Brand Fentora, generic fentanyl citrate buccal tablet, Lazanda, Subsys

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records demonstrating use is for the management of breakthrough pain associated with a cancer diagnosis (cancer diagnosis must be documented)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient must have at least a one week history of ONE of the following medications to demonstrate tolerance to opioids (Document drug and date of trial):</p> <ul style="list-style-type: none"> <li>• Morphine sulfate at a doses of greater than or equal to 60 milligrams per day</li> <li>• Fentanyl transdermal patch at a dose of greater than or equal to 25 micrograms per hour</li> <li>• Oxycodone at a dose of greater than or equal to 30 milligrams per day</li> <li>• Oral hydromorphone at a dose of greater than or equal to 8 milligrams per day</li> <li>• Oral oxymorphone at a dose of greater than or equal to 25 milligrams per day</li> <li>• An alternative opioid at an equianalgesic dose (e.g., oral methadone greater than or equal to 20 milligrams per day)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - The patient is currently taking a long-acting opioid around the clock for cancer pain (Document drug)</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - ONE of the following:</p> <p>4.1 The patient is not concurrently receiving an alternative fentanyl transmucosal product</p> <p style="text-align: center;"><b>OR</b></p> <p>4.2 BOTH of the following:</p> <p>4.2.1 The patient is currently receiving an alternative transmucosal fentanyl product</p>	

**AND**

**4.2.2** The prescriber is requesting the termination of all current authorizations for alternative transmucosal fentanyl products in order to begin treatment with the requested medication (Only one transmucosal fentanyl product will be approved at a time. If previous authorizations cannot be terminated, the PA request will be denied)

**AND**

**5** - History of failure, contraindication, or intolerance to Fentanyl citrate lozenges (generic Actiq) [Document date of trial]

## **2 . Revision History**

Date	Notes
6/8/2021	Arizona Medicaid 7.1 Implementation

Fexmid (cyclobenzaprine 7.5mg)- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99458
<b>Guideline Name</b>	Fexmid (cyclobenzaprine 7.5mg)- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Brand Fexmid 7.5mg, generic cyclobenzaprine 7.5mg	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of muscle spasm associated with acute, painful musculoskeletal conditions  <b>AND</b>	

2 - Reason or special circumstance the patient cannot use cyclobenzaprine 5 milligram (mg) or 10mg tablet

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Filspari (sparsentan)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-161585
<b>Guideline Name</b>	Filspari (sparsentan)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Filspari	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting diagnosis of primary immunoglobulin A nephropathy (IgAN) as confirmed by a kidney biopsy	

**AND**

**2** - Patient is at risk of rapid disease progression [e.g., proteinuria greater than 0.75 - 1 g/day, or by other criteria such as clinical risk scoring using the International IgAN Prediction Tool]

**AND**

**3** - Used to slow kidney function decline

**AND**

**4** - Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 30 mL/min/1.73 m<sup>2</sup>

**AND**

**5** - Submission of medical records (e.g., chart notes) demonstrating patient has been on a minimum 90-day trial of a maximally tolerated dose of one of the following (paid pharmacy claims may be used to confirm appropriate trial):

- An angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril)
- An angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan)

**AND**

**6** - Medication will not be used in combination with any of the following:

- Angiotensin receptor blockers or angiotensin receptor-neprilysin inhibitor (ARNI) [e.g., Entresto (sacubitril/valsartan)]
- Endothelin receptor antagonists (ERAs) [e.g., Letairis (ambrisentan), Tracleer (bosentan), Opsumit (macitentan)]
- Tekturna (aliskiren)

**AND**

**7** - Prescribed by or in consultation with a nephrologist

**AND**

**8** - Prescriber is certified/enrolled in the Filspari REMS (Risk Evaluation and Mitigation Strategy) Program

Product Name:Filspari	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting a positive clinical response to therapy as demonstrated by a decrease in urine protein-to-creatinine ratio (UPCR) from baseline</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Medication is not taken in combination with any of the following:</p> <ul style="list-style-type: none"><li>• Angiotensin receptor blockers or angiotensin receptor-neprilysin inhibitor (ARNI) [e.g., Entresto (sacubitril/valsartan)]</li><li>• Endothelin receptor antagonists (ERAs) [e.g., Letairis (ambrisentan), Tracleer (bosentan), Opsumit (macitentan)]</li><li>• Tekturna (aliskiren)</li></ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescriber is certified/enrolled in the Filspari REMS (Risk Evaluation and Mitigation Strategy) Program</p>	

## 2 . Revision History

Date	Notes
12/4/2024	Updates to criteria based on full FDA approval and updated indication

Filsuvez (birch triterpenes)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-146017
<b>Guideline Name</b>	Filsuvez (birch triterpenes)
<b>Formulary</b>	<ul style="list-style-type: none"><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Filsuvez	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of one of the following: <ul style="list-style-type: none"><li>• Dystrophic epidermolysis bullosa (DEB)</li></ul>	

- Junctional epidermolysis bullosa (JEB)

**AND**

**2** - Disease is confirmed by one of the following:

**2.1** Genetic testing confirms mutation in one of the following genes:

**2.1.1** For Dystrophic epidermolysis bullosa (DEB), collagen type VII (COL7A1)

**OR**

**2.1.2** For Junctional epidermolysis bullosa (JEB), one of the following:

- ITGA6
- ITGB4
- collagen type XVII (COL17A1)
- LAMA3
- LAMB3
- LAMC2
- ITGA3
- LAMA3A

**OR**

**2.2** Skin biopsy

**AND**

**3** - Patient is 6 months of age or older

**AND**

**4** - Medication is being used for the treatment of wounds

**AND**

**5** - DEB or JEB associated wounds are present for at least 21 days

**AND**

**6** - Patient does not have signs of infection for wound being treated

**AND**

**7** - Patient has no evidence or history of basal or squamous cell carcinoma for wound being treated

**AND**

**8** - Patient does not have history of stem cell transplant

**AND**

**9** - Medication is not being used concurrently with other FDA approved therapies (e.g., Vyjuvek) for the treatment epidermolysis bullosa

**AND**

**10** - Standard wound care management not adequate in healing wounds (e.g., daily wound dressings, pain management, controlling infections)

**AND**

**11** - Prescribed by or in consultation with a specialist with expertise in wound care

Product Name:Filsuvez	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient demonstrates positive clinical response to therapy as evidenced by wound is healing but not completely closed

**AND**

2 - Patient does not have signs of infection for wound being treated

**AND**

3 - Patient has no evidence or history of basal or squamous cell carcinoma for wound being treated

**AND**

4 - Prescribed by or in consultation with a specialist with expertise in wound care

**2 . Revision History**

Date	Notes
4/23/2024	New program

Firdapse

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116131
<b>Guideline Name</b>	Firdapse
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Firdapse	
Diagnosis	Lambert-Eaton myasthenic syndrome (LEMS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has a diagnosis of Lambert-Eaton myasthenic syndrome (LEMS)	

**AND**

**2** - Patient is not receiving Firdapse in combination with similar potassium channel blockers [e.g., Ampyra (dalfampridine), Ruzurgi (amiframpridine)]

**AND**

**3** - Patient is 6 years of age or older

Product Name:Firdapse	
Diagnosis	Lambert-Eaton myasthenic syndrome (LEMS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Firdapse therapy	
<b>AND</b>	
2 - Patient is not receiving Firdapse in combination with similar potassium channel blockers [e.g., Ampyra (dalfampridine), Ruzurgi (amifampridine)]	

## 2 . Revision History

Date	Notes
10/27/2022	Added age requirement

Flucytosine- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99520
<b>Guideline Name</b>	Flucytosine- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Ancobon, generic flucytosine	
Approval Length	2 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Diagnosis of septicemia, endocarditis or a urinary system infection caused by Candida species	

**OR**

**1.2** Diagnosis of meningitis or a pulmonary infection caused by Cryptococcus species

**AND**

**2** - If the patient is being treated for a systemic infection, flucytosine is being used in combination with amphotericin B

Product Name: Brand Ancobon, generic flucytosine*	
Diagnosis	Infectious Diseases Society of America (IDSA) Recommended Regimens
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The medication is being prescribed by or in consultation with an infectious disease specialist.	
Notes	*Approval duration based on provider recommended treatment durations, up to 12 months.

## 2 . Revision History

Date	Notes
5/13/2021	Arizona Medicaid 7.1 Implementation

Forteo, Prolia, Teriparatide, Tymlos - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-230268
<b>Guideline Name</b>	Forteo, Prolia, Teriparatide, Tymlos - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Preferred Drugs: Brand Forteo, Prolia	
Diagnosis	Patients with osteoporosis at high risk for fracture
Approval Length	12 Months**
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of osteoporosis	

**AND**

**2 - ONE of the following:**

**2.1** Bone Mineral Density (BMD) T-score less than or equal to -3.5 based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) [NOTE: Provider must submit patient specific BMD T-score]

**OR**

**2.2** BOTH of the following:

**2.2.1** BMD T-score between -2.5 and -3.5 (BMD T-score greater than -3.5 and less than or equal to -2.5) based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) [NOTE: Provider must submit patient specific BMD T-score]

**AND**

**2.2.2** ONE of the following:

**2.2.2.1** History of ONE of the following resulting from minimal trauma:

- Vertebral compression fracture
- Fracture of the hip
- Fracture of the distal radius
- Fracture of the pelvis
- Fracture of the proximal humerus

**OR**

**2.2.2.2** History of failure, contraindication, or intolerance to ONE conventional osteoporosis therapy [e.g., bisphosphonate or selective estrogen receptor modulator (SERM)] (Document drug, date, and duration of trial)\*\*

**OR**

**2.3** ALL of the following:

**2.3.1** BMD T-score between -1 and -2.5 (BMD T-score greater than -2.5 and less than or equal to -1) based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) [NOTE: Provider must submit patient specific BMD T-score]

**AND**

**2.3.2** ONE of the following:

**2.3.2.1** History of ONE of the following resulting from minimal trauma:

- Vertebral compression fracture
- Fracture of the hip
- Fracture of the distal radius
- Fracture of the pelvis
- Fracture of the proximal humerus

**OR**

**2.3.2.2** ONE of the following Fracture Risk Assessment Tool (FRAX) 10-year fracture probabilities:

- Major osteoporotic fracture at 20 percent or more
- Hip fracture at 3 percent or more

**AND**

**2.3.3** History of failure, contraindication, or intolerance to ONE conventional osteoporosis therapy [e.g., bisphosphonate or selective estrogen receptor modulator (SERM)] (Document drug, date, and duration of trial)\*

**AND**

**3 - For Brand Forteo Requests ONLY:** Treatment duration has not exceeded a total of 24 months\*\* of cumulative use of parathyroid hormone analogs (e.g., Teriparatide Injection, Forteo, Tymlos) during the patient's lifetime (APPLIES TO BRAND FORTEO ONLY)

Notes	*Claims history may be used in conjunction as documentation of drug, date, and duration of trial **Duration of coverage will be limited to 24 months of cumulative parat
-------	---

	thyroid hormone analog therapy (e.g., Forteo, Tymlos) in the patient's lifetime
--	---

Product Name: Non-Preferred Drugs: Brand Teriparatide, generic teriparatide, Tymlos	
Diagnosis	Patients with osteoporosis at high risk for fracture
Approval Length	12 Months**
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of osteoporosis

**AND**

2 - ONE of the following:

**2.1** Bone Mineral Density (BMD) T-score less than or equal to -3.5 based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) [NOTE: Provider must submit patient specific BMD T-score]

**OR**

**2.2** BOTH of the following:

**2.2.1** BMD T-score between -2.5 and -3.5 (BMD T-score greater than -3.5 and less than or equal to -2.5) based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) [NOTE: Provider must submit patient specific BMD T-score]

**AND**

**2.2.2** ONE of the following:

**2.2.2.1** History of ONE of the following resulting from minimal trauma:

- Vertebral compression fracture

- Fracture of the hip
- Fracture of the distal radius
- Fracture of the pelvis
- Fracture of the proximal humerus

**OR**

**2.2.2.2** History of failure, contraindication, or intolerance to ALL of the following (Document drug, date, and duration of trial)

- bisphosphonate (e.g. ALENDRONATE, IBANDRONATE)
- selective estrogen receptor modulator (SERM) (e.g RALOXIFENE)
- Prolia (denosumab)
- Brand Forteo (teriparatide)

**OR**

**2.3** ALL of the following:

**2.3.1** BMD T-score between -1 and -2.5 (BMD T-score greater than -2.5 and less than or equal to -1) based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) [NOTE: Provider must submit patient specific BMD T-score]

**AND**

**2.3.2** ONE of the following:

**2.3.2.1** History of ONE of the following resulting from minimal trauma:

- Vertebral compression fracture
- Fracture of the hip
- Fracture of the distal radius
- Fracture of the pelvis
- Fracture of the proximal humerus

**OR**

**2.3.2.2** ONE of the following Fracture Risk Assessment Tool (FRAX) 10-year fracture probabilities:

- Major osteoporotic fracture at 20 percent or more
- Hip fracture at 3 percent or more

**AND**

**2.3.3** History of failure, contraindication, or intolerance to ALL of the following (Document drug, date, and duration of trial)

- bisphosphonate (e.g. ALENDRONATE, IBANDRONATE)
- selective estrogen receptor modulator (SERM) (e.g RALOXIFENE)
- Prolia (denosumab)
- Brand Forteo (teriparatide)

**AND**

**3** - Treatment duration has not exceeded a total of 24 months\*\* of cumulative use of parathyroid hormone analogs (e.g., Teriparatide Injection, Forteo, Tymlos) during the patient's lifetime

Notes	<p>*Claims history may be used in conjunction as documentation of drug, date, and duration of trial</p> <p>**Duration of coverage will be limited to 24 months of cumulative parathyroid hormone analog therapy (e.g., Forteo, Tymlos) in the patient's lifetime</p>
-------	--

Product Name: Preferred Drugs: Brand Forteo, Prolia; Non-Preferred Drugs: Brand Teriparatide, generic teriparatide, Tymlos	
Diagnosis	Patients with osteoporosis at high risk for fracture
Approval Length	12 Months*
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient demonstrates positive clinical response to therapy</p>	

**AND**

**2** - Treatment duration has not exceeded a total of 24 months\* of cumulative use of parathyroid hormone analogs (e.g., Teriparatide Injection, Forteo, Tymlos) during the patient's lifetime) NOTE: DOES NOT APPLY TO PROLIA

Notes

\*Duration of coverage will be limited to 24 months of cumulative parathyroid hormone analog therapy (e.g., Forteo, Tymlos) in the patient's lifetime

## 2 . Revision History

Date	Notes
3/31/2025	GPI change

Furoscix (furosemide injection)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-224205
<b>Guideline Name</b>	Furoscix (furosemide injection)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Furoscix	
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting diagnosis of one of the following: <ul style="list-style-type: none"><li>chronic heart failure</li><li>chronic kidney disease</li></ul>	

**AND**

**2** - Patient is currently on maintenance oral diuretic therapy (e.g., bumetanide, furosemide, torsemide)

**AND**

**3** - Provider attests that patient will be closely monitored for fluid, electrolyte, and metabolic abnormalities throughout therapy (e.g., hypokalemia, hypovolemia, hyponatremia)

**AND**

**4** - Submission of medical records (e.g., chart notes) or paid claims documenting one of the following:

**4.1** History of failure to intravenous furosemide (Lasix IV)

**OR**

**4.2** Provider has submitted clinical rationale for why intravenous furosemide is inappropriate

## **2 . Revision History**

Date	Notes
3/26/2025	Updated dx criterion due to expanded indication

Galafold

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99613
<b>Guideline Name</b>	Galafold
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Galafold	
Diagnosis	Fabry disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Fabry disease	

**AND**

**2** - Patient has an amenable galactosidase alpha gene (GLA) variant based on in vitro assay data

**AND**

**3** - Patient is not receiving Galafold in combination with Fabrazyme (agalsidase beta)

Product Name:Galafold	
Diagnosis	Fabry disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Galafold therapy	
<b>AND</b>	
2 - Patient is not receiving Galafold in combination with Fabrazyme (agalsidase beta)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona SP to Medicaid Arizona SP for 7/1 eff

Gamifant (emapalumab-lzsg)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135435
<b>Guideline Name</b>	Gamifant (emapalumab-lzsg)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Gamifant	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of primary hemophagocytic lymphohistiocytosis (HLH)	

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims confirming one of the following:

**2.1** Disease is one of the following:

- Refractory
- Recurrent
- Progressive

**OR**

**2.2** Trial and failure, contraindication, or intolerance to conventional HLH therapy (e.g., etoposide, dexamethasone, cyclosporine A, intrathecal methotrexate)

**AND**

**3** - Prescribed by or in consultation with a hematologist/oncologist

**AND**

**4** - Patient has not received hematopoietic stem cell transplantation (HSCT)

Product Name: Gamifant	
Approval Length	6 Months [A]
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) demonstrating a positive clinical response to therapy (e.g., improvement in hemoglobin/lymphocyte/platelet counts, afebrile, normalization of inflammatory factors/markers)	

**AND**

**2** - Patient has not received HSCT

## **2 . Revision History**

Date	Notes
10/27/2023	New program

Gattex (teduglutide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135438
<b>Guideline Name</b>	Gattex (teduglutide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Gattex	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming all of the following: <b>1.1</b> Diagnosis of short bowel syndrome	

**AND**

**1.2** Patient is 1 year of age and older

**AND**

**1.3** Documentation that the patient is dependent on parenteral nutrition/intravenous (PN/IV) support for at least 12 consecutive months

**AND**

**2** - Prescribed by or in consultation with a gastroenterologist

Product Name:Gattex	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting that the patient has had a reduction in weekly parenteral nutrition/intravenous (PN/IV) support from baseline while on Gattex therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a gastroenterologist [C]	

## 2 . Revision History

Date	Notes
------	-------

10/27/2023	New program
------------	-------------

Gaucher's Disease Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-208206
<b>Guideline Name</b>	Gaucher's Disease Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Cerdelga	
Diagnosis	Type 1 Gaucher's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Type 1 Gaucher's disease	

**AND**

**2** - Patient is one of the following as detected by a Food and Drug Administration (FDA)-cleared test:

- CYP2D6 extensive metabolizer
- CYP2D6 intermediate metabolizer
- CYP2D6 poor metabolizer

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to generic miglustat

Product Name:Cerezyme	
Diagnosis	Type 1 Gaucher's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of Type 1 Gaucher's disease that results in one or more of the following conditions:</p> <ul style="list-style-type: none"><li>• Anemia</li><li>• Thrombocytopenia</li><li>• Bone disease</li><li>• Hepatomegaly or splenomegaly</li></ul> <p><b>AND</b></p> <p><b>2</b> - Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to BOTH of the following:</p> <ul style="list-style-type: none"><li>• generic miglustat</li></ul>	

- Eleyso

Product Name:Vpriv, Eleyso	
Diagnosis	Type 1 Gaucher's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of Type 1 Gaucher's disease</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - For VPRIV requests ONLY: Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to BOTH of the following:</p> <ul style="list-style-type: none"> <li>• generic miglustat</li> <li>• Eleyso</li> </ul>	

Product Name:Brand Zavesca, generic miglustat	
Diagnosis	Type 1 Gaucher's disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of mild to moderate Type 1 Gaucher's disease</p>	

**AND**

**2 - For BRAND Zavesca requests ONLY: Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to generic miglustat (applies to Brand Zavesca requests only)**

Product Name:Cerdelga, Cerezyme, Elelyso, Vpriv, Brand Zavesca, generic miglustat	
Diagnosis	Type 1 Gaucher's disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

## **2 . Revision History**

Date	Notes
3/26/2025	P&T Changes: Cerezyme, Cerdelga, and Vpriv to NP. Step through generic miglustat/Elelyso.Removed step through Brand Zavesca.

Generic fluticasone-salmeterol diskus, Wixela Inhub (authorized generic of Advair Diskus)



## Prior Authorization Guideline

<b>Guideline ID</b>	GL-145169
<b>Guideline Name</b>	Generic fluticasone-salmeterol diskus, Wixela Inhub (authorized generic of Advair Diskus)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/24/2024
-----------------	-----------

## 1 . Criteria

Product Name:Generic fluticasone-salmeterol diskus, Wixela Inhub (authorized generic of Advair Diskus)	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Trial and failure, contraindication, or intolerance to ALL of the following preferred agents: <ul style="list-style-type: none"><li>Brand Advair Diskus</li><li>Brand Advair HFA</li><li>Dulera</li></ul>	

- Brand Symbicort

## 2 . Revision History

Date	Notes
4/23/2024	Removed Airduo/generics as targets. Updated criteria to standard t/f verbiage.

Generic tretinoin cream and gel, generic Avita cream and gel, generic atralin gel

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144747
<b>Guideline Name</b>	Generic tretinoin cream and gel, generic Avita cream and gel, generic atralin gel
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/21/2024
-----------------	-----------

## 1 . Criteria

Product Name:generic tretinoin cream and gel, generic Avita cream and gel, generic atralin gel	
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Requests for generic tretinoin cream and gel, generic Avita cream and gel, generic atralin gel should be denied. The plan's preferred product is Brand Retin-A cream or gel.*	
Notes	*Brand Retin-A cream or gel may require PA Note: Clinical Program: Brand Over Generic-Not Covered

## 2 . Revision History

Date	Notes
3/21/2024	Updated guideline to add note that calls out brand is preferred

Global Quantity Limits

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-99460
<b>Guideline Name</b>	Global Quantity Limits
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Quantity Limit, Prescription Limit	
Diagnosis	Quantity limit review (General)
Approval Length	12 month(s)
Guideline Type	Administrative
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 The requested drug must be used for an FDA-approved indication	

**OR**

**1.2** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**AND**

**2** - The drug is being prescribed within the manufacturer's published dosing guidelines or falls within dosing guidelines found in ONE of the following compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**AND**

**3** - The requested dosage cannot be achieved using the plan accepted quantity limit of a different dose or formulation.

**AND**

**4** - The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plans' program.

Product Name:Quantity Limit, Prescription Limit	
Diagnosis	Quantity limit review for the treatment of gender dysphoria*
Approval Length	12 month(s)
Guideline Type	Administrative

**Approval Criteria**

1 - The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**AND**

2 - The drug is being prescribed for an indication that is recognized as a covered benefit by the applicable health plans' program.

Notes	* If the above criteria are not met, then refer for clinical review by an appropriate trained professional (physician or pharmacist) based on the applicable regulatory requirement.
-------	--

Product Name:Quantity Limit, Prescription Limit	
Diagnosis	Monthly prescription limit review for migraine therapy, benzodiazepines, or muscle relaxants
Approval Length	1 month(s)
Guideline Type	Administrative
<b>Approval Criteria</b>	
1 - Medical necessity rationale provided for why the member requires 5 or more fills of the same drug or drug class within a month.	
Notes	*If deemed medically necessary, longer authorization duration is permitted

Product Name:Quantity Limit, Prescription Limit	
Diagnosis	Topical products exceeding the allowable package size per fill OR the allowable quantity per month

Approval Length	12 month(s)
Guideline Type	Administrative
<p><b>Approval Criteria</b></p> <p>1 - The physician attests that a larger quantity is needed for treatment of a larger surface area.</p>	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

GLP-1 Agonists - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242224
<b>Guideline Name</b>	GLP-1 Agonists - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Preferred Drugs: Trulicity, Brand Victoza	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Both of the following:  1.1 Submission of medical records (e.g. chart notes) confirming both of the following: <ul style="list-style-type: none"><li>Diagnosis of type 2 diabetes mellitus</li><li>Baseline A1C greater than or equal to 6.5%</li></ul>	

**AND**

**1.2** Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to metformin at a minimum dose of 1500 milligrams (mg) daily for 90 days, or contraindication or intolerance to metformin

**AND**

**2** - Patient is 10 years of age or older

**AND**

**3** - Drug is not solely being used for weight loss

Notes

If requested medication is being used to treat appetite suppression/weight loss or improve conditions caused by obesity (i.e., Zepbound for OSA) – deny the case for Plan Exclusion. For all other indications deny the case for medical necessity and do not review for off-label use.

Product Name: Non-Preferred Drugs: Brand Exenatide, Brand Liraglutide, Mounjaro, Ozempic

Approval Length | 12 month(s)

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Both of the following:

**1.1** Submission of medical records (e.g. chart notes) confirming both of the following:

- Diagnosis of type 2 diabetes mellitus
- Baseline A1C greater than or equal to 6.5%

**AND**

**1.2** Submission of medical records (e.g., chart notes) or paid claims documenting history of

failure to metformin at a minimum dose of 1500 milligrams (mg) daily for 90 days, or contraindication or intolerance to metformin

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims documenting a history of a 90 day trial resulting in a therapeutic failure, contraindication, or intolerance to ALL of the following:

- Brand Victoza
- Trulicity

**AND**

**3** - One of the following:

- For Brand Liraglutide ONLY: Patient is 10 years of age or older
- For Brand Exenatide, Mounjaro and Ozempic ONLY: Patient is 18 years of age or older

**AND**

**4** - Drug is not solely being used for weight loss

Notes	If requested medication is being used to treat appetite suppression/weight loss or improve conditions caused by obesity (i.e., Zepbound for OSA) – deny the case for Plan Exclusion. For all other indications deny the case for medical necessity and do not review for off-label use.
-------	---

Product Name:Non-Preferred: Rybelsus	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  <b>1</b> - Both of the following:  <b>1.1</b> Submission of medical records (e.g. chart notes) confirming both of the following:	

- Diagnosis of type 2 diabetes mellitus
- Baseline A1C greater than or equal to 6.5%

**AND**

**1.2** Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to metformin at a minimum dose of 1500 milligrams (mg) daily for 90 days, or contraindication or intolerance to metformin

**AND**

**2** - ONE of the following:

**2.1** Submission of medical records (e.g., chart notes) or paid claims documenting a history of a 90 day trial resulting in a therapeutic failure, contraindication, or intolerance to ALL of the following:

- Brand Victoza
- Trulicity

**OR**

**2.2** BOTH of the following:

**2.2.1** Submission of medical records (e.g., chart notes) confirming that the patient is unable to self-inject due to ONE of the following:

- Physical impairment
- Visual impairment
- Lipohypertrophy
- Documented needle-phobia to the degree that the patient has previously refused any injectable therapy or medical procedure (refer to DSM-5 for specific phobia diagnostic criteria)

**AND**

**2.2.2** Submission of medical records (e.g., chart notes) or paid claims documenting a history of a 90 day trial resulting in a therapeutic failure, contraindication, or intolerance to ALL of the following:

- Farxiga
- Jardiance
- Synjardy
- Xigduo XR

**AND**

**3** - Patient is 18 years of age or older

**AND**

**4** - Drug is not solely being used for weight loss

Notes	If requested medication is being used to treat appetite suppression/weight loss or improve conditions caused by obesity (i.e., Zepbound for OSA) – deny the case for Plan Exclusion. For all other indications deny the case for medical necessity and do not review for off-label use.
-------	---

## 2 . Revision History

Date	Notes
4/24/2025	Removed inactive products. Added Exenatide. Updated/aligned verbiage regarding submission of medical records throughout.

Glycopyrrolate Products

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-108674
<b>Guideline Name</b>	Glycopyrrolate Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	7/1/2022
-----------------	----------

### 1 . Criteria

Product Name:Brand Cuvposa oral solution, Dartisla ODT, Brand Robinul, Brand Robinul Forte	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting requested drug is being used for a Food and Drug Administration (FDA)-approved indication</p> <p style="text-align: center;"><b>AND</b></p>	

2 - Trial and failure or intolerance to generic glycopyrrolate tablets or oral solution (verified via pharmacy paid claims or submission of medical records/chart notes)

Product Name: Glycopyrrolate injection 0.6mg/3ml

Approval Length 12 month(s)

Guideline Type Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting requested drug is being used for a Food and Drug Administration (FDA)-approved indication

**AND**

2 - Trial and failure or intolerance to preferred glycopyrrolate injection strengths (e.g., 0.2 mg/ml, 0.4mg/2ml, 1 mg/5ml, 4mg/20ml) (verified via pharmacy paid claims or submission of medical records/chart notes)

**2 . Revision History**

Date	Notes
6/24/2022	Added NP glycopyrrolate inj as target. Changed guideline name to Glycopyrrolate Products.

## Gonadotropin-Releasing Hormone Agonists

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-269209
<b>Guideline Name</b>	Gonadotropin-Releasing Hormone Agonists
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

#### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Lupron Depot Ped, Triptodur, Fensolvi	
Diagnosis	Central Precocious Puberty (CPP)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of central precocious puberty (idiopathic or neurogenic)	

**AND**

**2** - Onset of secondary sexual characteristics in one of the following:

**2.1** Females less than or equal to 8 years of age

**OR**

**2.2** Males less than or equal to 9 years of age

**AND**

**3** - Confirmation of diagnosis as defined by one of the following:

**3.1** Pubertal basal level of luteinizing hormone (based on laboratory reference ranges)

**OR**

**3.2** A pubertal luteinizing hormone response to a gonadotropin releasing hormone (GnRH) stimulation test

**OR**

**3.3** Bone age advanced one year beyond the chronological age

**AND**

**4** - If the request is for Triptodur or Fensolvi, history of failure, contraindication, or intolerance to Lupron-Depot Ped

Product Name:Lupron Depot Ped, Triptodur, Fensolvi	
Diagnosis	Central Precocious Puberty (CPP)
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is currently receiving therapy for central precocious puberty</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is ONE of the following (younger than the appropriate time point for the onset of puberty):</p> <p>3.1 Female younger than 11 years of age</p> <p style="text-align: center;"><b>OR</b></p> <p>3.2 Male younger than 12 years of age</p>	

Product Name:Lupron Depot 3.75 mg and 3-month 11.25 mg	
Diagnosis	Endometriosis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of endometriosis or endometriosis is suspected</p>	

**AND**

**2** - One of the following:

**2.1** History of failure, contraindication, or intolerance to both of the following:

**2.1.1** Oral contraceptives or depot medroxyprogesterone (e.g., Depo- Provera)

**AND**

**2.1.2** Non-steroidal anti-inflammatory drugs (NSAIDs)

**OR**

**2.2** Patient has had surgical ablation to prevent recurrence

Product Name:Lupron Depot 3.75 mg and 3-month 11.25 mg	
Diagnosis	Endometriosis
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of endometriosis or endometriosis is suspected	
<b>AND</b>	
2 - Recurrence of symptoms following an initial course of therapy	
<b>AND</b>	

**3** - Concurrently to be used with add-back therapy (e.g., progestin, estrogen, or bone sparing agents)

Product Name:Lupron Depot 3.75 mg and 3-month 11.25 mg	
Diagnosis	Uterine Leiomyomata (Fibroids)
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 All of the following:</p> <p>1.1.1 For the treatment of uterine leiomyomata-related anemia</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.2 Patient did not respond to iron therapy of 1 month duration</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.3 For use prior to surgery</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 For use prior to surgery to reduce the size of fibroids to facilitate a surgical procedure (e.g., myomectomy, hysterectomy)</p>	

Product Name:Lupron Depot, Lupron Depot-Ped, leuprolide acetate inj kit 5 mg/mL, Triptodur, Fensolvi, Leuprolide acetate (3 month) 22.5 mg inj	
Diagnosis	Gender dysphoria in adolescents
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of gender dysphoria, according to the current Diagnostic and Statistical Manual of Mental Disorders (i.e., DSM-5) criteria, by a mental health professional with expertise in child and adolescent psychiatry</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Medication is prescribed by or in consultation with an endocrinologist or a medical provider experienced in gender dysphoria hormone therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Patient has experienced puberty development to at least Tanner stage 2</p> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - One of the following laboratory tests, based upon the laboratory reference range, confirming:</p> <ul style="list-style-type: none"> <li>• Pubertal levels of estradiol in females</li> <li>• Pubertal levels of testosterone in males</li> <li>• Pubertal basal level of luteinizing hormone (based on laboratory reference ranges)</li> <li>• A pubertal luteinizing hormone response to a gonadotropin-releasing hormone (GnRH) stimulation test</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>5</b> - A letter from the prescriber and/or formal documentation stating all of the following:</p> <p><b>5.1</b> Patient has experienced pubertal changes that have resulted in an increase of their gender dysphoria that has significantly impaired psychological or social functioning</p>	

**AND**

**5.2** Coexisting psychiatric and medical comorbidities or social problems that may interfere with the diagnostic procedures or treatment have been addressed or removed

**AND**

**5.3** Both of the following:

**5.3.1** Current enrollment, attendance, and active participation in psychological and social support treatment program

**AND**

**5.3.2** Patient will continue enrollment, attendance and active participation in psychological and social support throughout the course of treatment

**AND**

**5.4** Patient demonstrates knowledge and understanding of the expected outcomes of treatment and related transgender therapies

**AND**

**6** - If the request is for leuprolide acetate, Triptodur, Fensolvi, history of failure, contraindication, or intolerance to Lupron Depot

Product Name:Lupron Depot, Lupron Depot-Ped, leuprolide acetate inj kit 5 mg/mL, Triptodur, Fensolvi, Leuprolide acetate (3 month) 22.5 mg inj

Diagnosis	Gender dysphoria in adolescents
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - One of the following:

- Documentation (within the last 6 months) of appropriate luteinizing hormone (LH) suppression
- Change in dosing

**AND**

**2** - Documented diagnosis of gender dysphoria, according to the current Diagnostic and Statistical Manual of Mental Disorders (i.e., DSM-5) criteria, by a mental health professional with expertise in child and adolescent psychiatry

**AND**

**3** - Medication is prescribed by or in consultation with an endocrinologist or a medical provider experienced in gender dysphoria hormone therapy

**AND**

**4** - A letter from the prescriber and/or formal documentation stating all of the following:

**4.1** Patient continues to meet their individual goals of therapy for gender dysphoria

**AND**

**4.2** Patient continues to have a strong affinity for the desired (opposite of natal) gender

**AND**

**4.3** Discontinuation of treatment and subsequent pubertal development would interfere with or impair psychological functioning and well-being

**AND**

**4.4** Coexisting psychiatric and medical comorbidities or social problems that may interfere with treatment continue to be addressed or removed

**AND**

**4.5** Both of the following:

**4.5.1** Current enrollment, attendance, and active participation in psychological and social support treatment program

**AND**

**4.5.2** Patient will continue enrollment, attendance and active participation in psychological and social support throughout the course of treatment

**AND**

**4.6** Patient demonstrates knowledge and understanding of the expected outcomes of treatment and related transgender therapies

Product Name:Lupron Depot, Lupron Depot-Ped, leuprolide acetate inj kit 5 mg/mL, Triptodur, Fensolvi, Leuprolide acetate (3 month) 22.5 mg inj

Diagnosis	Adjunct for Gender-Affirming Hormonal Therapy for Transgender Adults
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of gender dysphoria, according to the current Diagnostic and Statistical Manual of Mental Disorders (i.e., DSM-5) criteria, by a mental health professional

**AND**

**2** - Medication is prescribed by or in consultation with an endocrinologist or a medical provider experienced in transgender hormone therapy

**AND**

**3** - Gonads (i.e., testes, ovaries) have not been removed and are functional (e.g., hormone producing)

**AND**

**4** - Patient is currently receiving hormonal therapy (e.g., testosterone, estrogens, progesterones) to achieve the desired (e.g., non-natal) gender

**AND**

**5** - Inability of cross sex hormone therapy to inhibit natal secondary sex characteristics, luteinizing hormone (LH), or gonadotropins (e.g., menses, testosterone)

**AND**

**6** - A letter from the prescriber and/or formal documentation stating all of the following:

**6.1** Transgender patient has identified goals of gender-affirming hormone therapy

**AND**

**6.2** Coexisting psychiatric and medical comorbidities or social problems that may interfere with the diagnostic procedures or treatment have been addressed or removed

**AND**

**6.3** Both of the following:

**6.3.1** Current enrollment, attendance, and active participation in psychological and social support treatment program

**AND**

**6.3.2** Patient will continue enrollment, attendance and active participation in psychological and social support throughout the course of treatment

**AND**

**6.4** Patient demonstrates knowledge and understanding of the expected outcomes of treatment and related transgender therapies

**AND**

**7** - If the request is for leuprolide acetate, Triptodur, Fensolvi, history of failure, contraindication, or intolerance to Lupron Depot

Product Name:Lupron Depot, Lupron Depot-Ped, leuprolide acetate inj kit 5 mg/mL, Triptodur, Fensolvi, Leuprolide acetate (3 month) 22.5 mg inj

Diagnosis	Adjunct for Gender-Affirming Hormonal Therapy for Transgender Adults
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - One of the following:

- Documentation (within the last 6 months) of appropriate luteinizing hormone (LH) suppression
- Change in dosing

**AND**

**2** - Documented diagnosis of gender dysphoria, according to the current Diagnostic and Statistical Manual of Mental Disorders (i.e., DSM-5) criteria, by a mental health professional

**AND**

**3** - Medication is prescribed by or in consultation with an endocrinologist or a medical provider experienced in transgender hormone therapy

**AND**

**4** - Gonads (i.e., testes, ovaries) are intact

**AND**

**5** - Patient is currently receiving hormonal therapy (e.g., testosterone, estrogens, progesterones) to achieve the desired (e.g., non-natal) gender

**AND**

**6** - Inability of cross sex hormone therapy to inhibit natal secondary sex characteristics, luteinizing hormone (LH), or gonadotropins (e.g., menses, testosterone)

**AND**

**7** - A letter from the prescriber and/or formal documentation stating all of the following:

**7.1** Transgender patient continues to meet goals of gender-affirming hormone therapy

**AND**

**7.2** Coexisting psychiatric and medical comorbidities or social problems that may interfere with the diagnostic procedures or treatment continue to be addressed or removed

**AND**

**7.3** Both of the following:

**7.3.1** Current enrollment, attendance, and active participation in psychological and social support treatment program

**AND**

**7.3.2** Patient will continue enrollment, attendance and active participation in psychological and social support throughout the course of treatment

**AND**

**7.4** Patient demonstrates knowledge and understanding of the expected outcomes of treatment and related transgender therapies

Product Name:Lupron Depot, Lupron Depot-Ped, leuprolide acetate inj kit 5 mg/mL, Triptodur, Fensolvi, Leuprolide acetate (3 month) 22.5 mg inj

Diagnosis	Fertility Preservation
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - For use in pre-menopausal women

**AND**

**2** - Patient is receiving a cytotoxic agent that is associated with causing primary ovarian insufficiency (premature ovarian failure) [e.g., Cytoxan (cyclophosphamide), procarbazine, vinblastine, cisplatin]

**AND**

**3** - If the request is for leuprolide acetate, Triptodur, Fensolvi, history of failure, contraindication, or intolerance to Lupron Depot.

Product Name:Lupron Depot, Lupron Depot-Ped, leuprolide acetate inj kit 5 mg/mL, Triptodur, Fensolvi, Leuprolide acetate (3 month) 22.5 mg inj

Diagnosis	Fertility Preservation
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Patient is currently receiving gonadotropin-releasing hormone (GnRH) analog therapy for the purpose of fertility preservation

**AND**

**2** - Patient continues to receive a cytotoxic agent that is associated with causing primary ovarian insufficiency (premature ovarian failure) [e.g., Cytoxan (cyclophosphamide), procarbazine, vinblastine, cisplatin]

Product Name:Lupron Depot 7.5 mg, 22.5 mg, 30 mg and 45 mg, leuprolide acetate inj kit 5 mg/mL, Leuprolide acetate (3 month) 22.5 mg inj, Lutrate Depot

Diagnosis	Advanced or Metastatic Prostate Cancer
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of advanced or metastatic prostate cancer

## 2 . Revision History

Date	Notes
5/30/2025	Removed inactive product Lupanta, removed leuprolide acetate as target from CPP indication

Gralise, Horizant - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-141352
<b>Guideline Name</b>	Gralise, Horizant - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Brand Gralise, generic gabapentin (once-daily)	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of postherpetic neuralgia (PHN)  <b>AND</b>	

**2** - Trial and failure or intolerance to generic immediate-release gabapentin (generic for Neurontin)

**AND**

**3** - For generic gabapentin (once-daily) requests ONLY: Trial and failure or intolerance to Brand Gralise

Product Name:Horizant	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - One of the following:	
1.1 Both of the following:	
<ul style="list-style-type: none"><li>• Diagnosis of postherpetic neuralgia (PHN)</li><li>• Trial and failure or intolerance to generic immediate-release gabapentin (generic for Neurontin)</li></ul>	
<b>OR</b>	
1.2 Diagnosis of restless legs syndrome	

## 2 . Revision History

Date	Notes
2/28/2024	Added new GPIs for generic Gralise with step through Brand Gralise (preferred). Specified trial of preferred generic gabapentin is "immediate-release, generic for Neurontin". Added step through preferred IR gabapentin for Horizant PHN indication.

Growth Hormone, Growth Stimulating Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152804
<b>Guideline Name</b>	Growth Hormone, Growth Stimulating Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:All products	
Diagnosis	Idiopathic Short Stature (ISS)
Approval Length	N/A - Requests for non-approvable diagnoses should not be approved
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Requests for coverage for diagnosis of Idiopathic Short Stature (ISS) are not authorized and will not be approved	
Notes	Approval Length: N/A - Requests for Idiopathic Short Stature (ISS) should not be approved. Deny as a benefit exclusion.

Product Name:Non Preferred: Genotropin Cartridge, Humatrope, Ngenla, Nutropin AQ NuSpin, Omnitrope, Saizen, Saizen Click Easy, Skytrofa, Sogroya, Zomacton	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to ALL preferred products listed below:</p> <ul style="list-style-type: none"> <li>• Genotropin Miniquick</li> <li>• Norditropin</li> </ul>	

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Pediatric Growth Hormone Deficiency (GHD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 ONE of the following:</p> <p>1.1.1 All of the following:</p> <ul style="list-style-type: none"> <li>• Infant is less than 4 months of age</li> <li>• Infant has growth deficiency</li> <li>• Prescribed by an endocrinologist</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>1.1.2 BOTH of the following:</p> <ul style="list-style-type: none"> <li>• History of neonatal hypoglycemia associated with pituitary disease</li> </ul>	

- Prescribed by an endocrinologist

**OR**

**1.1.3 BOTH of the following:**

- Diagnosis of panhypopituitarism
- Prescribed by an endocrinologist

**OR**

**1.2 ALL of the following:**

**1.2.1** Diagnosis of pediatric growth hormone (GH) deficiency as confirmed by ONE of the following:

**1.2.1.1** Projected height (as determined by extrapolating pre-treatment growth trajectory along current channel to 18-20 year mark) is greater than 2.0 standard deviations [SD] below midparental height utilizing age and gender growth charts related to height

**OR**

**1.2.1.2** Height is greater than 2.25 SD below population mean (below the 1.2 percentile for age and gender) utilizing age and gender growth charts related to height

**OR**

**1.2.1.3** Growth velocity is greater than 2 SD below mean for age and gender

**OR**

**1.2.1.4** Delayed skeletal maturation of greater than 2 SD below mean for age and gender (e.g., delayed greater than 2 years compared with chronological age)

**AND**

**1.2.2 ONE of the following:**

**1.2.2.1** BOTH of the following:

- Patient is male
- Bone age less than 16 years

**OR**

**1.2.2.2** BOTH of the following:

- Patient is female
- Bone age less than 14 years

**AND**

**1.2.3** Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following:

**1.2.3.1** BOTH of the following:

**1.2.3.1.1** Patient has undergone TWO of the following provocative GH stimulation tests:

- Arginine
- Clonidine
- Glucagon
- Insulin
- Levodopa
- Growth hormone releasing hormone

**AND**

**1.2.3.1.2** BOTH GH response values are less than 10 micrograms per liter

**OR**

**1.2.3.2** BOTH of the following:

**1.2.3.2.1** Patient is less than 1 year of age

**AND**

**1.2.3.2.2** ONE of the following is below the age and gender adjusted normal range as provided by the physician's lab:

- Insulin-like Growth Factor 1 (IGF-1/Somatomedin-C)
- Insulin Growth Factor Binding Protein-3 (IGFBP-3)

**AND**

**1.2.4** ONE of the following:

**1.2.4.1** Request does not exceed a maximum supply limit of 0.3 milligrams per kilogram per week

**OR**

**1.2.4.2** BOTH of the following:

- Tanner Stage 3 or greater
- Request does not exceed a maximum supply limit of 0.7 milligrams per kilogram per week

**AND**

**1.2.5** Prescribed by an endocrinologist

Notes	*Includes children who have undergone brain radiation. If patient is a Transition Phase Adolescent or Adult who had childhood onset GH deficiency, utilize criteria for Transition Phase Adolescent or Adult GHD efficiency.
-------	--

Product Name: Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Pediatric Growth Hormone Deficiency (GHD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Height increase of at least 2 centimeters per year over the previous year documented by BOTH of the following:\*\*

- Previous height and date obtained
- Current height and date obtained

**AND**

**2** - BOTH of the following:\*\*

- Expected adult height not attained
- Documentation of expected adult height goal (e.g. genetic potential)

**AND**

**3** - Calculated height (growth) velocity over the past 12 months

**AND**

**4** - ONE of the following:

**4.1** BOTH of the following:

- Patient is male
- Bone age less than 16 years

**OR**

**4.2** BOTH of the following:

- Patient is female
- Bone age less than 14 years

**AND**

**5 - ONE** of the following:

**5.1** Request does not exceed a maximum supply limit of 0.3 milligrams per kilogram per week

**OR**

**5.2 BOTH** of the following:

- Tanner Stage 3 or greater
- Request does not exceed a maximum supply limit of 0.7 milligrams per kilogram per week

**AND**

**6 - Prescribed** by an endocrinologist

Notes	*Includes children who have undergone brain radiation. If patient is a Transition Phase Adolescent or Adult who had childhood onset GH deficiency, utilize criteria for Transition Phase Adolescent or Adult GH D deficiency. ** Documentation of previous height, current height and goal expected adult height will be required for renewal.
-------	---

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Prader-Willi Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Prader-Willi Syndrome	

**AND**

**2 - Prescribed by an endocrinologist**

Product Name: Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Prader-Willi Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1 - ONE of the following criteria:**

**1.1 BOTH of the following:**

**1.1.1** Evidence of positive response to therapy (e.g., increase in total lean body mass, decrease in fat mass)

**AND**

**1.1.2** Prescribed by an endocrinologist

**OR**

**1.2 ALL of the following:**

**1.2.1** Height increase of at least 2 centimeters per year over the previous year of treatment as documented by BOTH of the following:

- Previous height and date obtained
- Current height and date obtained

**AND**

**1.2.2 BOTH of the following:**

- Expected adult height not attained
- Documentation of expected adult height goal

**AND**

**1.2.3 Prescribed by an endocrinologist**

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Growth Failure in Children Small for Gestational Age (SGA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 -</b> Diagnosis of small for gestational age (SGA) based on demonstration of catch up growth failure in the first 24 months of life using a 0-36 month growth chart as confirmed by documentation that ONE of the following is below the third percentile for gestational age (more than 2 standard deviations [SD] below population mean):</p> <ul style="list-style-type: none"><li>• Birth weight</li><li>• Birth length</li></ul> <p><b>AND</b></p> <p><b>2 -</b> Documentation that height remains less than or equal to the third percentile (more than 2 SD below population mean)</p> <p><b>AND</b></p> <p><b>3 -</b> Prescribed by an endocrinologist</p>	

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Growth Failure in Children Small for Gestational Age (SGA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Height increase of at least 2 centimeters per year over the previous year documented by BOTH of the following:*</p> <ul style="list-style-type: none"> <li>• Previous height and date obtained</li> <li>• Current height and date obtained</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Documentation of BOTH of the following:*</p> <ul style="list-style-type: none"> <li>• Expected adult height not attained</li> <li>• Expected adult height goal</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by an endocrinologist</p>	
Notes	*Documentation of previous height, current height and goal expected adult height will be required for renewal.

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Turner Syndrome or Noonan Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## Approval Criteria

1 - Diagnosis of pediatric growth failure associated with ONE of the following:

1.1 BOTH of the following:

1.1.1 Turner Syndrome (Gonadal Dysgenesis)

**AND**

1.1.2 BOTH of the following:

- Patient is female
- Bone age less than 14 years

**OR**

1.2 BOTH of the following:

1.2.1 Noonan Syndrome

**AND**

1.2.2 ONE of the following:

1.2.2.1 BOTH of the following:

- Patient is male
- Bone age less than 16 years

**OR**

1.2.2.2 BOTH of the following:

- Patient is female
- Bone age less than 14 years

**AND**

**2** - Height is below the fifth percentile on growth charts for age and gender

**AND**

**3** - Prescribed by an endocrinologist

Product Name:Preferred: Genotropin Miniquick, Norditropin

Diagnosis	Turner Syndrome or Noonan Syndrome
-----------	------------------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Height increase of at least 2 centimeters per year over the previous year documented by BOTH of the following:\*

- Previous height and date obtained
- Current height and date obtained

**AND**

**2** - Documentation of BOTH of the following:\*

- Expected adult height not attained
- Expected adult height goal

**AND**

**3** - Prescribed by an endocrinologist

Notes	*Documentation of previous height, current height and goal expected adult height will be required for renewal.
-------	--

Product Name:Preferred: Genotropin Miniquick, Norditropin

Diagnosis	Short-Stature Homeobox (SHOX) Gene Deficiency
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of pediatric growth failure with short-stature homeobox (SHOX) gene deficiency as confirmed by genetic testing</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - ONE of the following:</p> <p><b>2.1</b> BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Patient is male</li> <li>• Bone age less than 16 years</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>2.2</b> BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Patient is female</li> <li>• Bone age less than 14 years</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by an endocrinologist</p>	

Product Name: Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Short-Stature Homeobox (SHOX) Gene Deficiency
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Height increase of at least 2 centimeters per year over the previous year documented by BOTH of the following:\*

- Previous height and date obtained
- Current height and date obtained

**AND**

2 - Documentation of BOTH of the following:\*

- Expected adult height not attained
- Expected adult height goal

**AND**

3 - Prescribed by an endocrinologist

Notes	*Documentation of previous height, current height and goal expected adult height will be required for renewal.
-------	--

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Growth Failure associated with Chronic Renal Insufficiency
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of pediatric growth failure associated with chronic renal insufficiency	
<b>AND</b>	

**2 - ONE of the following:**

**2.1 BOTH of the following:**

- Patient is male
- Bone age less than 16 years

**OR**

**2.2 BOTH of the following:**

- Patient is female
- Bone age less than 14 years

**AND**

**3 - Prescribed by ONE of the following:**

- Endocrinologist
- Nephrologist

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Growth Failure associated with Chronic Renal Insufficiency
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Height increase of at least 2 centimeters per year over the previous year documented by BOTH of the following:*	
<ul style="list-style-type: none"><li>• Previous height and date obtained</li><li>• Current height and date obtained</li></ul>	

**AND**

**2** - Documentation of BOTH of the following:\*

- Expected adult height not attained
- Expected adult height goal

**AND**

**3** - Prescribed by ONE of the following:

- Endocrinologist
- Nephrologist

Notes	*Documentation of previous height, current height and goal expected adult height will be required for renewal.
-------	--

Product Name:Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Adult Growth Hormone Deficiency
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of adult growth hormone deficiency (GHD) as a result of ONE of the following:	
1.1 Clinical records supporting a diagnosis of childhood-onset GHD	
<b>OR</b>	
1.2 BOTH of the following:	
1.2.1 Adult-onset GHD	

**AND**

**1.2.2** Clinical records documenting that hormone deficiency is a result of hypothalamic-pituitary disease from organic or known causes (e.g., damage from surgery, cranial irradiation, head trauma, or subarachnoid hemorrhage)

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) documenting **ONE** of the following:

**2.1** BOTH of the following:

**2.1.1** Patient has undergone **ONE** of the following GH (growth hormone) stimulation tests to confirm adult GH deficiency:

- Insulin tolerance test (ITT)
- ARG (Arginine) and GHRH (growth hormone releasing hormone)
- Glucagon
- ARG

**AND**

**2.1.2** **ONE** of the following peak GH values:

**2.1.2.1** ITT less than or equal to 5 micrograms per liter

**OR**

**2.1.2.2** GHRH and ARG of **ONE** of the following:

- Less than or equal to 11 micrograms per liter if body mass index [BMI] is less than 25 kilograms per meter squared
- Less than or equal to 8 micrograms per liter if BMI is greater than or equal to 25 and less than 30 kilograms per meter squared
- Less than or equal to 4 micrograms per liter if BMI is greater than or equal to 30 kilograms per meter squared

**OR**

**2.1.2.3** Glucagon less than or equal to 3 micrograms per liter

**OR**

**2.1.2.4** ARG less than or equal to 0.4 micrograms per liter

**OR**

**2.2** BOTH of the following:

**2.2.1** Submission of medical records (e.g., chart notes, laboratory values) documenting deficiency of THREE of the following anterior pituitary hormones:

- Prolactin
- ACTH (adrenocorticotrophic hormone)
- TSH (thyroid stimulating hormone)
- FSH/LH (follicle-stimulating hormone/luteinizing hormone)

**AND**

**2.2.2** Insulin-like Growth Factor 1 (IGF-1)/Somatomedin-C level is below the age and gender adjusted normal range as provided by the physician's lab

**AND**

**3** - ONE of the following:

**3.1** Diagnosis of panhypopituitarism

**OR**

**3.2** Other diagnosis and not used in combination with BOTH of the following:

- Aromatase inhibitors [e.g., Arimidex (anastrozole), Femara (letrozole)]

- Androgens [e.g., Delatestryl (testosterone enanthate), Depo-Testosterone (testosterone cypionate)]

**AND**

**4** - Request does not exceed a maximum supply limit of 0.3 milligrams per kilogram per week

**AND**

**5** - Prescribed by an endocrinologist

Product Name: Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Adult Growth Hormone Deficiency
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Documentation of Insulin-like Growth Factor 1 (IGF-1)/Somatomedin C level within the past 12 months</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - ONE of the following:</p> <p><b>2.1</b> Diagnosis of panhypopituitarism</p> <p style="text-align: center;"><b>OR</b></p> <p><b>2.2</b> Other diagnosis and not used in combination with BOTH of the following:</p> <ul style="list-style-type: none"> <li>• Aromatase inhibitors [e.g., Arimidex (anastrozole), Femara (letrozole)]</li> </ul>	

- Androgens [e.g., Delatestryl (testosterone enanthate), Depo-Testosterone (testosterone cypionate)]

**AND**

**3** - Request does not exceed a maximum supply limit of 0.3 milligrams per kilogram per week

**AND**

**4** - Prescribed by an endocrinologist

Product Name: Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Transition Phase Adolescent Patients
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Request does not exceed a maximum supply limit of 0.3 milligrams per kilogram per week</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Documentation of ONE of the following:</p> <ul style="list-style-type: none"> <li>• Attained expected adult height</li> <li>• Closed epiphyses on bone radiograph</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following:</p> <p><b>3.1</b> BOTH of the following:</p>	

**3.1.1** Documentation of high risk of growth hormone (GH) deficiency due to GH deficiency in childhood from ONE of the following:

**3.1.1.1** Embryopathic/congenital defects

**OR**

**3.1.1.2** Genetic mutations

**OR**

**3.1.1.3** Irreversible structural hypothalamic-pituitary disease

**OR**

**3.1.1.4** Panhypopituitarism

**OR**

**3.1.1.5** Deficiency of THREE of the following anterior pituitary hormones:

- ACTH (adrenocorticotrophic hormone)
- TSH (thyroid stimulating hormone)
- Prolactin
- FSH/LH (follicle-stimulating hormone/luteinizing hormone)

**AND**

**3.1.2** ONE of the following:

**3.1.2.1** Insulin-like Growth Factor 1 (IGF-1)/Somatomedin-C level is below the age and gender adjusted normal range as provided by the physician's lab

**OR**

**3.1.2.2** ALL of the following:

**3.1.2.2.1** Patient does not have a low IGF-1/Somatomedin C level

**AND**

**3.1.2.2.2** Discontinued GH therapy for at least 1 month

**AND**

**3.1.2.2.3** Patient has undergone ONE of the following GH stimulation tests after discontinuation of therapy for at least 1 month:

- Insulin tolerance test (ITT)
- ARG (Arginine) and GHRH (growth hormone releasing hormone)
- ARG
- Glucagon

**AND**

**3.1.2.2.4** ONE of the following peak GH values:

**3.1.2.2.4.1** ITT less than or equal to 5 micrograms per liter

**OR**

**3.1.2.2.4.2** GHRH and ARG of ONE of the following:

- Less than or equal to 11 micrograms per liter if body mass index [BMI] is less than 25 kilograms per meter squared
- Less than or equal to 8 micrograms per liter if BMI is greater than or equal to 25 and less than 30 kilograms per meter squared
- Less than or equal to 4 micrograms per liter if BMI is greater than or equal to 30 kilograms per meter squared

**OR**

**3.1.2.2.4.3** Glucagon less than or equal to 3 micrograms per liter

**OR**

**3.1.2.2.4.4** ARG less than or equal to 0.4 micrograms per liter

**OR**

**3.2** ALL of the following:

**3.2.1** At low risk of severe GH deficiency (e.g., due to isolated and/or idiopathic GH deficiency)

**AND**

**3.2.2** Discontinued GH therapy for at least 1 month

**AND**

**3.2.3** BOTH of the following:

**3.2.3.1** Patient has undergone ONE of the following GH stimulation tests after discontinuation of therapy for at least 1 month:

- ITT
- GHRH and ARG
- ARG
- Glucagon

**AND**

**3.2.3.2** ONE of the following peak GH values:

**3.2.3.2.1** ITT less than or equal to 5 micrograms per liter

**OR**

**3.2.3.2.2** GHRH and ARG of ONE of the following:

- Less than or equal to 11 micrograms per liter if body mass index [BMI] is less than 25 kilograms per meter squared
- Less than or equal to 8 micrograms per liter if BMI is greater than or equal to 25 and less than 30 kilograms per meter squared
- Less than or equal to 4 micrograms per liter if BMI is greater than or equal to 30 kilograms per meter squared

**OR**

**3.2.3.2.3** Glucagon less than or equal to 3 micrograms per liter

**OR**

**3.2.3.2.4** ARG less than or equal to 0.4 micrograms per liter

**AND**

**4** - Prescribed by an endocrinologist

Product Name: Preferred: Genotropin Miniquick, Norditropin	
Diagnosis	Transition Phase Adolescent Patients
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive response to therapy (e.g., increase in total lean body mass, exercise capacity or IGF-1 [Insulin-like Growth Factor 1] and IGFBP-3 [Insulin-like growth factor binding protein 3] levels)	

**AND**

**2** - Request does not exceed a maximum supply limit of 0.3 milligrams per kilogram per week

**AND**

**3** - Prescribed by an endocrinologist

Product Name: Serostim	
Diagnosis	Human Immunodeficiency Virus (HIV)-associated wasting syndrome or cachexia
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of human immunodeficiency virus (HIV)-associated wasting syndrome or cachexia</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Documentation of ONE of the following:</p> <p><b>2.1</b> Unintentional weight loss of greater than 10 percent over the last 12 months</p> <p style="text-align: center;"><b>OR</b></p> <p><b>2.2</b> Unintentional weight loss of greater than 7.5 percent over the last 6 months</p> <p style="text-align: center;"><b>OR</b></p>	

**2.3** Loss of 5 percent body cell mass (BCM) within 6 months

**OR**

**2.4** Body mass index (BMI) less than 20 kilograms per meter squared

**OR**

**2.5** ONE of the following:

**2.5.1** ALL of the following:

- Patient is male
- BCM less than 35 percent of total body weight
- BMI less than 27 kilograms per meter squared

**OR**

**2.5.2** ALL of the following:

- Patient is female
- BCM less than 23 percent of total body weight
- BMI less than 27 kilograms per meter squared

**AND**

**3** - A nutritional evaluation has been completed since onset of wasting first occurred

**AND**

**4** - Patient has not had weight loss as a result of other underlying treatable conditions (e.g., depression, mycobacterium avium complex, chronic infectious diarrhea, or malignancy with the exception of Kaposi's sarcoma limited to skin or mucous membranes)

**AND**

5 - Patient's anti-retroviral therapy has been optimized to decrease the viral load

Product Name: Serostim

Diagnosis	Human Immunodeficiency Virus (HIV)-associated wasting syndrome or cachexia
-----------	--

Approval Length	6 month(s)
-----------------	------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Evidence of positive response to therapy (i.e., greater than or equal to 2 percent increase in body weight and/or body cell mass [BCM])

**AND**

2 - ONE of the following targets or goals has not been achieved:

- Weight
- BCM
- Body Mass Index (BMI)

Product Name: Zorbitive\*

Diagnosis	Short Bowel Syndrome
-----------	----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of Short Bowel Syndrome

**AND**

**2** - Patient is currently receiving specialized nutritional support (e.g., intravenous parenteral nutrition, fluid, and micronutrient supplements)

**AND**

**3** - Patient has not previously received 4 weeks of treatment with Zorbtive\*

Notes

\*Treatment with Zorbtive will not be authorized beyond 4 weeks. Administration for more than 4 weeks has not been adequately studied.

Product Name:Increlex	
Diagnosis	Severe Primary IGF-1 Deficiency / Growth Hormone Gene Deletion
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following criteria:</p> <p>1.1 Documentation of ALL of the following:</p> <p>1.1.1 Diagnosis of severe primary Insulin-like Growth Factor 1 (IGF-1) deficiency</p> <p><b>AND</b></p> <p>1.1.2 Height standard deviation score less than or equal to -3.0</p> <p><b>AND</b></p> <p>1.1.3 Basal IGF-1 standard deviation score less than or equal to -3.0</p> <p><b>AND</b></p>	

**1.1.4** Normal or elevated growth hormone levels

**AND**

**1.1.5** Documentation of open epiphyses on last bone radiograph

**AND**

**1.1.6** The patient will not be treated with concurrent growth hormone therapy

**AND**

**1.1.7** Prescribed by an endocrinologist

**OR**

**1.2** ALL of the following:

**1.2.1** Diagnosis of growth hormone gene deletion and has developed neutralizing antibodies to growth hormone

**AND**

**1.2.2** Documentation of open epiphyses on last bone radiograph

**AND**

**1.2.3** The patient will not be treated with concurrent growth hormone therapy

**AND**

**1.2.4** Prescribed by an endocrinologist

Product Name:Increlex	
Diagnosis	Severe Primary IGF-1 Deficiency / Growth Hormone Gene Deletion
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Height increase of at least 2 centimeters per year over the previous year of treatment as documented by BOTH of the following:*</p> <ul style="list-style-type: none"> <li>• Previous height and date obtained</li> <li>• Current height and date obtained</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Documentation of BOTH of the following:*</p> <ul style="list-style-type: none"> <li>• Expected adult height not obtained</li> <li>• Expected adult height goal</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is not treated with concurrent growth hormone therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by an endocrinologist</p>	
Notes	*Documentation of previous height, current height and goal expected adult height will be required for renewal.

## 2 . Revision History

Date	Notes
------	-------

9/24/2024	Updated criteria due to formulary status change: Only Genotropin Miniquick and Norditropin are Preferred. Genotropin carts to NP, Omnitrope to NP, Zomacton to NP
-----------	---

HCG

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99463
<b>Guideline Name</b>	HCG
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Novarel, Ovidrel, Brand Pregnyl, generic chorionic gonadotropin	
Diagnosis	Prepubertal Cryptorchidism
Approval Length	6 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of prepubertal cryptorchidism not due to anatomical obstruction	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Hemangeol

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99464
<b>Guideline Name</b>	Hemangeol
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Hemangeol	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of proliferating infantile hemangioma  <b>AND</b>	

2 - Prescriber provides a reason or special circumstance the patient cannot use generic propranolol oral solution

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Hemophilia Clotting Factors

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-269194
<b>Guideline Name</b>	Hemophilia Clotting Factors
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	6/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Corifact	
Diagnosis	Congenital Factor XIII Deficiency (i.e., Fibrin Stabilizing Factor Deficiency)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of congenital factor XIII deficiency	

**AND**

**2 - ONE of the following:**

- Routine prophylactic treatment of bleeding
- Peri-operative management of surgical bleeding
- Treatment of bleeding episodes

Product Name:Tretten	
Diagnosis	Congenital Factor XIII Deficiency (i.e., Fibrin Stabilizing Factor Deficiency)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of congenital factor XIII A-subunit deficiency	
<b>AND</b>	
2 - ONE of the following:	
<ul style="list-style-type: none"><li>• Routine prophylactic treatment of bleeding</li><li>• Peri-operative management of surgical bleeding</li><li>• Treatment of bleeding episodes</li></ul>	

Product Name:Humate-P	
Diagnosis	Von Willebrand Disease (VWD)
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - One of the following:

1.1 Diagnosis of severe von Willebrand disease

**OR**

1.2 BOTH of the following:

- Diagnosis of mild or moderate von Willebrand disease
- History of failure, contraindication or intolerance to treatment with desmopressin

**AND**

2 - ONE of the following:

- Treatment of bleeding episodes
- Peri-operative management of surgical bleeding

Product Name:Alphanate	
Diagnosis	Von Willebrand Disease (VWD)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of mild or moderate von Willebrand disease	
<b>AND</b>	
2 - Used for peri-operative management of surgical bleeding	

**AND**

**3** - History of failure, contraindication or intolerance to treatment with desmopressin

Product Name:Wilate or Vonvendi	
Diagnosis	Von Willebrand Disease (VWD)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of von Willebrand disease	
<b>AND</b>	
2 - ONE of the following:	
<ul style="list-style-type: none"><li>• Treatment of bleeding episodes</li><li>• Peri-operative management of surgical bleeding</li><li>• Routine prophylactic treatment</li></ul>	

Product Name:NovoSeven RT	
Diagnosis	Congenital Factor VII Deficiency
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of congenital factor VII deficiency	

**AND**

**2 - ONE of the following:**

- Treatment of bleeding episodes
- Routine prophylactic treatment of bleeding

Product Name: Advate, Alphanate, Humate-P, Hemofilm, KoAte, KoAte-DVI, Kogenate FS, Kovaltry, NovoEight, Nuwiq, Recombinate, Xyntha, or Xyntha Solofuse

Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1 - Diagnosis of hemophilia A**

**AND**

**2 - ONE of the following:**

- Routine prophylactic treatment of bleeding
- Peri-operative management of surgical bleeding
- Treatment of bleeding episodes

Product Name: Eloctate

Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

## **Approval Criteria**

**1** - Diagnosis of hemophilia A

**AND**

**2** - ONE of the following:

- Routine prophylactic treatment of bleeding
- Peri-operative management of surgical bleeding
- Treatment of bleeding episodes

**AND**

**3** - Patient is not a suitable candidate for treatment with shorter half-life Factor VIII (recombinant) products [e.g., Advate, Kogenate FS, Kovaltry, Novoeight, Nuwiq, or Recombinate] as attested by the prescribing physician

**AND**

**4** - ONE of the following:

**4.1** BOTH of the following:

- Dose does not exceed 50 IU/kg
- Infusing no more frequently than every 4 days

**OR**

**4.2** Requested dosage regimen does not exceed 12.5 IU/kg/day

**OR**

**4.3** BOTH of the following:

**4.3.1** Patient is less than 6 years of age

**AND**

**4.3.2 ONE of the following:**

- Pharmacokinetic (PK) testing results suggest that dosing more intensive than 50 IU/kg is required
- PK testing results suggest that dosing more frequently than every 3 to 5 days is required
- PK testing results suggest that dosing more intensive than 14.5 IU/kg/day is required

Product Name: Jivi	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of hemophilia A	
<b>AND</b>	
2 - ONE of the following:	
<ul style="list-style-type: none"><li>• Peri-operative management of surgical bleeding</li><li>• Routine prophylactic treatment of bleeding</li><li>• Treatment of bleeding episodes</li></ul>	
<b>AND</b>	
3 - Patient has previously received Factor VIII replacement therapy	
<b>AND</b>	

4 - Patient is 12 years of age or older

**AND**

5 - Patient is not a candidate for treatment with shorter acting half-life Factor VIII (recombinant) products [e.g., Advate, Kogenate FS, Kovaltry, Novoeight, Nuwiq, or Recombinate] as attested by the prescribing physician

**AND**

6 - Patient is not to receive routine infusions more than 2 times per week

Product Name: Afstyla	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of hemophilia A	
<b>AND</b>	
2 - ONE of the following:	
<ul style="list-style-type: none"><li>• Routine prophylactic treatment of bleeding</li><li>• Peri-operative management of surgical bleeding</li><li>• Treatment of bleeding episodes</li></ul>	
<b>AND</b>	
3 - Patient is not a suitable candidate for treatment with shorter acting half-life Factor VIII (recombinant) products [e.g., Advate, Kogenate FS, Kovaltry, Novoeight, Nuwiq, Recombinate] as attested by the prescribing physician	

**AND**

**4** - ONE of the following:

**4.1** Patient is not to receive routine infusions more frequently than 3 times per week

**OR**

**4.2** BOTH of the following:

- Patient is less than 12 years of age
- Pharmacokinetic (PK) testing results suggest that more frequently than 3 times per week dosing is required

Product Name:Hemlibra	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - One of the following:</p> <p><b>1.1</b> All of the following:</p> <p><b>1.1.1</b> Diagnosis of severe hemophilia A</p> <p><b>AND</b></p> <p><b>1.1.2</b> Documentation of endogenous factor VIII level less than 1% of normal factor VIII ( &lt; 0.01 IU/mL)</p> <p><b>AND</b></p>	

**1.1.3** Physician attestation that the patient is not to receive extended half-life factor VIII replacement products (e.g., Eloctate, Adynovate, Afstyla, Jivi) for the treatment of breakthrough bleeding episodes

**OR**

**1.2** All of the following:

**1.2.1** One of the following:

**1.2.1.1** BOTH of the following:

- Diagnosis of moderate hemophilia A
- Documentation of endogenous factor VIII level greater than or equal to 1% to less than 5% (greater than or equal to 0.01 IU/mL to less than 0.05 IU/mL)

**OR**

**1.2.1.2** Both of the following:

- Diagnosis of mild hemophilia A
- Documentation of endogenous factor VIII level greater than or equal to 5% (greater than 0.05 IU/mL)

**AND**

**1.2.2** Submission of medical records (e.g., chart notes, laboratory values) documenting a failure to meet clinical goals (e.g., continuation of spontaneous bleeds, inability to achieve appropriate trough level, previous history of inhibitors) after a trial of prophylactic factor VIII replacement products

**AND**

**1.2.3** Physician attestation that the patient is not to receive extended half-life factor VIII replacement products (e.g., Eloctate, Adynovate, Afstyla, Jivi) for the treatment of breakthrough bleeding episodes

**OR**

**1.3 BOTH of the following:**

- Diagnosis of hemophilia A
- Patient has developed high-titer factor VIII inhibitors (greater than or equal to 5 Bethesda units [BU])

**AND**

**2 - Prescribed for the prevention of bleeding episodes (i.e., routine prophylaxis)**

Product Name:FEIBA	
Diagnosis	Hemophilia A
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - Diagnosis of hemophilia A</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>2 - Documentation of inhibitors (e.g., Bethesda inhibitor assay)</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>3 - One of the following:</b></p> <ul style="list-style-type: none"><li>• Routine prophylactic treatment of bleeding</li><li>• Peri-operative management of surgical bleeding</li><li>• Treatment of bleeding episodes</li></ul>	

Product Name:NovoSeven RT, Obizur	
Diagnosis	Acquired factor VIII Hemophilia

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of acquired factor VIII hemophilia (e.g., acquired hemophilia A, Factor VIII deficiency)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Treatment or prevention of bleeding episodes</p>	

Product Name:Adynovate	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of hemophilia A</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <ul style="list-style-type: none"> <li>• Routine prophylactic treatment of bleeding</li> <li>• Peri-operative management of surgical bleeding</li> <li>• Treatment of bleeding episodes</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is not a suitable candidate for treatment with shorter acting half-life Factor VIII (recombinant) products [Advate, Kogenate FS, Kovaltry, Novoeight, Nuwiq, or Recombinate] as attested by the prescribing physician</p>	

**AND**

**4** - One of the following:

**4.1** BOTH of the following:

- Patient is not to receive routine infusions more frequently than 2 times per week
- Patient is not to receive a routine dose greater than 50 IU/kg

**OR**

**4.2** ALL of the following:

- Patient is less than 12 years of age
- Patient is not to receive routine infusions more frequently than 2 times per week
- Patient is not to receive a routine dose greater than 70 IU/kg

Product Name:Esperoct	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of hemophilia A	
<b>AND</b>	
2 - ONE of the following:	
<ul style="list-style-type: none"><li>• Routine prophylactic treatment of bleeding</li><li>• Peri-operative management of surgical bleeding</li><li>• Treatment of bleeding episodes</li></ul>	

**AND**

**3** - ONE of the following:

**3.1** Patient is not to receive routine infusions more frequently than 2 times per week

**OR**

**3.2** BOTH of the following:

- Patient is less than 12 years of age
- Pharmacokinetic (PK) testing results suggest that more frequent than 2 times per week dosing is required

Product Name:Wilate	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of hemophilia A	
<b>AND</b>	
2 - ONE of the following:	
2.1 Routine prophylactic treatment of bleeding	
<b>OR</b>	
2.2 Treatment of bleeding episodes	

Product Name:NovoSeven RT, Sevenfact	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of hemophilia A</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Documentation of inhibitors (e.g., Bethesda inhibitor assay)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - One of the following:</p> <ul style="list-style-type: none"> <li>• Peri-operative management of surgical bleeding</li> <li>• Treatment of bleeding episodes</li> </ul>	

Product Name:Altuviiiio	
Diagnosis	Hemophilia A (i.e., Factor VIII Deficiency, Classical Hemophilia)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of hemophilia A</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p>	

- Treatment of bleeding episodes
- Prevention of bleeding in surgical interventions or invasive procedures (e.g., surgical prophylaxis)
- Prevention of bleeding episodes (i.e., routine prophylaxis)

**AND**

**3** - Patient is not a suitable candidate for treatment with shorter acting half-life Factor VIII (recombinant) products [e.g., Advate, Kogenate FS, Kovaltry, Novoeight, Nuwiq, or Recombinate] as attested by the prescribing physician

**AND**

**4** - Both of the following:

- Dose does not exceed 50 IU/kg
- Patient is infusing no more frequently than every 7 days

Product Name: AlphaNine SD, Mononine, Profilnine, Profilnine SD

Diagnosis	Hemophilia B (i.e., Congenital Factor IX Deficiency, Christmas Disease)
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of hemophilia B

**AND**

**2** - One of the following:

- Routine prophylactic treatment
- Treatment of bleeding episodes

Product Name: BeneFIX, Rixubis, Alprolix, Idelvion, Ixinity, or Rebinyn	
Diagnosis	Hemophilia B (i.e., Congenital Factor IX Deficiency, Christmas Disease)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of hemophilia B</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Routine prophylactic treatment</li> <li>• Peri-operative management of surgical bleeding</li> <li>• Treatment of bleeding episodes</li> </ul>	

Product Name: FEIBA	
Diagnosis	Hemophilia B (i.e., Congenital Factor IX Deficiency, Christmas Disease)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of hemophilia B</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Documentation of inhibitors (e.g., Bethesda inhibitor assay)</p>	

**AND**

**3 - ONE of the following:**

- Routine prophylactic treatment of bleeding
- Peri-operative management of surgical bleeding
- Treatment of bleeding episodes

Product Name:NovoSeven RT, Sevenfact

Diagnosis	Hemophilia B (i.e., Congenital Factor IX Deficiency, Christmas Disease)
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1 - Diagnosis of hemophilia B**

**AND**

**2 - Documentation of inhibitors (e.g., Bethesda inhibitor assay)**

**AND**

**3 - ONE of the following:**

- Peri-operative management of surgical bleeding
- Treatment of bleeding episodes

Product Name:Fibryga, RiaSTAP

Diagnosis	Fibrinogen Deficiency (i.e., Factor I deficiency)
-----------	---

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of congenital fibrinogen deficiency, including afibrinogenemia and hypofibrinogenemia</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Treatment of bleeding episodes</p>	

Product Name:NovoSeven RT	
Diagnosis	Glanzmann Thrombasthenia
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of Glanzmann's thrombasthenia</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Refractory to platelet transfusions</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Treatment of bleeding episodes</li> <li>• Peri-operative management of surgical bleeding</li> </ul>	

Product Name:Coagadex	
Diagnosis	Congenital Factor X Deficiency
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of congenital Factor X deficiency</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Treatment of bleeding episodes</li> <li>• Peri-operative management of surgical bleeding</li> <li>• Routine prophylactic treatment</li> </ul>	

Product Name:Hypavzi	
Diagnosis	Hemophilia A, Hemophilia B
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of one of the following:</p> <ul style="list-style-type: none"> <li>• hemophilia A (congenital factor VIII deficiency) without factor VIII inhibitors</li> <li>• hemophilia B (congenital factor IX deficiency) without factor IX inhibitors</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is 12 years of age or older</p>	

**AND**

**3** - Drug will be used for prophylaxis to prevent or reduce the frequency of bleeding episodes

Product Name:Alhemo	
Diagnosis	Hemophilia A, Hemophilia B
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of one of the following:	
<ul style="list-style-type: none"><li>• hemophilia A (congenital factor VIII deficiency) with factor VIII inhibitors</li><li>• hemophilia B (congenital factor IX deficiency) with factor IX inhibitors</li></ul>	
<b>AND</b>	
2 - Patient is 12 years of age or older	
<b>AND</b>	
3 - Drug will be used for prophylaxis to prevent or reduce the frequency of bleeding episodes	

Product Name:Qfitlia	
Diagnosis	Hemophilia A, Hemophilia B
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of one of the following:

- hemophilia A (congenital factor VIII deficiency) with or without factor VIII inhibitors
- hemophilia B (congenital factor IX deficiency) with or without factor IX inhibitors

**AND**

2 - Patient is 12 years of age or older

**AND**

3 - Drug will be used for prophylaxis to prevent or reduce the frequency of bleeding episodes

## 2 . Revision History

Date	Notes
5/29/2025	Added Sevenfact and Qfitlia as targets.

Hepatitis C - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157625
<b>Guideline Name</b>	Hepatitis C - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

### Note:

\*\*Preferred drugs Mavyret and sofosbuvir-velpatasvir (authorized generic of Epclusa) will be approved without requiring prior authorization ONE time per lifetime. Requests for retreatment or non-preferred drugs will require PA\*\*

## 1 . Criteria

Product Name:Preferred: sofosbuvir-velpatasvir (authorized generic of Epclusa)** , Mavyret**	
Diagnosis	Hepatitis C Retreatment
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic Hepatitis C infection status which has been confirmed by detectable	

serum hepatitis C virus (HCV) RNA by quantitative assay completed within the past 90 days from the date of the prior authorization request

**AND**

**2** - Age of the patient is Food and Drug Administration (FDA) approved for the specific HCV DAA (Direct Acting Antiviral) product

**AND**

**3** - The prescribing provider assesses the patient's ability to adhere to the HCV DAA treatment plan and attests the assessment has been documented within the clinical record. For patients that would benefit from adherence aids, the treating provider shall refer the patient to a treatment adherence program

**AND**

**4** - Patient agrees to adhere to the proposed course of treatment, including taking medications as prescribed, attending follow-up appointments, and, if applicable, participating in a treatment adherence program

**AND**

**5** - One of the following:

**5.1** Patient has been screened for Hepatitis A and B and has received one Hepatitis A and one Hepatitis B vaccine prior to requesting treatment

**OR**

**5.2** Patient demonstrates laboratory evidence of immunity to Hepatitis A and B

**AND**

**6** - The Prescriber must submit the following information with the request for HCV DAA medications to be considered:

**6.1** HCV treatment history and responses to treatment

**AND**

**6.2** Current medication list

**AND**

**6.3** Laboratory results for all of the following:

- HCV screen test results
- Genotype and current baseline HCV viral load
- Total bilirubin
- Albumin level
- International Normalized Ratio (INR)
- Creatinine Clearance (CrCl) or Glomerular Filtration Rate (GFR)
- Liver Function Tests (LFTs)
- Complete Blood Count (CBC)
- Viral resistance status (when applicable)
- Hepatic status (Child Pugh Score)

**AND**

**7** - If the HCV DAA product is being used in combination with ribavirin, the prescribing provider attests to monitoring hemoglobin levels periodically

**AND**

**8** - The prescribing provider attests to monitoring HCV RNA levels obtained at 12- and 24-weeks post therapy completion to demonstrate the Sustained Virologic Response (SVR)

**AND**

**9** - DAA HCV treatment coverage is NOT provided for ANY of the following:

**9.1** DAA dosages greater than the FDA approved maximum dosage

**OR**

**9.2** Patients currently using a potent P-gp inducer drug (St. John's wart, rifampin, carbamazepine, ritonavir, tipranavir, etc.)

**OR**

**9.3** Lost or stolen medication absent of good cause

**OR**

**9.4** Fraud, waste, or misuse of HCV DAA medications

Notes

\*Approval length: Mavyret = 8-16 Week(s), sofosbuvir-velpatasvir (authorized generic of Epclusa) = 12-24 Weeks(s). \*\*Preferred drugs Mavyret and sofosbuvir-velpatasvir (authorized generic of Epclusa) will be approved without requiring prior authorization ONE time per lifetime. Requests for retreatment or non-preferred drugs will require PA. Refer to AASLD for specific approval durations AASLD: <https://www.hcvguidelines.org/contents>

Product Name: Non-Preferred: Brand Epclusa, Brand Harvoni, ledipasvir-sofosbuvir (authorized generic of Harvoni), Sovaldi, Zepatier

Diagnosis Hepatitis C

Guideline Type Prior Authorization

**Approval Criteria**

1 - One of the following:

**1.1** Patient was adherent to previous DAA therapy as evidenced by submission of medical records and/or pharmacy prescription claims

**OR**

**1.2** If prior therapy was discontinued due to adverse effects from the DAA, the medical record shall be provided which documents these adverse effects and recommendation of discontinuation by treatment provider

**AND**

**2** - The patient's ability to adhere to the planned course of retreatment has been assessed by the treating provider and documented within the clinical record

**AND**

**3** - Resistance-associated polymorphism testing, when applicable, has been completed and submitted with the prior authorization request when BOTH of the following are true

- Required for regimens whereby the FDA (Food and Drug Administration) requires such testing prior to treatment to ensure clinical appropriateness
- Deemed medically necessary by the clinical reviewer prior to approval of the requested regimen

**AND**

**4** - HCV retreatment with a DAA shall NOT be approved for ANY of the following:

**4.1** Is considered an experimental service

**OR**

**4.2** Monotherapy of Sofosbuvir (Sovaldi)

**OR**

**4.3** DAA dosages greater than the FDA approved maximum dosage

**OR**

**4.4** Grazoprevir/elbasvir (Zepatier) if the NS5A polymorphism testing has not been completed and submitted with the prior authorization request

**OR**

**4.5** Patients currently using a potent P-gp inducer drug (St. John's wart, rifampin, carbamazepine, ritonavir, tipranavir, etc.)

**OR**

**4.6** Lost or stolen medication absent of good cause

**OR**

**4.7** Fraudulent use of HCV DAA medications

**AND**

**5** - If the request is for brand Epclusa or brand Harvoni BOTH of the following:

**5.1** The patient has a therapeutic failure, contraindication, or intolerance to the generic as evidenced by submission of medical records or claims history

**AND**

**5.2** The prescriber must submit the FDA MedWatch form

Notes	*NOTE: The approval length should be as recommended per AASLD. Refer to AASLD for specific approval durations. AASLD: <a href="https://www.hcvguidelines.org/contents">https://www.hcvguidelines.org/contents</a>
-------	---

Product Name: Non-Preferred: Brand Harvoni, ledipasvir-sofosbuvir (authorized generic of Harvoni)

Diagnosis Hepatitis C Retreatment

Approval Length 24 Week(s)

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of chronic hepatitis C infection</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has decompensated cirrhosis (e.g., Child-Pugh Class B or C)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - One of the following:</p> <p>3.1 Patient is ribavirin ineligible</p> <p style="text-align: center;"><b>OR</b></p> <p>3.2 Both of the following:</p> <ul style="list-style-type: none"> <li>• Prior failure (defined as viral relapse, breakthrough while on therapy, or non-responder therapy) to Sovaldi or NS5A-based therapy</li> <li>• Used in combination with ribavirin</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Not used in combination with another HCV direct acting antiviral agent (e.g., Sovaldi [sofosbuvir])</p>	

Product Name:Non-Preferred: Vosevi, Viekira Pak	
Diagnosis	Hepatitis C
Approval Length	12 Week(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of chronic hepatitis C infection

**AND**

2 - One of the following:

2.1 Patient is a previous relapser to an NS5A-based regimen (e.g., Daklinza [daclatasvir]; Eplclusa [sofosbuvir/velpatasvir]; Harvoni [ledipasvir/sofosbuvir]; Mavyret [glecaprevir/pibrentasvir]; Technivie [ombitasvir/paritaprevir/ritonavir]; Viekira [ombitasvir/paritaprevir/ritonavir & dasabuvir]; Zepatier [elbasvir/grazoprevir])

**OR**

2.2 Patient is a previous relapser to a sofosbuvir-based regimen without an NS5A inhibitor

**AND**

3 - Patient is without decompensated liver disease (e.g., Child-Pugh Class B or C)

**AND**

4 - Not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni (ledipasvir/sofosbuvir), Zepatier (elbasvir/grazoprevir)]

Product Name: Non-Preferred: Vosevi, Viekira Pak

Diagnosis	Hepatitis C: Prior Failure to Vosevi/Viekira Pak
-----------	--

Approval Length	24 Week(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of chronic hepatitis C infection

**AND**

2 - Both of the following:

2.1 Patient had a prior treatment failure with Vosevi or Viekira

**AND**

2.2 Used in combination with ribavirin

**AND**

3 - Patient is without decompensated liver disease (e.g., Child-Pugh Class B or C)

**AND**

4 - Not used in combination with another HCV direct acting antiviral agent [e.g., Harvoni (ledipasvir/sofosbuvir), Zepatier (elbasvir/grazoprevir)]

Product Name:Pegasys, PegIntron	
Diagnosis	Hepatitis C
Approval Length	48 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic hepatitis C infection	
<b>AND</b>	

**2** - Patient without decompensated liver disease (defined as Child-Pugh Class B or C)

**AND**

**3** - Will be used as part of a combination antiviral treatment regimen

Product Name: Ribavirin tablets and capsules

Diagnosis	Hepatitis C
-----------	-------------

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of chronic hepatitis C infection

**AND**

**2** - Used in combination with a direct-acting agent

## 2 . Revision History

Date	Notes
10/25/2024	Updated approval durations for Preferred drugs: Mavyret and sofosbuvir-velpatasvir.

Hereditary Angioedema (HAE) Agents - AZM

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-161706
<b>Guideline Name</b>	Hereditary Angioedema (HAE) Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	1/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Cinryze, Haegarda, Orladeyo or Takhzyro	
Diagnosis	Prophylaxis of HAE attacks
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting diagnosis of hereditary angioedema (HAE) confirmed by ONE of the following:</p> <p>1.1 C1 inhibitor (C1-INH) deficiency or dysfunction (Type I or II HAE) as documented by ONE of the following (per laboratory standard):</p>	

- C1-INH antigenic level below the lower limit of normal
- C1-INH functional level below the lower limit of normal

**OR**

**1.2** HAE with normal C1 inhibitor levels and ONE of the following:

- Confirmed presence of a FXII, angiotensin-1 or plasminogen gene mutation
- Recurring angioedema attacks that are refractory to high-dose antihistamines with confirmed family history of angioedema

**AND**

**2** - For prophylaxis against HAE attacks

**AND**

**3** - One of the following:

- Patient is 2 years of age or older (Applies to Takhzyro only)
- Patient is 6 years of age or older (applies to Cinryze and Haegarda only)
- Patient is 12 years of age or older (Applies to Orladeyo only)

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Immunologist
- Allergist

**AND**

**5** - ONE of the following: (APPLIES TO HAEGARDA, ORLADEYO, AND TAKHZYRO ONLY):

**5.1** Submission of medical records documenting a history of failure, contraindication, or intolerance to Cinryze

**OR**

**5.2** Submission of medical records documenting patient is currently on Haegarda, Orladeyo, or Takhzyro therapy

Notes

\*Please note: Preferred agent is Cinryze

Product Name: Berinert, Cinryze [off-label], Brand Firazyr, Generic icatibant, Kalbitor, Ruconest, or Sajazir

Diagnosis

Treatment of acute HAE attacks

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting diagnosis of hereditary angioedema (HAE) confirmed by **ONE** of the following:

**1.1** C1 inhibitor (C1-INH) deficiency or dysfunction (Type I or II HAE) as documented by **ONE** of the following (per laboratory standard):

- C1-INH antigenic level below the lower limit of normal
- C1-INH functional level below the lower limit of normal

**OR**

**1.2** HAE with normal C1 inhibitor levels and **ONE** of the following:

- Confirmed presence of a FXII, angiotensin-1 or plasminogen gene mutation
- Recurring angioedema attacks that are refractory to high-dose antihistamines with confirmed family history of angioedema

**AND**

**2** - For the treatment of acute HAE attacks

**AND**

**3** - Not used in combination with other approved treatments for acute HAE attacks

**AND**

**4** - One of the following:

- Patient is 6 years of age or older (applies to Cinryze only)
- Patient is 12 years of age or older (applies to Kalbitor)
- Patient is 18 years of age or older (applies to Brand Firazyr, generic icatibant, and Sajazir only)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Immunologist
- Allergist

**AND**

**6** - ONE of the following (APPLIES TO BRAND FIRAZYR, RUCONEST, AND SAJAZIR ONLY):

**6.1** Submission of medical records documenting a history of failure, contraindication, or intolerance to ALL of the following preferred HAE agents:

- Berinert
- Cinryze
- generic icatibant
- Kalbitor

**OR**

**6.2** Submission of medical records or paid claims documenting patient is currently on Brand Firazyr, Ruconest, or Sajazir therapy

Notes	Please note: Preferred HAE agents are Berinert, Cinryze, generic icatibant, and Kalbitor
-------	--

## 2 . Revision History

Date	Notes
12/6/2024	Haegarda to NP

Hetlioz, Hetlioz LQ (tasimelteon)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-149318
<b>Guideline Name</b>	Hetlioz, Hetlioz LQ (tasimelteon)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/2/2024
-----------------	----------

## 1 . Criteria

Product Name:Brand Hetlioz capsule, generic tasimelteon capsule	
Diagnosis	Non-24-Hour Sleep-Wake Disorder (Non-24)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting diagnosis of non-24-hour sleep-wake disorder (also known as free-running disorder, free-running or non-entrained type circadian rhythm sleep disorder, or hypernycthemeral syndrome) confirmed by meeting ONE of the following conditions:	

**1.1** Assessment of at least one physiologic circadian phase marker [e.g., measurement of urinary melatonin levels, dim light melatonin onset (as measured in blood or saliva), assessment of core body temperature]

**OR**

**1.2** If assessment of at least one physiologic circadian phase marker cannot be done, the diagnosis must be confirmed by actigraphy performed for at least 1 week plus evaluation of sleep logs recorded for at least 1 month

**AND**

**2** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting patient is totally blind (has no light perception) \*Requests for patients who are sighted (non-blinded) will be reviewed on a case-by-case basis

**AND**

**3** - Patient is 18 years of age or older

**AND**

**4** - Patient has received at least 3 months of continuous therapy (i.e., 3 consecutive months of daily treatment) under the guidance of a physician who specializes in the treatment of sleep disorders of BOTH of the following:

- Melatonin
- Rozerem (ramelteon)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Specialist in sleep disorders
- Neurologist

**AND**

**6** - For Brand Hetlioz capsule requests ONLY: History of failure, intolerance, or contraindication to generic tasimelteon

Product Name: Brand Hetlioz capsule, generic tasimelteon capsule

Diagnosis	Non-24-Hour Sleep-Wake Disorder (Non-24)
-----------	--

Approval Length	6 month(s)
-----------------	------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (e.g., entrainment, clinically meaningful or significant increases in nighttime sleep, clinically meaningful or significant decreases in daytime sleep)

**AND**

**2** - Submission of patient's sleep log demonstrating positive clinical response to therapy

**AND**

**3** - For Brand Hetlioz capsule requests ONLY: History of failure, intolerance, or contraindication to generic tasimelteon

Product Name: Brand Hetlioz capsule, generic tasimelteon capsule

Diagnosis	Smith-Magenis Syndrome (SMS)
-----------	------------------------------

Approval Length	6 month(s)
-----------------	------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

## **Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of Smith-Magenis Syndrome (SMS)

**AND**

**2** - Submission of test results confirming patient has microdeletion of the chromosome band 17p11.2 by fluorescent in situ hybridization (FISH) analysis

**AND**

**3** - Patient is 16 years of age or older

**AND**

**4** - Patient is experiencing nighttime sleep disturbances (i.e., difficulty falling asleep, frequent nighttime waking and early waking)

**AND**

**5** - Patient has received at least 3 months of continuous therapy (i.e., 3 consecutive months of daily treatment) under the guidance of a physician who specializes in the treatment of sleep disorders of BOTH of the following

- Melatonin
- Rozerem (ramelteon) (unless contraindicated due to patient age)

**AND**

**6** - Prescribed by or in consultation with one of the following:

- Specialist in sleep disorders
- Neurologist

**AND**

7 - For Brand Hetlioz capsule requests ONLY: History of failure, intolerance, or contraindication to generic tasimelteon

Product Name:Hetlioz LQ suspension	
Diagnosis	Smith-Magenis Syndrome (SMS)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of Smith-Magenis Syndrome (SMS)

**AND**

2 - Submission of test results confirming patient has microdeletion of the chromosome band 17p11.2 by fluorescent in situ hybridization (FISH) analysis

**AND**

3 - Patient is 3 through 15 years of age

**AND**

4 - Patient is experiencing nighttime sleep disturbances (i.e., difficulty falling asleep, frequent nighttime waking and early waking)

**AND**

5 - Patient has received at least 3 months of continuous therapy (i.e., 3 consecutive months of

daily treatment) of melatonin under the guidance of a physician who specializes in the treatment of sleep disorders

**AND**

**6** - Prescribed by or in consultation with one of the following:

- Specialist in sleep disorders
- Neurologist

Product Name: Brand Hetlioz capsule, generic tasimelteon capsule, Hetlioz LQ suspension

Diagnosis	Smith-Magenis Syndrome (SMS)
-----------	------------------------------

Approval Length	6 month(s)
-----------------	------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (i.e., improvement in nighttime total sleep time, improvement in nighttime sleep quality)

**AND**

**2** - Submission of patient's sleep log demonstrating positive clinical response to therapy

**AND**

**3** - For Brand Hetlioz capsule requests ONLY: History of failure, intolerance, or contraindication to generic tasimelteon

**2 . Revision History**

Date	Notes
7/2/2024	Updated reauth section for SMS indication to include Hetlioz LQ in product name

High Cost Oral Atypical Antipsychotics (Caplyta, Cobenfy, Fanapt, Lybalvi, Rexulti, Vraylar)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-161584
<b>Guideline Name</b>	High Cost Oral Atypical Antipsychotics (Caplyta, Cobenfy, Fanapt, Lybalvi, Rexulti, Vraylar)
<b>Formulary</b>	<ul style="list-style-type: none"> <li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li> </ul>

### Guideline Note:

Effective Date:	1/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Caplyta, Fanapt, Lybalvi, Rexulti, Vraylar	
Diagnosis	Schizophrenia
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of schizophrenia</p>	

**AND**

**2** - One of the following:

**2.1** Inadequate response (minimum 30 day trial) or adverse reaction to TWO, or contraindication to ALL of the following:

- Aripiprazole
- Clozapine
- Olanzapine
- Quetiapine
- Risperidone
- Ziprasidone

**OR**

**2.2** One of the following:

**2.2.1** The patient has been receiving treatment with the requested medication, and is new to the plan (enrollment effective date within the past 90 days)

**OR**

**2.2.2** The patient is currently receiving treatment with the requested medication in the hospital and must continue upon discharge

**AND**

**3** - For Lybalvi requests ONLY, One of the following:

- Patient has a BMI of 30 kg/m<sup>2</sup> or greater
- Patient has a BMI of 27 kg/m<sup>2</sup> or greater with a weight-related comorbidity (e.g., dyslipidemia, hypertension, type 2 diabetes, sleep apnea)
- Patient has a documented history of weight gain of greater than or equal to 10% of their baseline weight after initiating antipsychotic medication
- Physician has provided clinical rationale for requiring samidorphan in addition to olanzapine therapy

Product Name:Cobenfy	
Diagnosis	Schizophrenia
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of schizophrenia</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p>2.1 BOTH of the following:</p> <p>2.1.1 Inadequate response (minimum 30 day trial), adverse reaction, or contraindication to aripiprazole</p> <p style="text-align: center;"><b>AND</b></p> <p>2.1.2 Inadequate response (minimum 30 day trial) or adverse reaction to ONE, or contraindication to ALL of the following:</p> <ul style="list-style-type: none"> <li>• Clozapine</li> <li>• Olanzapine</li> <li>• Quetiapine</li> <li>• Risperidone</li> <li>• Ziprasidone</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>2.2 One of the following:</p> <p>2.2.1 The patient has been receiving treatment with the requested medication, and is new to the plan (enrollment effective date within the past 90 days)</p> <p style="text-align: center;"><b>OR</b></p>	

**2.2.2** The patient is currently receiving treatment with the requested medication in the hospital and must continue upon discharge

Product Name: Fanapt, Lybalvi, Vraylar

Diagnosis | Bipolar I Disorder

Approval Length | 12 month(s)

Guideline Type | Prior Authorization

**Approval Criteria**

1 - Diagnosis of bipolar I disorder

**AND**

2 - ONE of the following:

**2.1** Inadequate response (minimum 30 day trial) or adverse reaction to THREE, or contraindication to ALL of the following:

- Aripiprazole
- Lamotrigine
- Lithium
- Lurasidone
- Olanzapine
- Risperidone
- Valproate

**OR**

**2.2** One of the following:

**2.2.1** The patient has been receiving treatment with the requested medication, and is new to the plan (enrollment effective date within the past 90 days)

**OR**

**2.2.2** The patient is currently receiving treatment with requested medication in the hospital and must continue upon discharge

**AND**

**3** - For Lybalvi requests ONLY, One of the following:

- Patient has a BMI of 30 kg/m<sup>2</sup> or greater
- Patient has a BMI of 27 kg/m<sup>2</sup> or greater with a weight-related comorbidity (e.g., dyslipidemia, hypertension, type 2 diabetes, sleep apnea)
- Patient has a documented history of weight gain of greater than or equal to 10% of their baseline weight after initiating antipsychotic medication
- Physician has provided clinical rationale for requiring samidorphan in addition to olanzapine therapy

Product Name: Caplyta, Vraylar	
Diagnosis	Bipolar Depression
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of bipolar depression</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - ONE of the following:</p> <p><b>2.1</b> Inadequate response (minimum 30 day trial) or adverse reaction to THREE, or contraindication to ALL of the following:</p> <ul style="list-style-type: none"> <li>• Lurasidone</li> <li>• Quetiapine</li> <li>• Olanzapine plus fluoxetine</li> <li>• Valproate monotherapy</li> </ul>	

- Combination therapy (i.e., lithium plus lamotrigine/valproate, lurasidone plus lithium/valproate, quetiapine plus lithium/valproate)

**OR**

**2.2** One of the following:

**2.2.1** The patient has been receiving treatment with the requested medication, and is new to the plan (enrollment effective date within the past 90 days)

**OR**

**2.2.2** The patient is currently receiving treatment with requested medication in the hospital and must continue upon discharge

Product Name: Rexulti, Vraylar	
Diagnosis	Major Depressive Disorder (MDD)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of one of the following:</p> <ul style="list-style-type: none"> <li>• Major depressive disorder (MDD)</li> <li>• Treatment resistant depression (Applies to Vraylar only)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - One of the following:</p> <p><b>2.1</b> BOTH of the following:</p> <p><b>2.1.1</b> Inadequate response (minimum 30 day trial) or adverse reaction to TWO different antidepressants from the following classes (antidepressants MUST be from TWO different classes):</p>	

- Selective Serotonin Reuptake Inhibitors (SSRIs)
- Serotonin Norepinephrine Reuptake Inhibitors (SNRIs)
- New Generation Antidepressants (e.g., bupropion, mirtazapine, etc)
- Tricyclic Antidepressants (TCAs)

**AND**

**2.1.2** Inadequate response (minimum 30 day trial) or adverse reaction to TWO of the following:

- aripiprazole
- quetiapine ER
- risperidone
- Augmented therapy (e.g., addition of lithium, another antidepressant from a different class, thyroid hormone to current antidepressant regimen)

**OR**

**2.2** One of the following:

**2.2.1** The patient has been receiving treatment with the requested medication, and is new to the plan (enrollment effective date within the past 90 days)

**OR**

**2.2.2** The patient is currently receiving treatment with the requested medication in the hospital and must continue upon discharge

Product Name:Rexulti	
Diagnosis	Agitation Associated With Dementia Due To Alzheimer's Disease
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - The requested medication is being used for treatment of agitation associated with dementia due to Alzheimer's disease

Product Name: Caplyta, Cobenfy, Fanapt, Lybalvi, Rexulti, Vraylar

Diagnosis	Requests Exceeding Quantity Limit*
-----------	------------------------------------

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Quantity Limit
----------------	----------------

**Approval Criteria**

1 - ONE of the following:

1.1 The requested drug must be used for a Food and Drug Administration (FDA)-approved indication

**OR**

1.2 The use of this drug is supported by information from one of the following appropriate compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmacoeconomic studies
- Other drug reference resources

**AND**

2 - ONE of the following:

**2.1** The drug is being prescribed within the manufacturer's published dosing guidelines

**OR**

**2.2** The requested dose falls within dosing guidelines found in one of the following compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmacoeconomic studies
- Other drug reference resources

**AND**

**3** - The requested dosage cannot be achieved using the plan accepted quantity limit of a different dose or formulation

**AND**

**4** - The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plans' program

**AND**

**5** - Physician has provided rationale for needing to exceed the quantity limit (QL) of one of the following:

- For Caplyta requests, QL of 1 capsule/day
- For Cobenfy capsule requests, QL of 2 capsules/day
- For Cobenfy starter pack requests, QL of 2 packs/365 days

<ul style="list-style-type: none"> <li>• For Fanapt tablet requests, QL of 2 tablets/day</li> <li>• For Fanapt titration pack requests, QL of 1 pack/180 days</li> <li>• For Lybalvi requests, QL of 1 tablet/day</li> <li>• For Rexulti requests, QL of 1 tablet/day</li> <li>• For Vraylar 1.5 mg and 3 mg requests, QL of 2 capsules/day</li> <li>• For Vraylar 4.5 mg and 6 mg requests, QL of 1 capsule/day</li> <li>• For Vraylar therapy pack requests, 2 packs/365 days</li> </ul>	
Notes	*Prior authorization requests should be reviewed using the above criteria. This section is for quantity limit requests only.

**2 . Revision History**

Date	Notes
12/6/2024	Added criteria for Cobenfy, updated guideline name

HIV (Fuzeon, Selzentry)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-161707
<b>Guideline Name</b>	HIV (Fuzeon, Selzentry)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Selzentry tablets, generic maraviroc 150mg and 300mg tablets, Selzentry oral solution	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 All of the following:  1.1.1 Diagnosis of CCR5-tropic HIV-1 infection as confirmed by a highly sensitive tropism assay	

**AND**

**1.1.2** Patient is currently taking or will be prescribed an optimized background antiretroviral therapy regimen

**AND**

**1.1.3** Prescribed by or in consultation with a clinician with HIV expertise

**OR**

**1.2** For continuation of prior therapy

**AND**

**2** - For Brand Selzentry ONLY; history of failure or intolerance to generic maraviroc tablets

Product Name:Fuzeon	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - One of the following:</p> <p><b>1.1</b> All of the following:</p> <p><b>1.1.1</b> Patient has been diagnosed with multidrug-resistant HIV-1 infection</p> <p><b>AND</b></p> <p><b>1.1.2</b> Patient is currently taking or will be prescribed an optimized background antiretroviral therapy regimen</p>	

**AND**

**1.1.3** Prescribed by or in consultation with a clinician with HIV expertise

**OR**

**1.2** For continuation of prior therapy

## **2 . Revision History**

Date	Notes
12/6/2024	generic maraviroc to preferred. Brand Selzentry to NP.

Hydroxychloroquine

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99465
<b>Guideline Name</b>	Hydroxychloroquine
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Plaquenil, generic hydroxychloroquine	
Guideline Type	Quantity Limit
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 Treatment of chronic discoid lupus erythematosus or systemic lupus erythematosus  <b>OR</b>	

**1.2** Treatment of rheumatoid arthritis

**OR**

**1.3** Prophylaxis of malaria in geographic areas where chloroquine resistance is not reported

**OR**

**1.4** Treatment of uncomplicated malaria

Notes	Authorization will be issued for 6 months up to a quantity of 120 tablets per 30 days.
-------	--

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Hyftor (sirolimus) topical gel

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114463
<b>Guideline Name</b>	Hyftor (sirolimus) topical gel
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Hyftor	
Approval Length	4 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of facial angiofibroma associated with tuberous sclerosis complex  <b>AND</b>	

2 - Patient is 6 years of age or older

**AND**

3 - Patient is not a candidate for laser therapy or surgical treatments

**AND**

4 - Prescribed by or in consultation with a dermatologist

Product Name:Hyftor	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy (e.g., improvement in size or redness of facial angiofibroma)	

## 2 . Revision History

Date	Notes
9/26/2022	New program

Igalmi (dexmedetomidine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-110775
<b>Guideline Name</b>	Igalmi (dexmedetomidine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/15/2022
-----------------	-----------

## 1 . Criteria

Product Name:Igalmi	
Approval Length	14 Days [A]
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following diagnoses: <ul style="list-style-type: none"><li>Schizophrenia</li><li>Bipolar I or II disorder</li></ul>	

**AND**

**2** - For the treatment of acute agitation

**AND**

**3** - Trial and failure, contraindication or intolerance to at least two preferred products used in acute agitation (e.g., olanzapine, ziprasidone)

**AND**

**4** - Patient is currently being managed with maintenance medication for their underlying disorder (e.g., aripiprazole, olanzapine, quetiapine, lithium, valproic acid)

## **2 . Revision History**

Date	Notes
8/4/2022	New Program

Ilaris (canakinumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300288
<b>Guideline Name</b>	Ilaris (canakinumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Ilaris	
Diagnosis	Periodic Fever Syndromes [Cryopyrin-Associated Periodic Syndromes (CAPS), Tumor Necrosis Factor Receptor Associated Periodic Syndrome (TRAPS), Hyperimmunoglobulin D Syndrome (HIDS)/Mevalonate Kinase Deficiency(MKD), Familial Mediterranean Fever(FMF)]
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of one of the following periodic fever syndromes:

- Cryopyrin-associated periodic syndromes (CAPS), including familial cold autoinflammatory syndrome (FCAS) and Muckle-Wells syndrome (MWS)
- Tumor necrosis factor (TNF) receptor associated periodic syndrome (TRAPS)
- Hyperimmunoglobulin D (Hyper-IgD) syndrome (HIDS/mevalonate kinase deficiency (MKD))
- Familial Mediterranean Fever (FMF)

**AND**

2 - Prescribed by or in consultation with one of the following:

- Rheumatologist
- Immunologist

**AND**

3 - Both of the following:

- Patient is not receiving concomitant treatment with Tumor Necrosis Factor (TNF) inhibitors (e.g., Enbrel [etanercept], Humira [adalimumab], Remicade [infliximab])
- Patient is not receiving concomitant treatment with Interleukin-1 inhibitor (e.g., Arcalyst [rilonacept], Kineret [anakinra])

**AND**

4 - Patients diagnosed with Familial Mediterranean Fever (FMF) have a history of failure, contraindication, or intolerance to colchicine (applies to diagnosis of FMF ONLY)

Product Name: Ilaris	
Diagnosis	Periodic Fever Syndrome [CAPS, TRAPS, HIDS/MKD, FMF]
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy [defined as a decrease in frequency or severity of attacks, or a decrease in index disease flare or normalization of CRP (C-reactive protein)]

**AND**

2 - Both of the following:

- Patient is not receiving concomitant treatment with Tumor Necrosis Factor (TNF) inhibitors (e.g., Enbrel [etanercept], Humira [adalimumab], Remicade [infliximab])
- Patient is not receiving concomitant treatment with Interleukin-1 inhibitor (e.g., Arcalyst [rilonacept], Kineret [anakinra])

Product Name:Ilaris	
Diagnosis	Systemic Juvenile Idiopathic Arthritis (SJIA)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of active systemic juvenile idiopathic arthritis (SJIA)	
<b>AND</b>	
2 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:	
<ul style="list-style-type: none"><li>• Minimum duration of a 3-month trial and failure of methotrexate</li></ul>	

- Minimum duration of a 1-month trial of a nonsteroidal anti-inflammatory drug (NSAID) (e.g., ibuprofen, naproxen)
- Minimum duration of a 2-week trial of a systemic glucocorticoid (e.g., prednisone)

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to a preferred tocilizumab biosimilar

**AND**

**4** - Both of the following:

- Patient is not receiving concomitant treatment with Tumor Necrosis Factor (TNF) inhibitors (e.g., Enbrel [etanercept], Humira [adalimumab], Remicade [infliximab])
- Patient is not receiving concomitant treatment with Interleukin-1 inhibitor (e.g., Arcalyst [rilonacept], Kineret [anakinra])

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Rheumatologist
- Immunologist

Product Name: Ilaris	
Diagnosis	Systemic Juvenile Idiopathic Arthritis (SJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Documentation of positive clinical response to therapy as evidenced by at least one of the following:	

- Reduction in the total active (swollen and tender) joint count from baseline
- Improvement in clinical features or symptoms (e.g., pain, fever, inflammation, rash, lymphadenopathy, serositis) from baseline

**AND**

**2 - Both of the following:**

- Patient is not receiving concomitant treatment with Tumor Necrosis Factor (TNF) inhibitors (e.g., Enbrel [etanercept], Humira [adalimumab], Remicade [infliximab])
- Patient is not receiving concomitant treatment with Interleukin-1 inhibitor (e.g., Arcalyst [rilonacept], Kineret [anakinra])

Product Name: Ilaris	
Diagnosis	Still's Disease
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of Still's Disease, including Adult-Onset Still's Disease (AOSD)</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>2 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one of the following: [1-3]</b></p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., prednisone)</li> <li>• Methotrexate</li> <li>• Nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen)</li> </ul> <p style="text-align: center;"><b>AND</b></p>	

**3 - Both of the following:**

- Patient is not receiving concomitant treatment with Tumor Necrosis Factor (TNF) inhibitors (e.g., Enbrel [etanercept], Humira [adalimumab], Remicade [infliximab])
- Patient is not receiving concomitant treatment with Interleukin-1 inhibitor (e.g., Arcalyst [rilonacept], Kineret [anakinra])

**AND**

**4 - Prescribed by or in consultation with one of the following:**

- Rheumatologist
- Immunologist

Product Name:Ilaris

Diagnosis	Still's Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1 - Documentation of positive clinical response to therapy**

**AND**

**2 - Both of the following:**

- Patient is not receiving concomitant treatment with Tumor Necrosis Factor (TNF) inhibitors (e.g., Enbrel [etanercept], Humira [adalimumab], Remicade [infliximab])
- Patient is not receiving concomitant treatment with Interleukin-1 inhibitor (e.g., Arcalyst [rilonacept], Kineret [anakinra])

Product Name:Ilaris

Diagnosis	Gout Flares
Approval Length	12 Week(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of gout flares</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"> <li>• Nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen)</li> <li>• Colchicine</li> <li>• Corticosteroids (e.g., prednisone)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient has not received Ilaris in the last 12 weeks [A]</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Rheumatologist</li> <li>• Nephrologist</li> </ul>	

## 2 . Definitions

Definition	Description
------------	-------------

Cryopyrin-Associated Periodic Syndromes (CAPS):	A group of rare, autosomal dominantly inherited auto-inflammatory conditions comprising of Familial-Cold Auto-inflammatory Syndrome (FCAS), Muckle-Wells Syndrome (MWS), Neonatal-Onset Multisystem Inflammatory Disease (NOMID) or also known as Chronic Infantile Neurologic Cutaneous Articular Syndrome (CINCA), which are caused by the CIAS1 gene mutation and characterized by recurrent symptoms (urticaria-like skin lesions, fever chills, arthralgia, profuse sweating, sensorineural hearing/vision loss, and increased inflammation markers the blood). Approximately 300 people in the United States are affected by CAPS. [1, 4, 5]
Familial Cold Autoinflammatory Syndrome (FCAS):	The mildest form of CAPS, is characterized by cold-induced, daylong episodes of fever associated with rash, arthralgia, headaches and less frequently conjunctivitis, but without other signs of CNS inflammation. Symptoms usually begin during the first 6 months of life and are predominantly triggered by cold exposure. Duration of episodes usually is less than 24 hours. [5]
Muckle-Wells Syndrome (MWS):	A subtype of CAPS, which is characterized by episodic attacks of inflammation associated with a generalized urticaria-like rash, fever, malaise, arthralgia, and progressive hearing loss. Duration of symptoms usually lasts from 24-48 hours. [5]

**3 . Endnotes**

- A. The recommended dose of Ilaris for adult patients with a gout flare is 150 mg administered subcutaneously. In patients who require re-treatment, there should be an interval of at least 12 weeks before a new dose of Ilaris may be administered [1].

**4 . Revision History**

Date	Notes
7/3/2025	SJIA: Added step through preferred tocilizumab biosimilar

Ilumya (tildrakizumab-asmn)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152432
<b>Guideline Name</b>	Ilumya (tildrakizumab-asmn)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/20/2024
-----------------	-----------

## 1 . Criteria

Product Name:Ilumya	
Diagnosis	Chronic Moderate to Severe Plaque Psoriasis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 ALL of the following:  1.1.1 Diagnosis of chronic moderate to severe plaque psoriasis	

**AND**

**1.1.2** Greater than or equal to 3 percent body surface area involvement, palmoplantar, facial, genital involvement, or severe scalp psoriasis

**AND**

**1.1.3** Submission of medical records (e.g., chart notes) or paid claims documenting history of failure, to ONE of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Anthralin
- Coal tar

**AND**

**1.1.4** Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.1.5** Submission of medical records (e.g., chart notes) or paid claims documenting history of failure, contraindication, or intolerance to ALL of the following preferred biologic products:

- A preferred adalimumab biosimilar
- Enbrel (etanercept)
- Otezla (apremilast)

**AND**

**1.1.6** Patient is NOT receiving Ilumya in combination with ONE of the following:

- Biologic DMARD (disease modifying anti-rheumatic drug) [e.g., Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab), Cosentyx (secukinumab), Orencia (abatacept)]
- Janus kinase inhibitor [e.g., Xeljanz (tofacitinib)]
- Phosphodiesterase 4 (PDE4) inhibitor [e.g. Otezla (apremilast)]

**AND**

**1.1.7** Prescribed by or in consultation with a dermatologist

**OR**

**1.2** ALL of the following:

**1.2.1** Submission of medical records (e.g., chart notes) or paid claims documenting patient is currently on the requested therapy

**AND**

**1.2.2** Diagnosis of chronic moderate to severe plaque psoriasis

**AND**

**1.2.3** Patient is NOT receiving Ilumya in combination with ONE of the following:

- Biologic DMARD [e.g., Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab), Cosentyx (secukinumab), Orencia (abatacept)]
- Janus kinase inhibitor [e.g., Xeljanz (tofacitinib)]
- Phosphodiesterase 4 (PDE4) inhibitor [e.g. Otezla (apremilast)]

**AND**

**1.2.4** Prescribed by or in consultation with a dermatologist

Product Name: Ilumya	
Diagnosis	Chronic Moderate to Severe Plaque Psoriasis

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Ilumya therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is NOT receiving Ilumya in combination with ONE of the following:</p> <ul style="list-style-type: none"> <li>• Biologic DMARD (disease modifying anti-rheumatic drug) [e.g., Humira (adalimumab), Cimzia (certolizumab), Simponi (golimumab), Cosentyx (secukinumab), Orencia (abatacept)]</li> <li>• Janus kinase inhibitor [e.g., Xeljanz (tofacitinib)]</li> <li>• Phosphodiesterase 4 (PDE4) inhibitor [e.g. Otezla (apremilast)]</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a dermatologist</p>	

## 2 . Revision History

Date	Notes
8/20/2024	Updated submission of medical records verbiage in criteria

Imcivree (setmelanotide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-163778
<b>Guideline Name</b>	Imcivree (setmelanotide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Imcivree	
Diagnosis	POMC, PCSK1, LEPR Deficiency
Approval Length	n/a- requests for indications other than Bardet-Biedl syndrome (BBS) are excluded from coverage and will not be approved.
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Requests for indications other than Bardet-Biedl syndrome (BBS) are excluded from coverage and will not be approved.	
Notes	Requests for indications other than Bardet-Biedl syndrome (BBS) are excluded from coverage and will not be approved.

Product Name:Imcivree	
Diagnosis	Bardet-Biedl syndrome (BBS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting ALL of the following:</p> <p>1.1 Both of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of Bardet-Biedl syndrome (BBS)</li> <li>• Molecular genetic testing to confirm homozygous variants in a BBS gene that are interpreted as pathogenic or likely pathogenic (results of genetic testing must be submitted)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>1.2 One of the following:</p> <p>1.2.1 Patient has at least three of the following primary features of the disease:</p> <ul style="list-style-type: none"> <li>• Rod-cone dystrophy</li> <li>• polydactyly</li> <li>• learning disabilities</li> <li>• hypogonadotropic hypogonadism and/or genitourinary anomalies</li> <li>• renal anomalies</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>1.2.2 Both of the following:</p> <p>1.2.2.1 Patient has at least two of the following primary features of the disease:</p> <ul style="list-style-type: none"> <li>• Rod-cone dystrophy</li> <li>• polydactyly</li> <li>• learning disabilities</li> </ul>	

- hypogonadotropic hypogonadism and/or genitourinary anomalies
- renal anomalies

**AND**

**1.2.2.2** Patient has at least two of the following secondary features of the disease:

- Speech disorder/delay
- strabismus/cataracts/astigmatism
- brachydactyly/syndactyly
- developmental delay
- ataxia/poor coordination/imbalance
- mild spasticity (especially lower limbs)
- diabetes mellitus
- dental crowding/hypodontia/small roots/high arched palate
- left ventricular hypertrophy/congenital heart disease
- hepatic fibrosis

**AND**

**1.3** Patient has been diagnosed with obesity defined by one of the following:

- BMI greater than or equal to 30 kg/m<sup>2</sup> for adults 18 years of age or older
- Weight greater than or equal to 95th percentile using growth chart assessments for pediatric patients

**AND**

**1.4** Patient is 2 years of age or older

**AND**

**1.5** Other causes or types of obesity have been ruled out (e.g., obesity due to suspected POMC, PCSK1, or LEPR deficiency with POMC, PCSK1, or LEPR variants classified as benign or likely benign; obesity associated with other genetic syndromes; polygenic obesity)

**AND**

**2** - Prescribed by or in consultation with an endocrinologist

Notes	Requests for indications other than Bardet-Biedl syndrome (BBS) are excluded from coverage and will not be approved.
-------	--

Product Name:Imcivree	
Diagnosis	Bardet-Biedl syndrome (BBS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting BOTH of the following:</p> <p>1.1 Patient has been on therapy for 12 months or more</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Weight loss of greater than or equal to 5% of baseline body weight or BMI</p>	
Notes	Requests for indications other than Bardet-Biedl syndrome (BBS) are excluded from coverage and will not be approved.

## 2 . Revision History

Date	Notes
1/30/2025	Updated age criterion for BBS indication due to expanded approval.

Immune Globulins - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147675
<b>Guideline Name</b>	Immune Globulins - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Asthma (severe, persistent, high-dose steroid-dependent)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following diagnoses:	

- Severe asthma
- Persistent asthma
- High-dose steroid-dependent asthma

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Patient is receiving optimal conventional asthma therapy (e.g., high-dose inhaled glucocorticoids, short- and long-acting inhaled  $\beta$  agonists)

**AND**

**4** - History of failure, contraindication, or intolerance to at least TWO of the following:

- Anti-IgE therapy [e.g., Xolair (omalizumab)]
- Anti-interleukin 4 therapy [e.g., Dupixent (dupilumab)]
- Anti-interleukin 5 therapy [e.g., Nucala (mepolizumab), Cinqair (reslizumab), Fasenra (benralizumab)]

**AND**

**5** - Patient has required continuous oral glucocorticoid therapy for a minimum of 2 months prior to the decision to initiate immune globulin therapy

**AND**

**6** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

**AND**

**7** - Prescribed by or in consultation with a pulmonologist or allergist or immunologist

**AND**

**8** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Asthma (severe, persistent, high-dose steroid-dependent)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to immune globulin therapy

**AND**

**2** - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

**3** - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Autoimmune Bullous Disease [pemphigus vulgaris, pemphigus foliaceus, bullous pemphigoid, mucous membrane (cicatricial) pemphigoid, epidermolysis bullosa acquisita, pemphigoid gestationis, linear IgA bullous dermatosis]
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of Autoimmune Bullous Disease [pemphigus vulgaris, pemphigus foliaceus, bullous pemphigoid, mucous membrane (cicatricial) pemphigoid, epidermolysis bullosa acquisita, pemphigoid gestationis, linear IgA bullous dermatosis]

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Extensive and debilitating disease

**AND**

**4** - History of failure, contraindication, or intolerance to systemic corticosteroids with concurrent immunosuppressive treatment (e.g., azathioprine, cyclophosphamide, mycophenolate mofetil)

**AND**

**5** - Intravenous immunoglobulin (IVIG) dose does not exceed 1,000 to 2,000 milligrams (mg) per kilogram (kg) per month divided into 3 equal doses, each given over 3 consecutive days or 400 mg per kg per day given over 5 consecutive days per month. IVIG administration may be repeated monthly as needed for patients requiring maintenance therapy. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**6** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

**AND**

**7** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Autoimmune Bullous Disease [pemphigus vulgaris, pemphigus foliaceus, bullous pemphigoid, mucous membrane (cicatricial) pemphigoid, epidermolysis bullosa acquisita, pemphigoid gestationis, linear IgA bullous dermatosis]
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Bone Marrow Transplant (BMT)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following uses:</p> <ul style="list-style-type: none"> <li>Prevention of acute graft vs. host disease (GVHD)</li> </ul>	

- Prevention of infection

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Confirmed allogeneic bone marrow transplant within the last 100 days

**AND**

**4** - Documented severe hypogammaglobulinemia [Immunoglobulin (IgG) less than 400 milligrams (mg) per deciliter (dL)]

**AND**

**5** - Intravenous immunoglobulin (IVIG) dose does not exceed 500 mg per kilogram (kg) once weekly for the first 90 days of therapy, then monthly up to 360 days after transplantation

**AND**

**6** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen

- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Bone Marrow Transplant (BMT)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Chronic Inflammatory Demyelinating Polyneuropathy
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Diagnosis of chronic inflammatory demyelinating polyneuropathy as confirmed by ALL of the following:

**1.1** Progressive symptoms present for at least 2 months

**AND**

**1.2** Symptomatic polyradiculoneuropathy as indicated by progressive or relapsing motor or sensory impairment of more than one limb

**AND**

**1.3** Electrodiagnostic findings [consistent with European Federation of Neurological Societies/Peripheral Nerve Society (EFNS/PNS) guidelines for definite chronic inflammatory demyelinating polyradiculoneuropathy (CIDP)] indicating at least ONE of the following criteria are present:

- Motor distal latency prolongation in 2 nerves
- Reduction of motor conduction velocity in 2 nerves
- Prolongation of F-wave latency in 2 nerves
- Absence of F-waves in at least 1 nerve
- Partial motor conduction block of at least 1 motor nerve
- Abnormal temporal dispersion in at least 2 nerves
- Distal compound muscle action potential (CMAP) duration increase in at least 1 nerve

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Prescribed by or in consultation with a neurologist

**AND**

**4** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 consecutive days administered in up to six monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities.

**AND**

**5** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Chronic Inflammatory Demyelinating Polyneuropathy
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to therapy as measured by an objective scale [e.g., Rankin, Modified Rankin, Medical Research Council (MRC) scale]

**AND**

**2** - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

**AND**

**3** - Prescribed by or in consultation with a neurologist

**AND**

**4** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 consecutive days. IVIG administration may be repeated monthly as needed to prevent exacerbation. Dosing interval may need to be adjusted in patients with severe comorbidities.

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Prevention of infection in B-cell Chronic Lymphocytic Leukemia (CLL)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of B-cell chronic lymphocytic leukemia (CLL)

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3 - ONE of the following:**

- Documented hypogammaglobulinemia [Immunoglobulin (IgG) less than 500 milligrams (mg) per deciliter (dL)]
- History of bacterial infection(s) associated with B-cell CLL

**AND**

**4 - Intravenous immunoglobulin (IVIG) dose does not exceed 400 milligrams (mg) per kilogram (kg) every 3 to 4 weeks**

**AND**

**5 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:**

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Prevention of infection in B-cell Chronic Lymphocytic Leukemia (CLL)
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Dermatomyositis or polymyositis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of dermatomyositis or polymyositis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:</p> <ul style="list-style-type: none"> <li>• History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable</li> </ul>	

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - History of failure, contraindication, or intolerance to immunosuppressive therapy (e.g., azathioprine, corticosteroids, cyclophosphamide, methotrexate)

**AND**

**4** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 consecutive days administered as monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**5** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

**AND**

**6** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Dermatomyositis or polymyositis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Diabetes Mellitus
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is newly diagnosed with insulin dependent (type 1) diabetes mellitus</p> <p style="text-align: center;"><b>AND</b></p>	

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Patient is not a candidate for or is refractory to insulin therapy

**AND**

**4** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Diabetes Mellitus
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammalex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Feto-neonatal Alloimmune Thrombocytopenia (AIT)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - One of the following:

1.1 For pregnant women ALL of the following:

1.1.1 Diagnosis of feto-neonatal alloimmune thrombocytopenia (AIT)

**AND**

1.1.2 ONE of the following:

- Previously affected pregnancy
- Family history of the disease
- Platelet alloantibodies found on screening

**AND**

**1.1.3** ONE of the following:

**1.1.3.1** Intravenous immunoglobulin (IVIG) dose does not exceed 1,000 milligrams (mg) per kilogram (kg) once weekly until delivery

**OR**

**1.1.3.2** BOTH of the following:

- Fetus or newborn is considered to be at high risk for developing intracranial hemorrhage or other severe complication of AIT
- IVIG dose does not exceed 2,000 mg/kg once weekly until delivery

**OR**

**1.2** For newborns BOTH of the following:

**1.2.1** Diagnosis of feto-neonatal alloimmune thrombocytopenia

**AND**

**1.2.2** Thrombocytopenia that persists after transfusion of antigen-negative compatible platelets

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammalex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Feto-neonatal Alloimmune Thrombocytopenia (AIT)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to immune globulin therapy

**AND**

**2** - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

**3** - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Graves' ophthalmopathy Guillain-Barré syndrome (GBS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of Guillain-Barré Syndrome

**AND**

2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

3 - Severe disease requiring aid to walk

**AND**

4 - Onset of neuropathic symptoms within the last four weeks

**AND**

5 - Prescribed by or in consultation with a neurologist

**AND**

**6** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 consecutive days. IVIG administration may be repeated in up to three monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**7** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

**AND**

**8** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Graves' ophthalmopathy Guillain-Barré syndrome (GBS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to immune globulin therapy	

**AND**

**2** - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

**3** - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Prevention of bacterial infection in pediatric HIV
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of HIV disease

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Patient age less than or equal to 13 years of age

**AND**

**4 - ONE of the following:**

- Documented hypogammaglobulinemia [Immunoglobulin (IgG) less than 400 milligrams (mg) per deciliter (dL)]
- Functional antibody deficiency as demonstrated by either poor specific antibody titers or recurrent bacterial infections

**AND**

**5 - Intravenous immunoglobulin (IVIG ) dose does not exceed 400 mg per kilogram (kg) every 28 days**

**AND**

**6 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:**

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Prevention of bacterial infection in pediatric HIV
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Immune thrombocytopenia [Idiopathic thrombocytopenic purpura (ITP)]
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - One of the following:	
1.1 ALL of the following:	
<ul style="list-style-type: none"><li>• Diagnosis of acute thrombocytopenic purpura (ITP)</li><li>• Documented platelet count less than <math>50 \times 10^9</math> per Liter (L) (obtained within the past 30 days)</li><li>• Intravenous immunoglobulin (IVIG) dose does not exceed 1,000 milligrams (mg) per kilogram(kg) per day for 1 to 2 days</li></ul>	

**OR**

**1.2** All of the following:

**1.2.1** Diagnosis of chronic thrombocytopenic purpura (ITP)

**AND**

**1.2.2** History of failure, contraindication, or intolerance to at least ONE of the following:

- Corticosteroids
- Splenectomy

**AND**

**1.2.3** IVIG dose does not exceed 2,000 mg per kg per month given over 2 to 5 consecutive days. IVIG administration may be repeated monthly as needed to prevent exacerbation. Dosing interval should be adjusted depending upon response and titrated to the minimum effective dose that can be given at maximum intervals to maintain safe platelet levels.

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked

- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Immune thrombocytopenia [Idiopathic thrombocytopenic purpura (ITP)]
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Kawasaki Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization

Guideline Type	Prior Authorization
<p data-bbox="196 365 438 399"><b>Approval Criteria</b></p> <p data-bbox="196 432 646 466">1 - Diagnosis of Kawasaki disease</p> <p data-bbox="773 537 837 571" style="text-align: center;"><b>AND</b></p> <p data-bbox="196 642 1341 676">2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:</p> <ul data-bbox="245 714 1390 844" style="list-style-type: none"><li>• History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable</li><li>• Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested</li></ul> <p data-bbox="773 915 837 949" style="text-align: center;"><b>AND</b></p> <p data-bbox="196 1020 1325 1087">3 - Intravenous immunoglobulin (IVIG) dose does not exceed 4,000 milligrams (mg) per kilograms (kg) for five consecutive days or a single dose of 2,000 mg per kg</p> <p data-bbox="773 1159 837 1192" style="text-align: center;"><b>AND</b></p> <p data-bbox="196 1264 1237 1331">4 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:</p> <ul data-bbox="245 1369 545 1701" style="list-style-type: none"><li>• Bivigam</li><li>• Flebogamma</li><li>• Gammagard Liquid</li><li>• Gammagard S-D</li><li>• Gammaked</li><li>• Gamunex-C</li><li>• Hizentra</li><li>• Octagam</li><li>• Privigen</li><li>• Xembify</li></ul>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Kawasaki Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Lambert-Eaton Myasthenic Syndrome (LEMS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of Lambert-Eaton myasthenic syndrome (LEMS)</p>	

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - History of failure, contraindication, or intolerance to immunomodulator monotherapy (e.g., azathioprine, corticosteroids)

**AND**

**4** - Concomitant immunomodulator therapy (e.g., azathioprine, corticosteroids), unless contraindicated, will be used for long-term management of LEMS

**AND**

**5** - Prescribed by or in consultation with a neurologist

**AND**

**6** - Intravenous Immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 consecutive days. IVIG administration may be repeated monthly as needed to prevent exacerbation. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**7** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

**AND**

**8** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Lambert-Eaton Myasthenic Syndrome (LEMS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to immune globulin therapy

**AND**

**2** - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

**3** - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Lennox Gastaut Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - History of failure, contraindication or intolerance to initial treatment with traditional anti-epileptic pharmacotherapy (e.g., lamotrigine, phenytoin, valproic acid)

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Prescribed by or in consultation with a neurologist

**AND**

**4** - Intravenous immunoglobulin (IVIG) dose does not exceed 400 milligrams (mg) per kilogram (kg) per day given for 4 to 5 consecutive days. IVIG administration may be repeated monthly as needed in patients requiring maintenance therapy. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**5** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

**AND**

**6** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Lennox Gastaut Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Multifocal Motor Neuropathy (MMN)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of multifocal motor neuropathy as confirmed by ALL of the following:

- Weakness with slowly progressive or stepwise progressive course over at least one month
- Asymmetric involvement of two or more nerves
- Absence of motor neuron signs and bulbar signs

**AND**

2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

3 - Prescribed by or in consultation with a neurologist

**AND**

**4** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,400 milligram (mg) per kilogram (kg) per month given over 2 to 5 consecutive days. IVIG administration may be repeated monthly as needed to prevent exacerbation. Dosing interval may need to be adjusted in patients with severe comorbidities.

**AND**

**5** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Multifocal Motor Neuropathy (MMN)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to therapy as measured by an objective scale [e.g., Rankin, Modified Rankin, Medical Research Council (MRC) scale]

**AND**

**2** - Prescribed by or in consultation with a neurologist

**AND**

**3** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,400 milligram (mg) per kilogram (kg) per month given over 2 to 5 consecutive days. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**4** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Prevention of infection in Multiple Myeloma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of multiple myeloma

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3 - ONE of the following:**

- Documented hypogammaglobulinemia [immunoglobulin (IgG) less than 500 milligrams (mg) per deciliter (dL)]
- History of bacterial infection(s) associated with multiple myeloma

**AND**

**4 - Intravenous immunoglobulin (IVIG) dose does not exceed 400 mg per kilogram (kg) every 3 to 4 weeks**

**AND**

**5 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:**

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Prevention of infection in Multiple Myeloma
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Relapsing Multiple Sclerosis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of relapsing forms of multiple sclerosis (MS) (e.g., relapsing-remitting MS, secondary- progressive MS with relapses, progressive-relapsing MS with relapses)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:</p> <ul style="list-style-type: none"> <li>• History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable</li> </ul>	

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Documentation of an MS exacerbation or progression (worsening) of the patient's clinical status from the visit prior to the one prompting the decision to initiate immune globulin therapy

**AND**

**4** - History of failure, contraindication, or intolerance to at least TWO of the following agents:

- Aubagio (teriflunomide)
- Avonex (interferon beta-1a)
- Betaseron (interferon beta-1b)
- Copaxone/Glatopa (glatiramer acetate)
- Extavia (interferon beta-1b)
- Gilenya (fingolimod)
- Lemtrada (alemtuzumab)
- Mavenclad (cladribine)
- Mayzent (siponimod)
- Ocrevus (ocrelizumab)
- Plegridy (peginterferon beta-1a)
- Rebif (interferon beta-1a)
- Tecfidera (dimethyl fumarate)
- Tysabri (natalizumab)

**AND**

**5** - Prescribed by or in consultation with a neurologist

**AND**

**6** - Induction, when indicated, does not exceed a dose of 400 milligrams (mg) per kilogram (kg) daily for up to five days

**AND**

7 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammalex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Relapsing Multiple Sclerosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Medical records, including findings of interval examination including neurological deficits incurred and assessment of disability [e.g., Expanded Disability Status Scale (EDSS), Functional Systems Score (FSS), Multiple Sclerosis Functional Composite (MSFC), Disease Steps (DS)]

**AND**

2 - Stable or improved disability score (e.g., EDSS, FSS, MSFC, DS)

**AND**

3 - Documentation of decreased number of relapses since starting immune globulin therapy

**AND**

**4** - Diagnosis continues to be the relapsing forms of multiple sclerosis (MS)

**AND**

**5** - Prescribed by or in consultation with a neurologist

**AND**

**6** - Intravenous immunoglobulin (IVIG) dose does not exceed 1,000 milligram (mg) per kilogram (kg) monthly

**AND**

**7** - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammalex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Myasthenia Gravis - Exacerbation
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of generalized myasthenia gravis

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Evidence of myasthenia exacerbation, defined by at least ONE of the following symptoms in the last month

- Difficulty swallowing
- Acute respiratory failure
- Major functional disability responsible for the discontinuation of physical activity
- Recent immunotherapy treatment with a checkpoint inhibitor [e.g., Keytruda (pembrolizumab), Opdivo (nivolumab), Tecentriq (atezolizumab)]

**AND**

**4** - ONE of the following:

- History of failure, contraindication, or intolerance to immunomodulator therapy (e.g., azathioprine, mycophenolate mofetil, cyclosporine) for long-term management of myasthenia gravis
- Currently receiving immunomodulator therapy (e.g., azathioprine, mycophenolate mofetil, cyclosporine) for long-term management of myasthenia gravis

**AND**

**5** - Prescribed by or in consultation with a neurologist

**AND**

**6** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 days administered in up to three monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities.

**AND**

**7** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Refractory Myasthenia Gravis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of refractory generalized myasthenia gravis by or in consultation with a physician or center with expertise in management of myasthenia gravis

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Documentation that the disease status is unchanged or worsening (persistent or worsening symptoms that limit functioning) despite failure, contraindication, or intolerance to BOTH of the following (used in adequate doses and duration):

- Corticosteroids
- Two immunomodulator therapies (e.g., azathioprine, mycophenolate mofetil, cyclosporine, methotrexate, tacrolimus)

**AND**

**4** - Currently receiving immunomodulator therapy (e.g., corticosteroids, azathioprine, mycophenolate mofetil, cyclosporine, methotrexate, tacrolimus), used in adequate doses, for long-term management of myasthenia gravis

**AND**

**5** - Prescribed by or in consultation with a neurologist

**AND**

**6** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 days administered in up to three monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities.

**AND**

**7** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked

- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Myasthenia Gravis –Exacerbation and Refractory Myasthenia Gravis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Neuromyelitis Optica
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, laboratory values, etc.) to support the diagnosis of neuromyelitis optica spectrum disorder (NMOSD) by a neurologist confirming ALL of the following:

**1.1** Serologic testing for anti-aquaporin-4 immunoglobulin G (AQP4-IgG) or Neuromyelitis optica immunoglobulin G (NMO-IgG) antibodies has been performed

**AND**

**1.2** ONE of the following:

**1.2.1** If AQP4-IgG/NMO-IgG positive, past medical history of ONE of the following:

- Optic neuritis
- Acute myelitis
- Area postrema syndrome: episode of otherwise unexplained hiccups or nausea and vomiting
- Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions
- Symptomatic cerebral syndrome with NMOSD-typical brain lesions

**OR**

**1.2.2** If AQP4-IgG/NMO-IgG negative, past medical history of TWO of the following:

- Optic neuritis
- Acute myelitis
- Area postrema syndrome: episode of otherwise unexplained hiccups or nausea and vomiting
- Acute brainstem syndrome
- Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions
- Symptomatic cerebral syndrome with NMOSD-typical brain lesions

**AND**

**1.3** Diagnosis of multiple sclerosis or other diagnoses have been ruled out

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - History of failure, contraindication, or intolerance to at least TWO of the following:

- Azathioprine
- Corticosteroids
- Mycophenolate mofetil
- Rituximab
- Soliris (eculizumab)

**AND**

**4** - Patient is not receiving immune globulin in combination with either of the following:

- Rituximab
- Soliris (eculizumab)

**AND**

**5** - Prescribed by or in consultation with a neurologist

**AND**

**6** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligram (mg) per kilogram (kg) per month given over 2 to 5 days administered in up to six monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities.

**AND**

7 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Neuromyelitis Optica
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient has previously been treated with immune globulin

**AND**

2 - Submission of medical records (e.g., chart notes, laboratory tests) to demonstrate a positive clinical response from baseline as demonstrated by BOTH of the following:

2.1 Reduction in the number and or severity of relapses or signs and symptoms of neuromyelitis optica spectrum disorder (NMOSD)

**AND**

**2.2** Maintenance, reduction, or discontinuation of dose(s) of any baseline immunosuppressive therapy (IST) prior to starting immune globulin. (NOTE: Add on, dose escalation of IST, or additional rescue therapy from baseline to treat NMOSD or exacerbation of symptoms while on immune globulin therapy will be considered as treatment failure.)

**AND**

**3** - Patient is not receiving immune globulin in combination with either of the following:

- Rituximab
- Soliris (eculizumab)

**AND**

**4** - Prescribed by or in consultation with a neurologist

**AND**

**5** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 days administered in up to six monthly infusions. Dosing interval may need to be adjusted in patients with severe comorbidities

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Posttransfusion Purpura
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of posttransfusion purpura

**AND**

2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

3 - Intravenous immunoglobulin (IVIG) dose does not exceed 1,000 milligrams (mg) per kilogram (kg) for 2 days

**AND**

4 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Posttransfusion Purpura
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Post B-Cell Targeted Therapies
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation confirming previous treatment of B-cell targeted therapy within the last 100 days [e.g., CAR-T (e.g., Kymriah), Rituxan (rituximab), Besponsa (inotuzumab ozogamicin)]</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:</p> <ul style="list-style-type: none"> <li>History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable</li> </ul>	

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3 - BOTH of the following:**

- Documented hypogammaglobulinemia [immunoglobulin (IgG) less than 500 milligrams (mg) per deciliter (dL)]
- History of bacterial infection(s) associated with B-cell depletion

**AND**

**4 - Intravenous immunoglobulin (IVIG) dose does not exceed 400 mg per kilogram (kg) every 4 weeks, up to 360 days after discontinuation of B-cell depleting therapy**

**AND**

**5 - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:**

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Post B-Cell Targeted Therapies
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Primary Immunodeficiency Syndromes
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of primary immunodeficiency</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:</p> <ul style="list-style-type: none"> <li>• History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable</li> </ul>	

- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Clinically significant functional deficiency of humoral immunity as evidenced by ONE of the following:

- Documented failure to produce antibodies to specific antigens
- History of significant recurrent infections

**AND**

**4** - Initial intravenous immunoglobulin (IVIG) dose is 200 to 800 milligrams (mg) per kilogram (kg) every 3 to 4 weeks, based on product prescribing information, and titrated based upon patient response (For subcutaneous immune globulin (SCIG) products, FDA-labeled dosing and conversion guidelines will be used to determine benefit coverage.)

**AND**

**5** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Primary Immunodeficiency Syndromes
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to immune globulin therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Statement of expected frequency and duration of proposed immune globulin treatment</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response</p>	

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	Rasmussen Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of ONE of the following demonstrating that:</p> <ul style="list-style-type: none"> <li>• Short term amelioration of encephalitis is needed prior to definitive surgical therapy</li> <li>• Disease symptoms (e.g., seizures) persist despite surgical treatment</li> <li>• The patient is not a candidate for surgical treatment</li> </ul> <p style="text-align: center;"><b>AND</b></p>	

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 days

**AND**

**4** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Rasmussen Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to immune globulin therapy

**AND**

2 - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

3 - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammalex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Stiff-Person Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of stiff-person syndrome

**AND**

2 - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - History of failure, contraindication or intolerance to GABAergic (gamma-aminobutyric acid analogs) medication (e.g., baclofen, benzodiazepines)

**AND**

**4** - Prescribed by or in consultation with a neurologist

**AND**

**5** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 days. IVIG administration may be repeated monthly as needed for patients requiring maintenance therapy. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

**6** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Stiff-Person Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of a positive clinical improvement from baseline

**AND**

2 - Prescribed by or in consultation with a neurologist

**AND**

3 - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligrams (mg) per kilogram (kg) per month given over 2 to 5 days. IVIG administration may be repeated monthly as needed for patients requiring maintenance therapy. Dosing interval may need to be adjusted in patients with severe comorbidities

**AND**

4 - For long term treatment, documentation of titration to the minimum dose and frequency needed to maintain a sustained clinical effect

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Thrombocytopenia, secondary to Hepatitis C Virus (HCV), Human Immunodeficiency Virus (HIV), or pregnancy
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - One of the following:

1.1 Both of the following:

- Diagnosis of thrombocytopenia secondary to Hepatitis C Virus (HCV) infection
- Patient is receiving concurrent antiviral therapy, unless contraindicated

**OR**

**1.2** Both of the following:

- Diagnosis of thrombocytopenia secondary Human Immunodeficiency Virus (HIV) infection
- Patient is receiving concurrent antiviral therapy, unless contraindicated

**OR**

**1.3** Diagnosis of thrombocytopenia secondary to pregnancy

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - Documented platelet count less than  $50 \times 10^9$  per liter (L) (obtained within the past 30 days)

**AND**

**4** - Intravenous immunoglobulin (IVIG) dose does not exceed 1,000 milligrams (mg) per kilogram (kg) per day for 1 to 2 days

**AND**

**5** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	Thrombocytopenia, secondary to Hepatitis C Virus (HCV), Human Immunodeficiency Virus (HIV), or pregnancy
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - One of the following:

**1.1** Both of the following:

- Diagnosis of thrombocytopenia secondary to Hepatitis C Virus (HCV) infection
- Patient is receiving concurrent antiviral therapy, unless contraindicated

**OR**

**1.2** Both of the following:

- Diagnosis of thrombocytopenia secondary Human Immunodeficiency Virus (HIV) infection

- Patient is receiving concurrent antiviral therapy, unless contraindicated

**OR**

**1.3** Diagnosis of thrombocytopenia secondary to pregnancy

**AND**

**2** - Intravenous immunoglobulin (IVIG) dose does not exceed 2,000 milligram (mg) per kilogram (kg) per month given over 2 to 5 consecutive days. IVIG administration may be repeated monthly as needed to prevent exacerbation. Dosing interval should be adjusted depending upon response and titrated to the minimum effective dose that can be given at maximum intervals to maintain safe platelet levels.

Product Name: Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify

Diagnosis	All other indications
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - One of the following diagnoses:

- Autoimmune Uveitis
- Cytomegalovirus (CMV) induced pneumonitis in solid organ transplants
- Enteroviral Meningoencephalitis
- IgM antimyelin-associated glycoprotein paraprotein-associated peripheral neuropathy
- Lymphoproliferative disease (treatment of bacterial infections)
- Monoclonal gammopathy
- Paraproteinemic neuropathy
- Renal transplantation (prevention or treatment of acute humoral rejection)
- Severe Rheumatoid arthritis
- Rotaviral enterocolitis
- Staphylococcal toxic shock
- Toxic epidermal necrolysis or Stevens-Johnson syndrome

- Urticaria (delayed pressure)

**AND**

**2** - Submission of medical records(e.g., chart notes) documenting BOTH of the following:

- History and physical examination documenting the severity of the condition, including frequency and severity of infections where applicable
- Laboratory results or diagnostic evidence supporting the indication for which immune globulin is requested

**AND**

**3** - If the request is for a non-preferred product, there must be a history of failure, contraindication or intolerance to ALL the following products:

- Bivigam
- Flebogamma
- Gammagard Liquid
- Gammagard S-D
- Gammaked
- Gamunex-C
- Hizentra
- Octagam
- Privigen
- Xembify

Product Name:Alyglo, HyQvia, Gammagard S-D, Gammagard liquid, Cuvitru, Gamastan, Gamastan S-D, Gamunex-C, Carimune NF Nanofiltered, Privigen, Hizentra, Bivigam, Flebogamma Dif, Gammaplex, Octagam, Panzyga, Gammaked, Xembify	
Diagnosis	All other indications
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to immune globulin therapy	

**AND**

**2** - Statement of expected frequency and duration of proposed immune globulin treatment

**AND**

**3** - For long term treatment, documentation of titration to the minimum effective dose and frequency needed to maintain a sustained clinical response

## **2 . Revision History**

Date	Notes
5/23/2024	Added Alyglo as NP target

Inbrija

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99466
<b>Guideline Name</b>	Inbrija
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name: Inbrija	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Parkinson's disease  <b>AND</b>	

**2** - Inbrija will be used as intermittent treatment for OFF episodes

**AND**

**3** - Prescribed by, or in consultation with, a neurologist or specialist in the treatment of Parkinson's disease

**AND**

**4** - Patient is currently on a stable dose of a carbidopa/levodopa-containing medication and will continue receiving treatment with a carbidopa/levodopa-containing medication while on therapy

**AND**

**5** - Patient continues to experience greater than or equal to 2 hours of OFF time per day despite optimal management of carbidopa/levodopa therapy including BOTH of the following:

- Taking carbidopa/levodopa on an empty stomach or at least one half-hour or more before or one hour after a meal or avoidance of high protein diet
- Dose and dosing interval optimization

**AND**

**6** - History of failure, contraindication, or intolerance to TWO anti-Parkinson's disease therapies from the following adjunctive pharmacotherapy classes (trial must be from two different classes):

- Dopamine agonists (e.g., pramipexole, ropinirole)
- Catechol-O-methyl transferase (COMT) inhibitors (e.g., entacapone)
- Monoamine oxidase (MAO) B inhibitors (e.g., selegiline)

Product Name: Inbrija	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to Inbrija therapy

**AND**

2 - Patient will continue to receive treatment with a carbidopa/levodopa-containing medication

**2 . Revision History**

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

## Infliximab Products

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-146015
<b>Guideline Name</b>	Infliximab Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

#### Guideline Note:

Effective Date:	5/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active RA	

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a minimum duration of a 3-month trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:

- methotrexate
- leflunomide
- sulfasalazine

**AND**

**4** - Used in combination with methotrexate

**AND**

**5** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Documentation of positive clinical response to therapy as evidenced by at least one of the following:	

- Reduction in the total active (swollen and tender) joint count from baseline
- Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of active PsA</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <ul style="list-style-type: none"> <li>• Actively inflamed joints</li> <li>• Dactylitis</li> <li>• Enthesitis</li> <li>• Axial disease</li> <li>• Active skin and/or nail involvement</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Dermatologist</li> <li>• Rheumatologist</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and</p>	

failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis

Diagnosis Psoriatic Arthritis (PsA)

Approval Length 12 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy as evidenced by at least one of the following:

- Reduction in the total active (swollen and tender) joint count from baseline
- Improvement in symptoms (e.g., pain, stiffness, pruritus, inflammation) from baseline
- Reduction in the body surface area (BSA) involvement from baseline

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis

Diagnosis Plaque Psoriasis (PsO)

Approval Length 6 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of chronic severe (i.e., extensive and/or disabling) plaque psoriasis

**AND**

2 - One of the following:

- Greater than or equal to 3% body surface area involvement
- Severe scalp psoriasis
- Palmoplantar (i.e., palms, soles), facial, or genital involvement

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies:

- corticosteroids (e.g., betamethasone, clobetasol)
- vitamin D analogs (e.g., calcitriol, calcipotriene)
- tazarotene
- calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- anthralin
- coal tar

**AND**

**4** - Prescribed by or in consultation with a dermatologist

**AND**

**5** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)

Product Name:Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to infliximab therapy as evidenced by ONE of the following:

- Reduction the body surface area (BSA) involvement from baseline
- Improvement in symptoms (e.g., pruritus, inflammation) from baseline

Product Name:Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Ankylosing Spondylitis (AS)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of active ankylosing spondylitis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Paid claims or submission of medical records (e.g., chart notes) confirming a minimum duration of one month trial and failure, contraindication, or intolerance to two different NSAIDs (e.g., ibuprofen, naproxen) at maximally tolerated doses</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) *DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)</p>	

Product Name:Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Ankylosing Spondylitis (AS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy as evidenced by improvement from baseline for least one of the following:</p> <ul style="list-style-type: none"> <li>• Disease activity (e.g., pain, fatigue, inflammation, stiffness)</li> <li>• Lab values (erythrocyte sedimentation rate, C-reactive protein level)</li> <li>• Function</li> <li>• Axial status (e.g., lumbar spine motion, chest expansion)</li> <li>• Total active (swollen and tender) joint count</li> </ul>	

Product Name:Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Crohn's Disease (CD) or Fistulizing Crohn's Disease
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of one of the following:</p> <ul style="list-style-type: none"> <li>• Moderately to severely active Crohn's disease</li> <li>• Fistulizing Crohn's disease</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p>	

- Frequent diarrhea and abdominal pain
- At least 10% weight loss
- Complications such as obstruction, fever, abdominal mass
- Abnormal lab values (e.g., C-reactive protein [CRP])
- CD Activity Index (CDAI) greater than 220

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one of the following conventional therapies:

- 6-mercaptopurine
- Azathioprine
- Corticosteroids (e.g., prednisone)
- Methotrexate

**AND**

**5** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)

<b>Product Name:Zymfentra</b>	
Diagnosis	Crohn's Disease (CD)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's disease	

**AND**

**2** - Prescribed by or in consultation with a gastroenterologist

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)

**AND**

**4** - Patient has achieved a clinical response following a minimum of 10 weeks of IV infliximab (Janssen manufacturer)

**AND**

**5** - Provider attests that continued IV administration is not appropriate for the patient (e.g., problems with IV access)

Product Name:Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis, Zymfentra	
Diagnosis	Crohn's Disease (CD) or Fistulizing Crohn's Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy as evidenced by at least one of the following:	

- Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline
- Reversal of high fecal output state

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <ul style="list-style-type: none"> <li>• Greater than 6 stools per day</li> <li>• Frequent blood in the stools</li> <li>• Frequent urgency</li> <li>• Presence of ulcers</li> <li>• Abnormal lab values (e.g., hemoglobin, ESR, CRP)</li> <li>• Dependent on, or refractory to, corticosteroids</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a gastroenterologist</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one of the following conventional therapies:</p>	

- 6-mercaptopurine
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)
- Azathioprine
- Corticosteroids (e.g., prednisone)

**AND**

**5 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)**

Product Name: Zymfentra	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis</b></p> <p><b>AND</b></p> <p><b>2 - Prescribed by or in consultation with a gastroenterologist</b></p> <p><b>AND</b></p> <p><b>3 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) *DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)</b></p> <p><b>AND</b></p>	

**4** - Patient has achieved a clinical response following a minimum of 10 weeks of IV infliximab (Janssen manufacturer)

**AND**

**5** - Provider attests that continued IV administration is not appropriate for the patient (e.g., problems with IV access)

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis, Zymfentra

Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to therapy as evidenced by at least one of the following:

- Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline
- Reversal of high fecal output state

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis

Diagnosis	Sarcoidosis [Off-label] [12-15]
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of sarcoidosis

**AND**

2 - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Dermatologist
- Ophthalmologist

**AND**

3 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one corticosteroid (e.g., prednisone)

**AND**

4 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one immunosuppressant (e.g., methotrexate, cyclophosphamide, or azathioprine)

**AND**

5 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure or intolerance to Infliximab (Janssen manufacturer) \*DOES NOT APPLY TO REQUESTS FOR INFLIXIMAB (JANSSEN MANUFACTURER)

Product Name: Avsola, Inflectra, Infliximab (Janssen manufacturer), Remicade, Renflexis	
Diagnosis	Sarcoidosis [Off-label] [12-15]
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to infliximab therapy

## 2 . Revision History

Date	Notes
4/23/2024	Added Zymfentra as NP target

Ingrezza (valbenazine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-155390
<b>Guideline Name</b>	Ingrezza (valbenazine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Ingrezza	
Diagnosis	Moderate to Severe Tardive Dyskinesia
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderate to severe tardive dyskinesia (TD) secondary to a centrally acting dopamine receptor blocking agent (DRBA)	

**AND**

**2** - Prescribed by or in consultation with a psychiatrist or neurologist

**AND**

**3** - Patient is 18 years of age or older

**AND**

**4** - Patient has an Abnormal Involuntary Movement Scale (AIMS) score of 3 or 4 on any one of the AIMS items 1 through 9

**AND**

**5** - Ingrezza is not prescribed concurrently with Austedo or tetrabenazine

**AND**

**6** - Dose does not exceed 80 mg per day

Notes

\*NOTE: Patients will be approved for ONE strength of Ingrezza ONLY : Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Ingrezza products. End-date/Retire all other active PA's for Ingrezza products.

Product Name:Ingrezza	
Diagnosis	Moderate to Severe Tardive Dyskinesia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient is responding positively to therapy as evidenced by a reduction in the baseline AIMS score in any one of the AIMS items 1 through 9

**AND**

2 - Ingrezza is not prescribed concurrently with Austedo or tetrabenazine

**AND**

3 - Dose does not exceed 80 mg per day

Notes

\*NOTE: Patients will be approved for ONE strength of Ingrezza ONLY : Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Ingrezza products. End-date/Retire all other active PA's for Ingrezza products.

Product Name:Ingrezza

Diagnosis Chorea Associated with Huntington's Disease

Approval Length 6 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Diagnosis of chorea in patients with Huntington's disease

**AND**

2 - Prescribed by or in consultation with a neurologist

**AND**

3 - Patient is 18 years of age or older

<b>AND</b>	
4 - Dose does not exceed 80 mg per day	
Notes	*NOTE: Patients will be approved for ONE strength of Ingrezza ONLY : Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Ingrezza products. End-date/Retire all other active PA's for Ingrezza products.

Product Name:Ingrezza	
Diagnosis	Chorea Associated with Huntington's Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Dose does not exceed 80 mg per day.</p>	
Notes	*NOTE: Patients will be approved for ONE strength of Ingrezza ONLY : Approve ALL requests at GPI-14. Confirm there is only 1 active PA on file for ALL Ingrezza products. End-date/Retire all other active PA's for Ingrezza products.

**2 . Revision History**

Date	Notes
9/24/2024	Added note for PA Team, only 1 strength of Ingrezza can be approved.

Inhaled Corticosteroids - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-105180
<b>Guideline Name</b>	Inhaled Corticosteroids - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

### 1 . Criteria

Product Name:Alvesco, Arnuity Ellipta, Asmanex HFA, Qvar Redihaler	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of asthma  <b>AND</b>	

**2** - History of failure, contraindication, intolerance to a majority (not more than 3) of the preferred inhaled corticosteroids:

- Asmanex Twisthaler (mometasone)
- Flovent Diskus (fluticasone)
- Flovent HFA (fluticasone)
- Pulmicort Flexhaler (budesonide)
- budesonide respule (generic)

## **2 . Revision History**

Date	Notes
3/24/2022	Removed Pulmicort and budesonide respules as targets

Injectable Oncology Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-269191
<b>Guideline Name</b>	Injectable Oncology Agents
<b>Formulary</b>	<ul style="list-style-type: none"> <li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li> </ul>

**Guideline Note:**

Effective Date:	6/1/2025
-----------------	----------

### 1 . Criteria

Product Name:*Injectable Oncology Drugs: Abecma, Abraxane, Adcetris, Brand Alimta, generic pemetrexed, Anktiva, Brand Axtle, generic pemetrexed, Bavencio, Belrapzo, Bendeka, Besponsa, Blynicyto, Brand Bicnu, generic carmustine, Bizengri, Brand Bortezomib, Breyanzi, Boruzu, Brand Carmustine, Carvykti, Columvi, cyclophosphamide, Darzalex, Darzalex Faspro, Datroway, Docetaxel, Elahere, Elrexfio, Enhertu, Epkinly, fludarabine, Folutyn, Grafapex, Brand Hycamtin injection, generic topotecan injection, Imdelltra, Imfinzi, Imjudo, Ivra, Jemperli, Keytruda, Kymriah, Kyprolis, Libtayo, Loqtorzi, Lunsumio, methotrexate, Onivyde, Opdivo, Opdivo Qvantig, Paclitaxel, Brand Pemetrexed, Brand Pemfexy, Pemrydi RTU, Polivy, Pralatrexate, Rybrevant, Sarclisa, Synribo, Talvey, Tecartus, Tecentriq, Tecentriq Hybreza, Tecvayli, Temodar IV, Teylute, Tevimbra, Tivdak, Brand Treanda, generic bendamustine, Trodelvy, Vectibix, Brand Velcade, generic bortezomib, Brand Vidaza, generic azacitidine, Vivimusta, Vyloy, Yervoy, Yescarta, Ziihera, Zynyz	
Diagnosis	Cancer Indications
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - The drug is being used as indicated by National Comprehensive Cancer Network (NCCN) guidelines with a Category of Evidence and Consensus of 1, 2A, or 2B

Notes

\*For medical PA, Final approval must be approved by the member's plan. If approved, the transaction will go through POS.

**2 . Revision History**

Date	Notes
5/29/2025	Added Tepylute as target

Inlyta

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99682
<b>Guideline Name</b>	Inlyta
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name:Inlyta	
Diagnosis	Advanced Renal Cell Carcinoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of renal cell cancer	

**AND**

**2** - One of the following:

**2.1** Disease has relapsed

**OR**

**2.2** Diagnosis of Stage IV disease

Product Name: Inlyta	
Diagnosis	Thyroid Carcinoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - ONE of the following diagnosis:</p> <ul style="list-style-type: none"><li>• Follicular Carcinoma</li><li>• Hürthle Cell Carcinoma</li><li>• Papillary Carcinoma</li></ul> <p><b>AND</b></p> <p><b>2</b> - ONE of the following:</p> <ul style="list-style-type: none"><li>• Unresectable recurrent</li><li>• Persistent locoregional disease</li><li>• Metastatic disease</li></ul> <p><b>AND</b></p>	

**3** - Disease is refractory to radioactive iodine treatment

Product Name: Inlyta

Diagnosis	Advanced Renal Cell Carcinoma, Thyroid Carcinoma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Patient does not show evidence of progressive disease while on Inlyta therapy

Product Name: Inlyta

Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Inlyta will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.

Product Name: Inlyta

Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to Inlyta therapy

**2 . Revision History**

Date	Notes
4/8/2021	7/1 Implementation

Insulins, Concentrated- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-154074
<b>Guideline Name</b>	Insulins, Concentrated- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Humulin R U-500 vial and kwikpen	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, intolerance, or contraindication to ALL of the following: <ul style="list-style-type: none"><li>Insulin aspart or insulin lispro</li><li>Lantus</li></ul>	

**OR**

**2** - There is a reason or special circumstance the patient needs to use a concentrated insulin product

## **2 . Revision History**

Date	Notes
9/24/2024	Updated embedded step requirement

Inzirqo (hydrochlorothiazide) oral suspension

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-226190
<b>Guideline Name</b>	Inzirqo (hydrochlorothiazide) oral suspension
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name: Inzirqo	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Patient is under 10 years of age</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** One of the following:

**1.2.1** Trial and failure, or intolerance to hydrochlorothiazide capsules/tablets

**OR**

**1.2.2** Patient is unable to swallow oral capsules/tablets

## **2 . Revision History**

Date	Notes
3/26/2025	New program

Iqirvo (elafibranor)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152668
<b>Guideline Name</b>	Iqirvo (elafibranor)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Iqirvo	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of primary biliary cholangitis (PBC) (also known as primary biliary cirrhosis)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting one of the following:

**2.1** Both of the following:

**2.1.1** Patient has failed to achieve an alkaline phosphatase (ALP) level of less than 1.67 times the upper limit of normal (ULN) after at least 12 consecutive months of treatment with ursodeoxycholic acid (UDCA) (e.g., Urso, Urso Forte, ursodiol)

**AND**

**2.1.2** Used in combination with ursodeoxycholic acid (UDCA)

**OR**

**2.2** History of contraindication or intolerance to ursodeoxycholic acid (UDCA)

**AND**

**3** - Requested drug will not be used in combination with Ocaliva (obeticholic acid)

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Hepatologist
- Gastroenterologist

Product Name: Iqirvo	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., ALP level less than 1.67 times ULN, total bilirubin less than or equal to ULN, ALP decrease greater than or equal to 15% from baseline)

**AND**

2 - Requested drug will not be used in combination with Ocaliva (obeticholic acid)

**2 . Revision History**

Date	Notes
8/27/2024	New Program

Iron Chelators

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-104870
<b>Guideline Name</b>	Iron Chelators
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	3/17/2022
-----------------	-----------

### 1 . Criteria

Product Name:Brand Exjade, Brand Jadenu, generic deferasirox	
Diagnosis	Chronic Iron Overload due to Blood Transfusion
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of chronic iron overload (e.g., sickle cell anemia, thalassemia, etc.) due to blood transfusion	

Product Name:Brand Exjade, Brand Jadenu, generic deferasirox	
Diagnosis	Chronic Iron Overload due to Blood Transfusion
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

Product Name:Brand Ferriprox, generic deferiprone	
Diagnosis	Chronic Iron Overload due to Blood Transfusion
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - BOTH of the following</p> <p>1.1 Diagnosis of transfusional iron overload due to thalassemia syndromes</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Current chelation therapy is inadequate [e.g., Desferal (deferoxamine), Exjade (deferasirox)]</p>	

Product Name:Brand Ferriprox, generic deferiprone	
Diagnosis	Chronic Iron Overload due to Blood Transfusion
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

Product Name: Brand Exjade, Brand Jadenu, generic deferasirox

Diagnosis	Chronic Iron Overload in Non-Transfusion Dependent Thalassemia Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - ALL of the following:

1.1 Diagnosis of chronic iron overload in non-transfusion dependent thalassemia syndrome

**AND**

1.2 Patient has liver iron (Fe) concentration (LIC) levels consistently greater than or equal to 5 mg Fe per gram of dry weight prior to initiation of treatment with Exjade or Jadenu

**AND**

1.3 Patient has serum ferritin levels consistently greater than 300 micrograms per liter prior to initiation of treatment with Exjade or Jadenu

Product Name: Brand Exjade, Brand Jadenu, generic deferasirox

Diagnosis	Chronic Iron Overload in Non-Transfusion Dependent Thalassemia Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
3/16/2022	Added new generic deferiprone tabs

Irritable Bowel Syndrome-Diarrhea

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99468
<b>Guideline Name</b>	Irritable Bowel Syndrome-Diarrhea
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Lotronex, generic alosetron	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of severe diarrhea-predominant irritable bowel syndrome (IBS)  <b>AND</b>	

2 - Symptoms for at least 6 months

**AND**

3 - Patient was female at birth

**AND**

4 - Age greater than or equal to 18 years

**AND**

5 - History of failure, contraindication, or intolerance to TWO of the following:

- Antispasmodic agent (e.g. dicyclomine)
- Antidiarrheal agents (e.g. loperamide)
- Tricyclic antidepressant (e.g. amitriptyline)

Product Name: Brand Lotronex, generic alosetron

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to Lotronex therapy

Product Name: Viberzi

Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of irritable bowel syndrome with diarrhea (IBS-D)

**AND**

2 - History of failure, contraindication, or intolerance to TWO of the following:

- Antispasmodic agent (e.g. dicyclomine)
- Antidiarrheal agents (e.g. loperamide)
- Tricyclic antidepressant (e.g. amitriptyline)

Product Name:Viberzi	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Viberzi therapy	

**2 . Revision History**

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Isotretinoin - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269200
<b>Guideline Name</b>	Isotretinoin - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Absorica, Absorica LD, Accutane, Amnesteem, Claravis, generic isotretinoin, Myorisan, Zenatane	
Diagnosis	Oncology Uses (Off Label)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Used for oncology indication meeting National Comprehensive Cancer Network (NCCN) with a Category of Evidence and Consensus of 1, 2A, or 2B. or from ONE of the following appropriate compendia of current literature: American Hospital Formulary Service Drug Information, Thomson Micromedex DrugDex, or Clinical Pharmacology	

Product Name: Brand Absorica, Absorica LD, Accutane, Amnesteem, Claravis, generic isotretinoin, Myorisan, Zenatane

Approval Length | 5 month(s)

Therapy Stage | Initial Authorization

Guideline Type | Prior Authorization

**Approval Criteria**

1 - ONE of the following:

1.1 Diagnosis of severe recalcitrant nodular acne unresponsive to conventional therapy

**OR**

1.2 Diagnosis of treatment resistant acne

**AND**

2 - History of failure, contraindication, or intolerance to an adequate trial on TWO of the following conventional therapy regimens:

- Topical retinoid or retinoid-like agent [eg, Retin-A/Retin-A Micro (tretinoin)]
- Oral antibiotic [eg, Ery-Tab (erythromycin), Biaxin (clarithromycin), Minocin (minocycline)]
- Topical antibiotic with or without benzoyl peroxide [eg, Cleocin-T (clindamycin), erythromycin, BenzaClin (benzoyl peroxide/clindamycin), Benzamycin (benzoyl peroxide/erythromycin)]

**AND**

3 - If the request is for a non-preferred medication, there must be a reason or special circumstance that the patient must be treated with a non-preferred medication (see table in Background section)

Product Name: Brand Absorica, Absorica LD, Accutane, Amnesteem, Claravis, generic isotretinoin, Myorisan, Zenatane	
Diagnosis	Persistent or Recurring Acne After 2 Months Off Therapy
Approval Length	5 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - After greater than or equal to 2 months OFF therapy, persistent or recurring severe recalcitrant nodular acne is still present</p>	
Notes	Authorization will be given only by clinical pharmacist review for up to 5 months.

Product Name: Brand Absorica, Absorica LD, Accutane, Amnesteem, Claravis, generic isotretinoin, Myorisan, Zenatane	
Diagnosis	Dose Titration
Approval Length	1 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Confirmation that the cumulative dose is less than 150 mg/kg (there is little therapeutic benefit to be gained by increasing the cumulative dose beyond 150 mg/kg)*</p>	
Notes	Authorization will be given only by clinical pharmacist review for 1 month to allow for titration up to the target dose *See background section for dosing regimens

## 2 . Background

<b>Benefit/Coverage/Program Information</b>
---

**Formulary****Preferred Agents:**

Accutane, Myorisan (isotretinoin), Claravis (isotretinoin), Amnesteem (isotretinoin), Zenatane (isotretinoin), generic isotretinoin

**Non-Preferred Agents:**

Absorica (isotretinoin)

Absorica LD (isotretinoin)

**Dosing by Body Weight (based on administration with food):**

Body Weight		Daily Dose		
Kg	Lbs	0.5 mg/kg/day	1 mg/kg/day	2 mg/kg/day
40	88	20	40	80
50	110	25	50	100
60	132	30	60	120
70	154	35	70	140
80	176	40	80	160
90	198	45	90	180
100	220	50	100	200

--

### 3 . Revision History

Date	Notes
5/29/2025	Added GPI for Amnesteem 30mg

Isturisa (osilodrostat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269196
<b>Guideline Name</b>	Isturisa (osilodrostat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Isturisa	
Diagnosis	Cushing's Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Cushing's syndrome	

**AND**

**2** - Used for treatment of endogenous hypercortisolemia

**AND**

**3** - One of the following:

- Patient is not a candidate for surgery (e.g., adrenalectomy, transsphenoidal surgery)
- Surgery has not been curative for the patient

Product Name:Isturisa	
Diagnosis	Cushing's Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy (e.g., a clinically meaningful reduction in 24-hour urinary free cortisol levels, improvement in signs or symptoms of the disease)	

Product Name:Isturisa	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Isturisa will be approved for uses supported by The National Comprehensive Cancer	

Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.

## 2 . Revision History

Date	Notes
5/29/2025	Updated Cushing's syndrome criteria due to revised indication, removed reauth for NCCN approved indications section (NCCN auth is 12 months).

Izervay (avacincaptad pegol)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135329
<b>Guideline Name</b>	Izervay (avacincaptad pegol)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Izervay	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of geographic atrophy (GA) secondary to age-related macular degeneration (AMD) as confirmed by one of the following: <ul style="list-style-type: none"><li>Fundus photography (e.g. fundus autofluorescence [FAF])</li></ul>	

- Optical coherence tomography (OCT)
- Fluorescein angiography

**AND**

**2** - GA is not secondary to any other conditions (e.g., Stargardt disease, cone rod dystrophy, toxic maculopathies)

**AND**

**3** - Prescribed by or in consultation with an ophthalmologist experienced in the treatment of retinal diseases

Product Name: IZERVAY	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., reduction in growth rate of GA lesion)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient has not exceeded a total of 12 months treatment</p>	

## 2 . Revision History

Date	Notes
10/23/2023	New program

Jesduvroq (daprodustat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-143522
<b>Guideline Name</b>	Jesduvroq (daprodustat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Jesduvroq	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of chronic kidney disease (CKD)  <b>AND</b>	

2 - Patient has been on dialysis for at least 4 months

**AND**

3 - Adequate iron stores confirmed by both of the following:

- Patient's ferritin level is greater than 100mcg/L
- Patient's transferrin saturation (TSAT) is greater than 20%

**AND**

4 - Hemoglobin level less than 11 g/dL

**AND**

5 - Trial and failure, contraindication or intolerance to one of the following:

- Epogen
- Procrit
- Retacrit

**AND**

6 - Prescribed by or in consultation with one of the following:

- hematologist
- nephrologist

**AND**

7 - Patient is not on concurrent treatment with an erythropoietin stimulating agent [ESA] (e.g., Aranesp, Epogen, Procrit)

Product Name: Jesduvroq	
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy (e.g., increase in hemoglobin)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Hemoglobin level does not exceed 12g/dL</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Adequate iron stores confirmed by both of the following:</p> <ul style="list-style-type: none"> <li>• Patient's ferritin level is greater than 100mcg/L</li> <li>• Patient's transferrin saturation (TSAT) is greater than 20%</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Patient is not on concurrent treatment with an erythropoietin stimulating agent [ESA] (e.g., Aranesp, Epogen, Procrit)</p>	

## 2 . Revision History

Date	Notes
2/28/2024	New program

Joenja (leniolisib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-127086
<b>Guideline Name</b>	Joenja (leniolisib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Joenja	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, laboratory values) documenting ALL of the following:  1.1 Diagnosis of activated phosphoinositide 3-kinase delta syndrome (APDS)	

**AND**

**1.2** Molecular genetic testing confirms mutations in the PIK3CD or PIK3R1 gene

**AND**

**1.3** Both of the following:

- Presence of nodal and/or extranodal proliferation (e.g., lymphadenopathy, splenomegaly, hepatomegaly)
- Presence of other clinical findings and manifestations consistent with APDS (e.g., recurrent sino-pulmonary infections, bronchiectasis, enteropathy)

**AND**

**1.4** Trial and failure, contraindication, or intolerance to at least one standard of care treatment for APDS (e.g., Immunoglobulin replacement therapy, antimicrobial prophylaxis [e.g., azithromycin, bactrim], rituximab, tacrolimus, etc.)

**AND**

**2** - Patient is 12 years of age or older

**AND**

**3** - Patient weighs greater than or equal to 45kg

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Hematologist
- Immunologist

Product Name:Joenja	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., reduced lymph node size, increased naïve B-cell percentage, decreased severity or frequency of infections/hospitalizations)</p>	

## 2 . Revision History

Date	Notes
6/26/2023	New program

Juxtapid - AZ

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99791
<b>Guideline Name</b>	Juxtapid - AZ
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Juxtapid	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of homozygous familial hypercholesterolemia (HoFH) as confirmed by BOTH of the following:*	
1.1 ONE of the following:	

- Pre-treatment low density lipoprotein cholesterol (LDL-C) greater than 500 milligrams per deciliter
- Treated LDL-C greater than 300 milligrams per deciliter

**AND**

**1.2 ONE** of the following:

- Xanthoma before 10 years of age
- Evidence of heterozygous familial hypercholesterolemia (HeFH) in both parents

**AND**

**2** - Used as an adjunct to a low-fat diet and exercise

**AND**

**3** - Patient is receiving other lipid-lowering therapy (e.g., statin, ezetimibe, LDL apheresis)

**AND**

**4** - Prescribed by **ONE** of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

**AND**

**5** - Patient has tried, failed or intolerant to Repatha and Praluent

**AND**

**6** - Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor

Notes	Results of prior genetic testing can be submitted as confirmation of diagnosis of HoFH.
-------	---

Product Name:Juxtapid	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is continuing a low-fat diet and exercise regimen</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient continues to receive other lipid-lowering therapy (e.g., statin, low density lipoprotein [LDL] apheresis)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Submission of medical records (e.g. chart notes, laboratory values) documenting low density lipoprotein cholesterol (LDL-C) reduction while on Juxtapid therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by ONE of the following:</p> <ul style="list-style-type: none"> <li>• Cardiologist</li> <li>• Endocrinologist</li> <li>• Lipid specialist</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>5 - Not used in combination with a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor</p>	

**2 . Revision History**

Date	Notes
7/13/2021	Arizona Medicaid 7.1 Implementation

Jynarque

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-100644
<b>Guideline Name</b>	Jynarque
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name: Jynarque, Jynarque Pak	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of autosomal dominant polycystic kidney disease (ADPKD)	

Product Name:Jynarque, Jynarque Pak	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Jynarque therapy</p>	

## 2 . Revision History

Date	Notes
12/16/2021	Added new Jynarque GPs

Kalydeco (ivacaftor)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135314
<b>Guideline Name</b>	Kalydeco (ivacaftor)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

### 1 . Criteria

Product Name:Kalydeco, Kalydeco packet	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystic fibrosis (CF)  <b>AND</b>	

**2** - Submission of laboratory results confirming that patient has ONE of the mutations in the cystic fibrosis transmembrane conductance regulator (CFTR) gene listed in the table in the Background section:

**AND**

**3** - Prescribed by, or in consultation with, a specialist affiliated with a CF care center

Product Name:Kalydeco, Kalydeco packet

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Provider attests that the patient has achieved a clinically meaningful response while on Kalydeco therapy to ONE of the following:

- Lung function as demonstrated by percent predicted expiratory volume in 1 second (ppFEV1)
- Body mass index (BMI)
- Pulmonary exacerbations
- Quality of life as demonstrated by Cystic Fibrosis Questionnaire-Revised (CFQ-R) respiratory domain score

**AND**

**2** - Prescribed by, or in consultation with, specialist affiliated with a cystic fibrosis (CF) care center

**2 . Background**

Benefit/Coverage/Program Information

**CFTR Gene Mutations that are Responsive to Kalydeco**

**List of CFTR Gene Mutations that Produce CFTR Protein and are Responsive to KALYDECO**

<i>711+3A→G *</i>	<i>F311del</i>	<i>I148T</i>	<i>R75Q</i>	<i>S589N</i>
<i>2789+5G→A *</i>	<i>F311L</i>	<i>I175V</i>	<i>R117C *</i>	<i>S737F</i>
<i>3272-26A→G *</i>	<i>F508C</i>	<i>I807M</i>	<i>R117G</i>	<i>S945L *</i>
<i>3849+10kbC→T *</i>	<i>F508C;S1251N †</i>	<i>I1027T</i>	<i>R117H *</i>	<i>S977F *</i>
<i>A120T</i>	<i>F1052V</i>	<i>I1139V</i>	<i>R117L</i>	<i>S1159F</i>
<i>A234D</i>	<i>F1074L</i>	<i>K1060T</i>	<i>R117P</i>	<i>S1159P</i>
<i>A349V</i>	<i>G178E</i>	<i>L206W *</i>	<i>R170H</i>	<i>S1251N *</i>
<i>A455E *</i>	<i>G178R *</i>	<i>L320V</i>	<i>R347H *</i>	<i>S1255P *</i>
<i>A1067T</i>	<i>G194R</i>	<i>L967S</i>	<i>R347L</i>	<i>T338I</i>
<i>D110E</i>	<i>G314E</i>	<i>L997F</i>	<i>R352Q *</i>	<i>T1053I</i>
<i>D110H</i>	<i>G551D *</i>	<i>L1480P</i>	<i>R553Q</i>	<i>V232D</i>
<i>D192G</i>	<i>G551S *</i>	<i>M152V</i>	<i>R668C</i>	<i>V562I</i>
<i>D579G *</i>	<i>G576A</i>	<i>M952I</i>	<i>R792G</i>	<i>V754M</i>
<i>D924N</i>	<i>G970D</i>	<i>M952T</i>	<i>R933G</i>	<i>V1293G</i>
<i>D1152H *</i>	<i>G1069R</i>	<i>P67L *</i>	<i>R1070Q</i>	<i>W1282R</i>
<i>D1270N</i>	<i>G1244E *</i>	<i>Q237E</i>	<i>R1070W *</i>	<i>Y1014C</i>
<i>E56K</i>	<i>G1249R</i>	<i>Q237H</i>	<i>R1162L</i>	<i>Y1032C</i>
<i>E193K</i>	<i>G1349D *</i>	<i>Q359R</i>	<i>R1283M</i>	
<i>E822K</i>	<i>H939R</i>	<i>Q1291R</i>	<i>S549N *</i>	
<i>E831X *</i>	<i>H1375P</i>	<i>R74W</i>	<i>S549R *</i>	

\* Clinical data exist for these mutations.

† Complex/compound mutations where a single allele of the CFTR gene has multiple mutations, these exist independent of the presence of mutations on the other allele.

### 3 . Revision History

Date	Notes
10/23/2023	Added GPI for 5.8 mg packs

Katerzia, Norliqva (amlodipine oral solution)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-137600
<b>Guideline Name</b>	Katerzia, Norliqva (amlodipine oral solution)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Katerzia, Norliqva	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Patient is 8 years of age or younger</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** Both of the following:

**1.2.1** Requested medication is being used for one of the following diagnoses:

- Hypertension
- Chronic stable angina
- Confirmed or suspected vasoplastic angina
- Angiographically documented Coronary Artery Disease (CAD)

**AND**

**1.2.2** One of the following:

**1.2.2.1** Trial and failure, contraindication, or intolerance to generic amlodipine tablets (verified via paid pharmacy claims or submitted chart notes)

**OR**

**1.2.2.2** Patient is unable to swallow oral tablets/capsules

**AND**

**2 - For Norliqva requests: trial and failure, contraindication, or intolerance to Katerzia (verified via paid pharmacy claims or submitted chart notes) APPLIES TO NORLIQVA REQUESTS ONLY**

## **2 . Revision History**

Date	Notes
12/11/2023	Added step through Katerzia for Norliqva (now NP).

Kepivance (palifermin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-136964
<b>Guideline Name</b>	Kepivance (palifermin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Kepivance	
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting all of the following:  1.1 Medication will be used to prevent or treat severe (WHO Grade 3 or higher) oral mucositis	

**AND**

**1.2** Inadequate response to an oral mouthwash formulated with diphenhydramine/antacid and lidocaine (e.g., magic or miracle mouthwash)

**AND**

**2** - Prescribed by a hematologist or oncologist

## **2 . Revision History**

Date	Notes
12/1/2023	New program

Kerendia (finerenone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-126131
<b>Guideline Name</b>	Kerendia (finerenone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Kerendia	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of chronic kidney disease (CKD) associated with type 2 diabetes (T2D)  <b>AND</b>	

2 - Urine albumin-to-creatinine ratio (UACR) greater than or equal to 30 mg/g

**AND**

3 - Estimated glomerular filtration rate (eGFR) greater than or equal to 25 mL/min/1.73 m<sup>2</sup>

**AND**

4 - Serum potassium level less than or equal to 5.0 mEq/L prior to initiating treatment

**AND**

5 - One of the following:

5.1 Minimum 30-day supply trial of a maximally tolerated dose and will continue therapy with one of the following [2]:

- Generic angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril)
- Generic angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan)

**OR**

5.2 Patient has a contraindication or intolerance to ACE inhibitors and ARBs

Product Name:Kerendia	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

**AND**

**2 - One of the following:**

**2.1 Patient continues to be on a maximally tolerated dose of ACE inhibitor or ARB**

**OR**

**2.2 Patient has a contraindication or intolerance to ACE inhibitors and ARBs**

## **2 . Revision History**

Date	Notes
5/30/2023	Updated initial auth criteria

Keveyis

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99621
<b>Guideline Name</b>	Keveyis
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Keveyis	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 Diagnosis of primary hyperkalemic periodic paralysis or related variant	

**OR**

**1.2** Diagnosis of primary hypokalemic periodic paralysis or related variant

Product Name:Keveyis	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Keveyis therapy	

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Kevzara (sarilumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-329255
<b>Guideline Name</b>	Kevzara (sarilumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Kevzara	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g. chart notes) or verification of paid claims documenting ALL of the following:  1.1 Diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**1.2** History of failure to a 3 month trial of ONE non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Oencia (abatacept)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred tocilizumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Kevzara	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by at least one of the following	
<ul style="list-style-type: none"><li>• Reduction in the total active (swollen and tender) joint count from baseline</li><li>• Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline</li></ul>	

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Kevzara

Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
-----------	--

Approval Length	6 month(s)
-----------------	------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of active polyarticular juvenile idiopathic arthritis (PJIA)

**AND**

**2** - Patient weighs at least 63 kg

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) confirming a minimum duration of a 6-week trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:

- leflunomide
- methotrexate

**AND**

**5** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Orencia (abatacept)
- Xeljanz (tofacitinib) oral tablet
- A preferred tocilizumab biosimilar

Product Name:Kevzara	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
 <b>Approval Criteria</b>  <b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by at least one of the following: <ul style="list-style-type: none"><li>• Reduction in the total active (swollen and tender) joint count from baseline</li><li>• Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline</li></ul> <p style="text-align: center;"><b>AND</b></p> <b>2</b> - Prescribed by or in consultation with a rheumatologist	

Product Name:Kevzara	
Diagnosis	Polymyalgia Rheumatica (PMR)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of polymyalgia rheumatica (PMR)

**AND**

2 - One of the following:

2.1 Patient has had an inadequate response to corticosteroids (e.g., prednisone)

**OR**

2.2 Patient cannot tolerate tapering of corticosteroids (e.g., prednisone)

**AND**

3 - Prescribed by or in consultation with a rheumatologist

Product Name:Kevzara	
Diagnosis	Polymyalgia Rheumatica (PMR)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by at least one of the following:	
<ul style="list-style-type: none"><li>Improvement in symptoms (e.g., pain, stiffness) or lab values (e.g., C-reactive protein) from baseline</li></ul>	

- Reduced need for corticosteroids (e.g., prednisone)

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

## **2 . Revision History**

Date	Notes
7/16/2025	Updated preferred/embedded steps for RA and PJIA, removed COT f or RA. PJIA: removed "SC" from Orencia

Kimmtrak (tebentafusp-tebn)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-104978
<b>Guideline Name</b>	Kimmtrak (tebentafusp-tebn)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Kimmtrak	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of uveal melanoma  <b>AND</b>	

2 - Disease is unresectable or metastatic

**AND**

3 - Patient is HLA-A\*02:01 genotype positive as determined by a high-resolution genotyping test [2]

**AND**

4 - Prescribed by or in consultation with an oncologist

Product Name:Kimmtrak	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting patient does not show evidence of progressive disease while on therapy	

## 2 . Revision History

Date	Notes
3/22/2022	New Program mirrors ORx with Submission of Records added to initial and reauth

Kineret (anakinra)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300290
<b>Guideline Name</b>	Kineret (anakinra)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Kineret	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**1.2** History of failure to a 3 month trial of one non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Oencia (abatacept)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred tocilizumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Kineret	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a rheumatologist	

Product Name:Kineret	
Diagnosis	Neonatal-Onset Multisystem Inflammatory Disease (NOMID)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming the diagnosis of neonatal-onset multisystem inflammatory disease (NOMID)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Diagnosis of NOMID has been confirmed by one of the following:</p> <p>2.1 NLRP-3 (nucleotide-binding domain, leucine rich family (NLR), pyrin domain containing 3-gene (also known as Cold-Induced Auto-inflammatory Syndrome-1 [CIAS1]) mutation</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Both of the following:</p> <p>2.2.1 Two of the following clinical symptoms:</p> <ul style="list-style-type: none"> <li>• Urticaria-like rash</li> <li>• Cold/stress triggered episodes</li> <li>• Sensorineural hearing loss</li> <li>• Musculoskeletal symptoms (e.g., arthralgia, arthritis, myalgia)</li> <li>• Chronic aseptic meningitis</li> <li>• Skeletal abnormalities (e.g., epiphyseal overgrowth, frontal bossing)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2.2.2 Elevated acute phase reactants (e.g., erythrocyte sedimentation rate [ESR], C-reactive protein [CRP], serum amyloid A [SAA])</p>	

**AND**

**3** - Prescribed by or in consultation with one of the following

- Allergist/Immunologist
- Rheumatologist
- Pediatrician

Product Name:Kineret	
Diagnosis	Neonatal-Onset Multisystem Inflammatory Disease (NOMID)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

Product Name:Kineret	
Diagnosis	Deficiency of Interleukin-1 Receptor Antagonist (DIRA)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming the diagnosis of deficiency of interleukin-1 receptor antagonist (DIRA)	

Product Name:Kineret	
Diagnosis	Systemic Juvenile Idiopathic Arthritis (SJIA)

Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming the diagnosis of active systemic juvenile idiopathic arthritis</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a rheumatologist</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Paid claims or submission of medical records (e.g., chart notes) confirming trial and failure, contraindication, or intolerance to ONE of the following:</p> <ul style="list-style-type: none"> <li>• Nonsteroidal anti-inflammatory drug (NSAID) (e.g., Motrin [ibuprofen], Naprosyn [naproxen])</li> <li>• Systemic glucocorticoid (e.g., prednisone)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Paid claims or submission of medical records (e.g., chart notes) confirming trial and failure, contraindication, or intolerance to a preferred tocilizumab biosimilar</p>	

Product Name:Kineret	
Diagnosis	Systemic Juvenile Idiopathic Arthritis (SJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with a rheumatologist

**2 . Revision History**

Date	Notes
7/3/2025	Updated preferred agents/embedded steps and criteria for RA and S JIA.

Korlym

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99622
<b>Guideline Name</b>	Korlym
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Korlym	
Diagnosis	Endogenous Cushing's Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ALL of the following:  1.1 Diagnosis of Endogenous Cushing's Syndrome (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids)	

**AND**

**1.2 ONE of the following:**

- Diagnosis of type 2 diabetes mellitus
- Diagnosis of glucose intolerance

**AND**

**1.3 ONE of the following:**

- Patient has failed surgery
- Patient is not a candidate for surgery

Product Name:Korlym	
Diagnosis	Endogenous Cushing's Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of ONE of the following:	
<ul style="list-style-type: none"><li>• Patient has improved glucose tolerance while on Korlym therapy</li><li>• Patient has stable glucose tolerance while on Korlym therapy</li></ul>	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1



Korsuva (difelikefalin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-107424
<b>Guideline Name</b>	Korsuva (difelikefalin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Korsuva	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting all of the following:  1.1 Diagnosis of chronic kidney disease (CKD)	

**AND**

**1.2** Patient is currently undergoing hemodialysis (HD) at an optimal dialysis dose (e.g., Kt/V greater than or equal to 1.2)

**AND**

**1.3** Patient is experiencing moderate to severe pruritus associated with CKD (CKD-aP)

**AND**

**1.4** Exclusion of other causes of pruritus (e. g., eczema, infections, drug-induced skin dryness)

**AND**

**1.5** Trial and failure, contraindication, or intolerance to ONE topical anti-pruritic treatment:

- emollient cream
- analgesics (e.g., pramoxine lotion, capsaicin)
- corticosteroids (e.g., hydrocortisone, triamcinolone)

**AND**

**1.6** Trial and failure, contraindication, or intolerance to ONE oral treatment\*:

- antihistamine (e.g., diphenhydramine, hydroxyzine, loratadine)
- gabapentin
- pregabalin

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Nephrologist

<ul style="list-style-type: none"> <li>• Dermatologist</li> </ul>	
Notes	*PA may be required

Product Name:Korsuva	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting both of the following:</p> <p>1.1 Patient is currently undergoing hemodialysis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 Documentation of positive clinical response to therapy (e.g., improved quality of life, improved worst itching intensity numerical rating score from baseline)</p>	

## 2 . Revision History

Date	Notes
5/24/2022	New Program

Krystexxa (pegloticase)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117642
<b>Guideline Name</b>	Krystexxa (pegloticase)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Krystexxa	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of gout  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) confirming trial and failure, contraindication, or intolerance to maximum recommended doses to both of the following conventional therapies:

- Xanthine oxidase inhibitor (i.e., allopurinol, febuxostat)
- Uricosuric agent (e.g., probenecid)

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting one of the following:

- History of at least two gout flares in the previous 12 months
- At least 1 gouty tophus

**AND**

**4** - Prescribed by or in consultation with a rheumatologist or nephrologist

Product Name:Krystexxa	
Approval Length	12 Months [B]
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g, chart notes) documenting positive clinical response to Krystexxa therapy demonstrated by both of the following:	
<ul style="list-style-type: none"><li>• Serum urate level has decreased since initiating therapy</li><li>• Clinical improvement in the signs and symptoms of gout (e.g., decrease in tophi size or frequency of gouty flares per year from baseline or improvement in chronic arthropathy or quality of life)</li></ul>	

## 2 . Revision History

Date	Notes
12/4/2022	New program

Kuvan

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99623
<b>Guideline Name</b>	Kuvan
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Kuvan	
Diagnosis	Phenylketonuria (PKU)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of phenylketonuria (PKU)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

LAMA, LABA - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269199
<b>Guideline Name</b>	LAMA, LABA - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Anoro Ellipta, Brand Umeclidinium-vilanterol, Bevespi, Stiolto	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of chronic obstructive pulmonary disease (COPD)  <b>AND</b>  2 - One of the following:	

**2.1** History of failure, contraindication, or intolerance to treatment with a 30 day trial of a long-acting beta-agonist (e.g. Foradil, Serevent, Striverdi, Arcapta)

**OR**

**2.2** History of failure, contraindication, or intolerance to treatment with a 30 day trial of an orally inhaled anticholinergic agent (e.g. Spiriva, Atrovent, Combivent, Tudorza)

**AND**

**3** - For Bevespi requests ONLY: history of failure, contraindication, or intolerance to treatment with a 30 day trial of both of the following Preferred drugs:

- Brand Anoro Ellipta
- Stiolto Respimat

**AND**

**4** - For Umeclidinium-vilanterol requests ONLY, history of failure or intolerance to Brand Anoro Ellipta

## 2 . Revision History

Date	Notes
5/29/2025	Added umeclidinium-vilanterol as NP target, with step through Brand Anoro Ellipta

Lamzede (velmanase alfa-tycv)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-125941
<b>Guideline Name</b>	Lamzede (velmanase alfa-tycv)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Lamzede	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of alpha-mannosidosis  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) confirming diagnosis by one of the following:

- Deficiency in alpha-mannosidase enzyme activity as measured in fibroblasts or leukocytes
- Molecular genetic testing confirms mutations in the MAN2B1 gene

**AND**

**3** - Treatment is only for non-central nervous system disease manifestations (e.g., large head, prominent forehead, protruding jaw, skeletal abnormalities)

Product Name:Lamzede	
Approval Length	24 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming patient does not show evidence of progressive disease while on therapy as evidenced by one of the following:</p> <ul style="list-style-type: none"> <li>• Reduction in serum oligosaccharide concentration from baseline</li> <li>• Improvement in clinical signs and symptoms from baseline (e.g., 3-minute stair climbing test, 6-minute walking test, pulmonary function, quality of life)</li> </ul>	

## 2 . Revision History

Date	Notes
5/22/2023	New program

Lantidra (donislecel-jujn)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-136956
<b>Guideline Name</b>	Lantidra (donislecel-jujn)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Lantidra	
Approval Length	One Time Approval
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting ALL of the following: <b>1.1</b> Diagnosis of Type 1 diabetes	

**AND**

**1.2** Patient is insulin dependent

**AND**

**1.3** Patient is unable to approach target HbA1c because of current repeated episodes of severe hypoglycemia despite intensive diabetes management and education

**AND**

**1.4** Patient has reduced awareness of hypoglycemia, as defined by the absence of adequate autonomic symptoms at glucose levels of less than 54 mg/dL

**AND**

**1.5** Patient has had at least one episode of severe hypoglycemia in the past 3 years with both of the following:

**1.5.1** Patient required assistance of another person

**AND**

**1.5.2** One of the following:

**1.5.2.1** Symptoms were associated with a blood glucose level less than 50 mg/dL

**OR**

**1.5.2.2** Prompt recovery after oral carbohydrate, intravenous glucose, or glucagon administration

**AND**

**1.6** Patient will be on concomitant immunosuppression (e.g., daclizumab, sirolimus, tacrolimus, etanercept, mycophenolate mofetil, etc.)

**AND**

**2** - Prescribed by or in consultation with an endocrinologist

Product Name:Lantidra	
Approval Length	One Time Approval
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting that patient has not achieved independence from exogenous insulin within one year of infusion or within one year after losing independence from exogenous insulin after previous infusion</p> <p><b>AND</b></p> <p><b>2</b> - Patient has not had more than three infusions of Lantidra in their lifetime*</p>	
Notes	*There are no data regarding the effectiveness or safety for patients receiving more than three infusions.

## 2 . Revision History

Date	Notes
11/27/2023	New Program

Lenmeldy (atidarsagene autotemcel)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147672
<b>Guideline Name</b>	Lenmeldy (atidarsagene autotemcel)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Lenmeldy	
Approval Length	1 Time Authorization in Lifetime*
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting both of the following: <ul style="list-style-type: none"><li>Diagnosis of metachromatic leukodystrophy (MLD)</li><li>Molecular genetic testing confirms mutation in the arylsulfatase A (ARSA) gene</li></ul>	

**AND**

**2** - Disease is one of the following:

**2.1** Pre-symptomatic late infantile (PSLI) as confirmed by both of the following:

**2.1.1** Disease onset at less than or equal to 30 months of age

**AND**

**2.1.2** One of the following:

- Absence of neurological signs and symptoms of MLD (e.g., peripheral neuropathy, gait difficulties, hypotonia)
- Abnormal reflexes or abnormalities on brain magnetic resonance imaging (MRI) and/or nerve conduction tests not associated with functional impairment (e.g., no tremor, no peripheral ataxia)

**OR**

**2.2** Pre-symptomatic early juvenile (PSEJ) as confirmed by both of the following:

**2.2.1** Disease onset at greater than 30 months and less than 7 years of age

**AND**

**2.2.2** One of the following:

- Absence of neurological signs and symptoms of MLD (e.g., peripheral neuropathy, gait difficulties, hypotonia)
- Abnormal reflexes or abnormalities on brain magnetic resonance imaging (MRI) and/or nerve conduction tests not associated with functional impairment (e.g., no tremor, no peripheral ataxia)

**OR**

**2.3** Early-symptomatic early juvenile (ESEJ) as confirmed by all of the following:

- Disease onset at greater than 30 months and less than 7 years of age

- Gross motor function classification (GMFC)-MLD score less than or equal to 1
- Intelligence quotient (IQ) of greater than or equal to 85

**AND**

**3** - Prescribed by a specialist with expertise in MLD at an authorized treatment center

**AND**

**4** - Both of the following:

- Patient has never received Lenmeldy treatment in their lifetime
- Patient has never received prior hematopoietic stem cell transplant (HSCT)

Notes	*Per prescribing information, Lenmeldy is for one-time, single dose intravenous use only.
-------	---

## 2 . Revision History

Date	Notes
5/23/2024	New program

Leqembi (lecanemab-irmb)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-143555
<b>Guideline Name</b>	Leqembi (lecanemab-irmb)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Leqembi	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Both of the following:  1.1 Based on the National Institute on Aging and the Alzheimer's Association (NIA-AA) criteria, one of the following:	

- Diagnosis of mild cognitive impairment due to Alzheimer's disease
- Diagnosis of probable Alzheimer's disease dementia

**AND**

**1.2** Submission of medical records (e.g., chart notes) confirming all of the following:

- Global Clinical Dementia Rating (CDR) score of 0.5 or 1.0
- CDR Memory Box score of 0.5 or greater
- Mini-Mental State Examination score of 22 or greater

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming the presence of beta-amyloid protein deposition, as evidenced by one of the following:

**2.1** Positive amyloid positron emission tomography (PET) scan

**OR**

**2.2** Both of the following:

- Attestation that the patient does not have access to amyloid PET scanning
- Cerebrospinal fluid (CSF) biomarker or blood testing documents abnormalities suggestive of beta-amyloid accumulation (e.g., A $\beta$ 42 level, A $\beta$ 42:A $\beta$ 40 ratio)

**AND**

**3** - Provider attests that the patient's ApoE e4 carrier status is known prior to initiating treatment and a shared decision-making conversation regarding the results has been completed

**AND**

**4** - Other differential diagnoses (e.g., dementia with Lewy bodies (DLB), frontotemporal dementia (FTD), vascular dementia, pseudodementia due to mood disorder, vitamin B12 deficiency, encephalopathy, etc.) have been ruled out

**AND**

**5** - Both of the following:

- Patient is not currently taking an anticoagulant (e.g., warfarin, dabigatran)
- Patient has no history of intracerebral hemorrhage (e.g., transient ischemic attack [TIA], stroke) within the previous year prior to initiating treatment

**AND**

**6** - Counseling has been provided on the risk of amyloid-related imaging abnormalities (ARIA-E and ARIA-H) and patient and/or caregiver are aware to monitor for headache, dizziness, visual disturbances, nausea, and vomiting

**AND**

**7** - Submission of medical records (e.g., chart notes) confirming a baseline brain magnetic resonance imaging (MRI) has been completed within 12 months prior to initiating treatment

**AND**

**8** - Not used in combination with other A $\beta$  monoclonal antibodies (mAbs) for Alzheimer's Disease (e.g., Aduhelm)

**AND**

**9** - One of the following:

**9.1** Prescribed by a geriatrician or geriatric psychiatrist

**OR**

**9.2** Prescribed by or in consultation with a neurologist

Product Name:Legembi	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - Patient is benefitting from therapy as defined by both of the following:</b></p> <p><b>1.1</b> Based on the National Institute on Aging and the Alzheimer's Association (NIA-AA) criteria, one of the following [2,3]:</p> <ul style="list-style-type: none"> <li>• Patient continues to have a diagnosis of mild cognitive impairment due to Alzheimer's disease</li> <li>• Patient continues to have a diagnosis of probable Alzheimer's disease dementia</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> Submission of medical records (e.g., chart notes) confirming all of the following [4-5]:</p> <ul style="list-style-type: none"> <li>• Global Clinical Dementia Rating (CDR) score of 0.5 or 1.0</li> <li>• CDR Memory Box score of 0.5 or greater</li> <li>• Mini-Mental State Examination score of 22 or greater</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2 - Submission of medical records (e.g., chart notes) confirming follow-up brain magnetic resonance imaging (MRI) has been completed after the initiation of therapy prior to the 5th and 7th infusion treatment to show one of the following:</b></p> <p><b>2.1</b> Both of the following:</p> <ul style="list-style-type: none"> <li>• Less than 10 new incident microhemorrhages</li> <li>• 2 or less focal areas of superficial siderosis</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>2.2</b> If 10 or more new incident microhemorrhages or greater than 2 focal areas of superficial siderosis are present, then both of the following:</p>	

- Patient has been clinically evaluated for ARIA related signs or symptoms (e.g., dizziness, visual disturbances)
- Follow-up MRI demonstrates radiographic stabilization (i.e., no increase in size or number of ARIA-H)

**AND**

**3** - Not used in combination with other A $\beta$  monoclonal antibodies (mAbs) for Alzheimer's Disease (e.g., Aduhelm)

**AND**

**4** - One of the following:

**4.1** Prescribed by a geriatrician or geriatric psychiatrist

**OR**

**4.2** Prescribed by or in consultation with a neurologist

## 2 . Definitions

Definition	Description
ARIA-E	Amyloid related imaging abnormality due to edema/effusion
ARIA-H	Amyloid related imaging abnormality due to micro hemorrhages and hemosiderin deposits

## 3 . Revision History

Date	Notes
2/28/2024	Updated specialist prescriber verbiage



Leqvio (inclisiran)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-129086
<b>Guideline Name</b>	Leqvio (inclisiran)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Leqvio	
Diagnosis	Heterozygous Familial Hypercholesterolemia (HeFH), Atherosclerotic Cardiovascular Disease (ASCVD)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting one of the following diagnoses:	

**1.1** Heterozygous familial hypercholesterolemia (HeFH) as confirmed by one of the following:

**1.1.1** Both of the following: [5]

**1.1.1.1** Untreated/pre-treatment LDL-cholesterol (LDL-C) greater than 190 mg/dL

**AND**

**1.1.1.2** One of the following:

- Family history of myocardial infarction in first-degree relative less than 60 years of age
- Family history of myocardial infarction in second-degree relative less than 50 years of age
- Family history of LDL-C greater than 190 mg/dL in first- or second-degree relative
- Family history of familial hypercholesterolemia in first- or second-degree relative
- Family history of tendinous xanthomata and/or arcus cornealis in first- or second-degree relative

**OR**

**1.1.2** Both of the following: [5]

**1.1.2.1** Untreated/pre-treatment LDL-cholesterol (LDL-C) greater than 190 mg/dL

**AND**

**1.1.2.2** Submission of medical records (e.g., chart notes, laboratory values) documenting one of the following:

- Functional mutation in the LDL receptor, ApoB, or PCSK9 gene
- Tendinous xanthomata
- Arcus cornealis before age 45

**OR**

**1.2** Atherosclerotic cardiovascular disease (ASCVD) as confirmed by one of the following: [2,4]

- Acute coronary syndromes
- History of myocardial infarction

- Stable or unstable angina
- Coronary or other arterial revascularization
- Stroke
- Transient ischemic attack
- Peripheral arterial disease presumed to be of atherosclerotic origin

**AND**

**2** - One of the following: [4]

**2.1** Patient has been receiving at least 12 consecutive weeks of HIGH-INTENSITY statin therapy [i.e., atorvastatin 40-80 mg, rosuvastatin 20-40 mg] and will continue to receive a HIGH-INTENSITY statin at maximally tolerated dose

**OR**

**2.2** Both of the following:

**2.2.1** Patient is unable to tolerate high-intensity statin as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

**AND**

**2.2.2** One of the following:

- Patient has been receiving at least 12 consecutive weeks of MODERATE-INTENSITY statin therapy [i.e., atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin 20-40 mg, pravastatin 40-80 mg, lovastatin 40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 40 mg twice daily, or Livalo (pitavastatin) 2-4 mg] and will continue to receive a MODERATE-INTENSITY statin at maximally tolerated dose
- Patient has been receiving at least 12 consecutive weeks of LOW-INTENSITY statin therapy [i.e., simvastatin 10 mg, pravastatin 10-20 mg, lovastatin 20 mg, fluvastatin 20-40 mg, Livalo (pitavastatin) 1 mg] and will continue to receive a LOW-INTENSITY statin at maximally tolerated dose

**OR**

**2.3** Patient is unable to tolerate low- or moderate-, and high-intensity statins as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms for low- or moderate-, and high-intensity statins:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times ULN)

**OR**

**2.4** Patient has a labeled contraindication to all statins

**OR**

**2.5** Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations greater than 10 times ULN [4]

**AND**

**3** - One of the following:

**3.1** Patient has been receiving at least 12 consecutive weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy

**OR**

**3.2** Patient has a history of contraindication or intolerance to ezetimibe

**AND**

**4** - Patient is unable to maintain adherence to proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor therapy

**AND**

**5** - Submission of medical records (e.g., laboratory values) documenting one of the following LDL-C values while on maximally tolerated lipid lowering therapy within the last 120 days:

- LDL-C greater than or equal to 55 mg/dL for diagnosis of ASCVD [2]
- LDL-C greater than or equal to 100 mg/dL for diagnosis of HeFH [3]

**AND**

**6** - Prescribed by or in consultation with one of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

**AND**

**7** - Medication will not be used in combination with PCSK9 inhibitor therapy [2,3]

Product Name:Leqvio	
Diagnosis	Heterozygous Familial Hypercholesterolemia (HeFH), Atherosclerotic Cardiovascular Disease (ASCVD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting LDL-C reduction from baseline while on therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - One of the following:</p> <p><b>2.1</b> Patient continues to receive other lipid-lowering therapy (e.g., statins, ezetimibe) at the maximally tolerated dose</p>	

**OR**

**2.2** Patient has a documented inability to take other lipid-lowering therapy (e.g., statins, ezetimibe)

**AND**

**3** - Medication will not be used in combination with PCSK9 inhibitor therapy [2,3]

## 2 . Revision History

Date	Notes
9/1/2023	Update to account for 2022 ACC recommendations of a lower LDL th reshould of 55mg/dl for patients with ASCVD at very high risk.

Leucovorin- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99469
<b>Guideline Name</b>	Leucovorin- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Leucovorin tabs	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Methotrexate toxicity prophylaxis</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** Treatment of hematologic toxicity from folic acid antagonists (i.e., pyrimethamine toxicity treatment or trimethoprim toxicity treatment)

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Libervant, Nayzilam, Valtoco

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147227
<b>Guideline Name</b>	Libervant, Nayzilam, Valtoco
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Libervant, Nayzilam, Valtoco	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of epilepsy  <b>AND</b>	

**2** - Requested medication is being prescribed for the acute treatment of intermittent, stereotypic episodes of frequent seizure activity that are distinct from a patient's usual seizure pattern

Product Name: Libervant, Nayzilam, Valtoco	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

Product Name: Nayzilam, Valtoco	
Diagnosis	Requests Exceeding Quantity Limit
Approval Length	12 month(s)
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - Physician has provided rationale for needing to exceed the quantity limit of 2 boxes per 30 days</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - The requested dose is within the FDA (Food and Drug Administration) maximum dose per day</p>	

**2 . Revision History**

Date	Notes
------	-------

5/23/2024	Added Libervant as NP target
-----------	------------------------------

Lidoderm (lidocaine) 5% patches

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117420
<b>Guideline Name</b>	Lidoderm (lidocaine) 5% patches
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Brand Lidoderm patch, generic lidocaine patch	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 The requested drug must be used for a Food and Drug Administration (FDA)-approved indication	

**OR**

**1.2** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmacoeconomic studies
- Other drug reference resources

## **2 . Revision History**

Date	Notes
11/29/2022	Updated approval duration

Likmez (metronidazole) oral suspension

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-143794
<b>Guideline Name</b>	Likmez (metronidazole) oral suspension
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Likmez	
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Both of the following:  1.1 One of the following diagnoses:  1.1.1 Trichomoniasis caused by Trichomonas vaginalis	

**OR**

**1.1.2** Acute intestinal amebiasis (amoebic dysentery) and amebic liver abscess

**OR**

**1.1.3** Treatment of one the following serious infections caused by susceptible anaerobic bacteria:

- Intra-abdominal infections, including peritonitis, intra-abdominal abscess, and liver abscess, caused by Bacteroides species including the B. fragilis group (B. fragilis, B. ovatus, B. thetaiotaomicron, B. vulgatus), Parabacteroides distasonis, Clostridium species, Eubacterium species, Peptococcus species, and Peptostreptococcus species
- Skin and skin structure infections caused by Bacteroides species including the B. fragilis group, Clostridium species, Peptococcus species, Peptostreptococcus species, and Fusobacterium species
- Gynecologic infections, including endometritis, endomyometritis, tubo-ovarian abscess, and postsurgical vaginal cuff infection, caused by Bacteroides species including the B. fragilis group, Clostridium species, Peptococcus species, Peptostreptococcus species, and Fusobacterium species
- Bacterial septicemia caused by Bacteroides species including the B. fragilis group and Clostridium species
- Bone and joint infections, (as adjunctive therapy), caused by Bacteroides species including the B. fragilis group
- Central nervous system (CNS) infections, including meningitis and brain abscess, caused by Bacteroides species including the B. fragilis group
- Lower respiratory tract infections, including pneumonia, empyema, and lung abscess, caused by Bacteroides species including the B. fragilis group
- Endocarditis caused by Bacteroides species including the B. fragilis group

**AND**

**1.2** One of the following:

**1.2.1** Patient has a history of failure, contraindication, or intolerance to metronidazole tablets as evidenced by submission of medical records or claims history

**OR**

**1.2.2** Patient has a swallowing disorder and cannot swallow solid oral dosage forms

## 2 . Revision History

Date	Notes
3/1/2024	Changed guideline name to reflect suspension formulation

Livdelzi (seladelpar)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157798
<b>Guideline Name</b>	Livdelzi (seladelpar)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Livdelzi	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of primary biliary cholangitis (PBC) (also known as primary biliary cirrhosis)	

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims documenting one of the following:

**2.1** Both of the following:

**2.1.1** Patient has failed to achieve an alkaline phosphatase (ALP) level of less than 1.67 times the upper limit of normal (ULN) after at least 12 consecutive months of treatment with ursodeoxycholic acid (UDCA) (e.g., Urso, Urso Forte, ursodiol)

**AND**

**2.1.2** Used in combination with ursodeoxycholic acid (UDCA)

**OR**

**2.2** History of contraindication or intolerance to ursodeoxycholic acid (UDCA)

**AND**

**3** - Requested drug will not be used in combination with Ocaliva (obeticholic acid) or Iqirvo (elafibranor)

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Hepatologist
- Gastroenterologist

Product Name:Livdelzi	
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., ALP level less than 1.67 times ULN, total bilirubin less than or equal to ULN, ALP decrease greater than or equal to 15% from baseline )</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Requested drug will not be used in combination with Ocaliva (obeticholic acid) or Iqirvo (elafibranor)</p>	

## 2 . Revision History

Date	Notes
10/25/2024	New program

Livmarli (maralixibat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269195
<b>Guideline Name</b>	Livmarli (maralixibat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Livmarli	
Diagnosis	Alagille Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming both of the following: <b>1.1</b> Diagnosis of Alagille Syndrome (ALGS)	

**AND**

**1.2** Molecular genetic testing confirms mutations in the JAG1 or NOTCH2 gene

**AND**

**2** - Documentation of ONE of the following:

- Total serum bile acid > 3x the upper limit of normal (ULN)
- Conjugated bilirubin > 1 mg/dL
- Fat soluble vitamin deficiency otherwise unexplainable
- Gammaglutamyl transpeptidase (GGT) > 3x ULN

**AND**

**3** - Patient is experiencing moderate to severe cholestatic pruritus

**AND**

**4** - Patient has had an inadequate response to at least two of the following treatments used for the relief of pruritus:

- Ursodeoxycholic acid (e.g., Ursodiol)
- Antihistamines (e.g., diphenhydramine, hydroxyzine)
- Rifampin
- Bile acid sequestrants (e.g., Questran, Colestid, Welchol)

**AND**

**5** - Patient is 3 months of age or older

**AND**

**6** - Prescribed by or in consultation with a hepatologist

Product Name: Livmarli	
Diagnosis	Progressive Familial Intrahepatic Cholestasis (PFIC)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming both of the following:</p> <p><b>1.1</b> Diagnosis of Progressive familial intrahepatic cholestasis (PFIC)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> Molecular genetic testing confirms mutations in the ATP8B1, ABCB11, ABCB4, TJP2, NR1H4, or MYO5B gene</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is experiencing both of the following:</p> <ul style="list-style-type: none"> <li>• Moderate to severe pruritus</li> <li>• Patient has a serum bile acid concentration above the upper limit of the normal reference for the reporting laboratory</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Patient is 12 months of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Patient has had an inadequate response to at least two of the following treatments used for the relief of pruritus:</p> <ul style="list-style-type: none"> <li>• Ursodeoxycholic acid (e.g., Ursodiol)</li> <li>• Antihistamines (e.g., diphenhydramine, hydroxyzine)</li> </ul>	

- Rifampin
- Bile acid sequestrants (e.g., Questran, Colestid, Welchol)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Hepatologist
- Gastroenterologist

Product Name: Livmarli	
Diagnosis	All indications listed above
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., reduced bile acids, reduced pruritus severity score)</p>	

## 2 . Revision History

Date	Notes
5/29/2025	Added GPIs for new tablet formulations

Livtency (maribavir)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-113529
<b>Guideline Name</b>	Livtency (maribavir)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/8/2022
-----------------	----------

## 1 . Criteria

Product Name:Livtency	
Diagnosis	CMV infection/disease
Approval Length	8 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of cytomegalovirus (CMV) infection/disease as confirmed by one of the following methods: <ul style="list-style-type: none"><li>quantitative polymerase chain reaction (qPCR)</li></ul>	

- CMV pp65 antigenemia

**AND**

**2** - Patient is a recipient of one of the following:

- Hematopoietic stem cell transplant
- Solid organ transplant

**AND**

**3** - Trial and failure of a minimum 2 weeks duration, contraindication, or intolerance to one of the following therapies at an appropriately indicated dose:

- Intravenous (IV) ganciclovir
- Oral valganciclovir
- IV foscarnet
- IV cidofovir

**AND**

**4** - Patient is 12 years of age or older

**AND**

**5** - Patient weighs greater than or equal to 35kg

**AND**

**6** - Prescribed by or in consultation with a provider who specializes in one of the following areas:

- Transplant
- Infectious Disease

## **2 . Revision History**

Date	Notes
9/8/2022	Removed references and end note, no changes to clinical criteria.

Lodoco (colchicine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135381
<b>Guideline Name</b>	Lodoco (colchicine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Lodoco	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cardiovascular disease (CV)  <b>AND</b>	

**2** - Used for the secondary prevention of CV disease (e.g., very high-risk patients – see Table 1)

**AND**

**3** - Patient is on guideline therapy management for multiple risk factors (e.g., dyslipidemia, hypertension, hyperglycemia) associated with CV disease

**AND**

**4** - Submission of medical records (e.g., chart notes) or paid claims documenting trial and failure or intolerance to colchicine 0.6 mg tablets

Product Name:Lodoco	
Approval Length	6 months [C, 4]
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy(e.g., reduced risk of cardiovascular death, myocardial infarction, ischemia-driven coronary revascularization)	

## 2 . Background

<b>Clinical Practice Guidelines</b>
Table 1 [3]
Definition of Very High-Risk
History of multiple major ASCVD events
OR

<b>One Major ASCVD event AND 2 or more high risk conditions</b>
<b>Major ASCVD Events</b>
Recent ACS (within the past 12 months)
History of MI (other than recent ACS events listed above)
History of ischemic stroke
Symptomatic peripheral artery disease (history of claudication with ABI <0.85, or previous revascularization or amputation)
<b>High-Risk Conditions</b>
Age 65 or older
Familial hypercholesterolemia
History of previous coronary artery bypass graft surgery or percutaneous coronary intervention outside of the major ASCVD event(s)
Diabetes
Hypertension
Chronic kidney disease (eGFR, 15–59 mL/min/1.73 m <sup>2</sup> )
Current tobacco smoking
Persistently elevated LDL-C ≥100 mg/dL despite maximally tolerated statin therapy and ezetimibe
History of congestive heart failure
<b>ABI</b> indicates ankle brachial index; <b>ACS</b> , acute coronary syndrome; <b>ASCVD</b> , atherosclerotic cardiovascular disease; <b>CKD</b> , chronic kidney disease; <b>eGFR</b> , estimated glomerular filtration rate; <b>LDL-C</b> , low-density lipoprotein cholesterol; and <b>MI</b> , myocardial infarction.

### 3 . Revision History

Date	Notes
10/27/2023	New program

Long-Acting Opioid Products - AZM



### Prior Authorization Guideline

<b>Guideline ID</b>	GL-163760
<b>Guideline Name</b>	Long-Acting Opioid Products - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	2/1/2025
-----------------	----------

**Note:**

\*\*PLEASE NOTE: This guideline contains criteria for multiple DUR rejection codes. PA Reviewers: Please confirm the correct criteria is being utilized for reviews based on Diagnosis and Guideline Type. Criteria for MME EXCEEDED rejections can be found towards the end of the guideline.

### 1 . Criteria

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets	
Diagnosis	PA REQUIRED for use of MAT and other Opioids (Reject 88)
Guideline Type	DUR

**Approval Criteria**

1 - Provider attests to notify the prescriber of the MAT therapy and the prescriber of the MAT therapy approves the concurrent opioid therapy.

**AND**

2 - The days supply does not exceed 14 days for a surgical procedure.

**AND**

3 - The days supply does not exceed 5 days for all other requests.

**AND**

4 - There has not been a previous approval in the last 6 months.

Notes	Approval Length: 14 Days for surgical procedure, 5 Days for all other requests
-------	--

Product Name: PREFERRED: fentanyl transdermal 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr, 100 mcg/hr; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets	
Diagnosis	Cancer related pain/Hospice care/end-of-life care*
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - ONE of the following:	
1.1 Patient is being treated for cancer	

**OR**

**1.2 Patient is receiving hospice or end-of-life care**

Notes

\*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 30 day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment.

Product Name:NON-PREFERRED: fentanyl transdermal 37.5mcg/hr, 62.5mcg/hr, and 87.5mcg/hr; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER

Diagnosis

Cancer related pain/Hospice care/end-of-life care\*

Approval Length

12 month(s)

Guideline Type

Prior Authorization

**Approval Criteria**

**1 - ONE of the following:**

**1.1 Patient is being treated for cancer**

**OR**

**1.2 Patient is receiving hospice or end-of-life care**

**AND**

**2 - BOTH of the following:**

**2.1 ONE of the following:**

**2.1.1** The patient has a history of failure, contraindication or intolerance to a trial of at least THREE of the following (Document drugs and date of trials):\*

- morphine sulfate controlled release tablets (specifically generic MS Contin)
- preferred fentanyl transdermal (12mcg, 25mcg, 50mcg, 75mcg, 100mcg)\*\*
- Brand Butrans (buprenorphine) transdermal
- Brand Oxycontin
- tramadol extended release tablets (non-biphasic release tablets)

**OR**

**2.1.2** Patient is established on pain therapy with the requested medication for cancer, hospice care, or end-of-life care pain, and the medication is not a new regimen for treatment of cancer, hospice care, or end-of-life care pain (Document date regimen was started)

**AND**

**2.2** Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

Notes	<p>*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. *Note: If the request is for a non-preferred product and the member is currently taking the requested long-acting opioid for at least 30 days and has met the medical necessity authorization criteria requirements for treatment with an opioid, but has not tried the preferred alternatives a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. **NOTE: Fentanyl transdermal 37.5mcg/hr, 62.5mcg/hr, and 87.5mcg/hr are non-preferred. *Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial.</p>
-------	---

Product Name:NON-PREFERRED: Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets	
Diagnosis	Cancer related pain/Hospice care/end-of-life care*
Approval Length	12 month(s)

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 Patient is being treated for cancer</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 Patient is receiving hospice or end-of-life care</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - BOTH of the following:</p> <p>2.1 ONE of the following:</p> <p>2.1.1 The patient has a history of failure, contraindication or intolerance to a trial of BOTH of the following (Document drugs and date of trials):*</p> <ul style="list-style-type: none"> <li>• tramadol immediate release (IR)</li> <li>• tramadol extended release tablets (non-biphasic release tablets)</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>2.1.2 Patient is established on pain therapy with the requested medication for cancer, hospice care, or end-of-life care pain, and the medication is not a new regimen for treatment of cancer, hospice care, or end-of-life care pain (Document date regimen was started)</p> <p style="text-align: center;"><b>AND</b></p> <p>2.2 Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided</p>	
Notes	*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial sho

	uld be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. *Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial.
--	--

Product Name: PREFERRED: fentanyl transdermal 12 mcg/hr, 25 mcg/hr, 50 mcg/hr, 75 mcg/hr, 100 mcg/hr; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets

Diagnosis	Non-cancer pain/Non-hospice care/Non-end-of-life care pain*
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Prescriber attests to ALL of the following:

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment
- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression
- Pain is moderate to severe and expected to persist for an extended period of time
- Pain is chronic
- Pain is not postoperative (unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time)
- Pain management is required around the clock with a long-acting opioid

**AND**

2 - ONE of the following:

2.1 Prior to the start of therapy with the long-acting opioid, the patient has failed an adequate

(minimum of 2 week) trial of a short-acting opioid within the last 30 days (Document drug(s) and date of trial)\*

**OR**

**2.2** The patient is already receiving chronic opioid therapy prior to surgery for postoperative pain

**OR**

**2.3** Postoperative pain is expected to be moderate to severe and persist for an extended period of time

**AND**

**3** - If the request for neuropathic pain (examples of neuropathic pain include neuralgias, neuropathies, fibromyalgia), BOTH of the following:

**3.1** Unless it is contraindicated, the patient has not exhibited an adequate response to 8 weeks of treatment with gabapentin titrated to a therapeutic dose (Document date of trial)\*

**AND**

**3.2** Unless it is contraindicated, the patient has not exhibited an adequate response to at least 6 weeks of treatment with a tricyclic antidepressant titrated to the maximum tolerated dose. (Document drug and date of trial)\*

Notes

\*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. \*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial \*\*NOTE: Fentanyl transdermal 37.5mcg/hr, 62.5mcg/hr, and 87.5 mcg/hr are non-preferred.

Product Name:NON-PREFERRED: fentanyl transdermal 37.5mcg/hr, 62.5mcg/hr, and 87.5mcg/hr; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone

tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER

Diagnosis	Non-cancer pain/Non-hospice care/Non-end-of-life care pain*
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Prescriber attests to ALL of the following:

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment
- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression
- Pain is moderate to severe and expected to persist for an extended period of time
- Pain is chronic
- Pain is not postoperative (unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time)
- Pain management is required around the clock with a long-acting opioid

**AND**

2 - ONE of the following:

**2.1** Prior to the start of therapy with the long-acting opioid, the patient has failed an adequate (minimum of 2 week) trial of a short-acting opioid within the last 30 days (Document drug(s) and date of trial)\*

**OR**

**2.2** The patient is already receiving chronic opioid therapy prior to surgery for postoperative pain

**OR**

**2.3** Postoperative pain is expected to be moderate to severe and persist for an extended period of time

**AND**

**3** - The patient has a history of failure, contraindication or intolerance to at least **THREE** of the following (Document drugs and date of trials):\*

- morphine sulfate controlled release tablets (specifically generic MS Contin)
- preferred fentanyl transdermal (12mcg, 25mcg, 50mcg, 75mcg, 100mcg)\*\*
- Brand Butrans (buprenorphine) transdermal
- Brand Oxycontin
- tramadol extended release tablets (non-biphasic release tablets)

**AND**

**4** - If the request for neuropathic pain (examples of neuropathic pain include neuralgias, neuropathies, fibromyalgia), **BOTH** of the following:

**4.1** Unless it is contraindicated, the patient has not exhibited an adequate response to 8 weeks of treatment with gabapentin titrated to a therapeutic dose (Document date of trial) )\*

**AND**

**4.2** Unless it is contraindicated, the patient has not exhibited an adequate response to at least 6 weeks of treatment with a tricyclic antidepressant titrated to the maximum tolerated dose. (Document drug and date of trial) )\*

Notes

\*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. \*Note: If the request is for a non-preferred product and the member is currently taking the requested long-acting opioid for at least 30 days and has met th

	<p>e medical necessity authorization criteria requirements for treatment with an opioid, but has not tried the preferred alternatives a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. Additionally **NOTE: Fentanyl transdermal 37.5mcg/hr, 62.5mcg/hr, and 87.5 mcg/hr are non-preferred. *Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial.</p>
--	---

<b>Product Name:NON-PREFERRED: Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets</b>	
Diagnosis	Non-cancer pain/Non-hospice care/Non-end-of-life care pain*
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Prescriber attests to ALL of the following:</p> <ul style="list-style-type: none"> <li>• The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided</li> <li>• Treatment goals are defined, including estimated duration of treatment</li> <li>• Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention</li> <li>• Patient has been screened for substance abuse/opioid dependence</li> <li>• If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression</li> <li>• Pain is moderate to severe and expected to persist for an extended period of time</li> <li>• Pain is chronic</li> <li>• Pain is not postoperative (unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time)</li> <li>• Pain management is required around the clock with a long-acting opioid</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p>	

**2.1** Prior to the start of therapy with the long-acting opioid, the patient has failed an adequate (minimum of 2 week) trial of a short-acting opioid within the last 30 days (Document drug(s) and date of trial)\*

**OR**

**2.2** The patient is already receiving chronic opioid therapy prior to surgery for postoperative pain

**OR**

**2.3** Postoperative pain is expected to be moderate to severe and persist for an extended period of time

**AND**

**3** - If the request for neuropathic pain (examples of neuropathic pain include neuralgias, neuropathies, fibromyalgia), BOTH of the following:

**3.1** Unless it is contraindicated, the patient has not exhibited an adequate response to 8 weeks of treatment with gabapentin titrated to a therapeutic dose (Document date of trial)\*

**AND**

**3.2** Unless it is contraindicated, the patient has not exhibited an adequate response to at least 6 weeks of treatment with a tricyclic antidepressant titrated to the maximum tolerated dose. (Document drug and date of trial)\*

**AND**

**4** - The patient has a history of failure, contraindication or intolerance to BOTH of the following (Document drugs and date of trials): )\*

- tramadol immediate release (IR)\*\*
- tramadol extended release tablets (non-biphasic release tablets)\*\*

Notes	<p>*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. *Note: If the request is for tramadol extended release capsules or tramadol extended release biphasic release tablets and the member is currently taking the requested long-acting opioid for at least 30 days and has met the medical necessity authorization criteria requirements for treatment with an opioid, but has not tried the preferred alternatives a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. *Drug may require prior authorization *Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial.</p>
-------	---

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets	
Diagnosis	Non-cancer pain/Non-hospice care/Non-end-of-life care pain*
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates meaningful improvement in pain and function (Document improvement in function or pain score improvement)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Identify rationale for not tapering and discontinuing opioid (Document rationale)</p> <p style="text-align: center;"><b>AND</b></p>	

**3 - Prescriber attests to ALL of the following:**

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment
- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression
- Pain is moderate to severe and expected to persist for an extended period of time
- Pain is chronic
- Pain is not postoperative (unless the patient is already receiving chronic opioid therapy prior to surgery, or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time)
- Pain management is required around the clock with a long-acting opioid

**Notes**

\*Note: If the member is currently taking the requested long-acting opioid for at least 30 days and does not meet the medical necessity authorization criteria requirements for treatment with an opioid, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment. \*Note: If the request is for a non-preferred product and the member is currently taking the requested long-acting opioid for at least 30 days and has met the medical necessity authorization criteria requirements for treatment with an opioid, but has not tried the preferred alternatives a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested drug/strength combination up to the requested quantity for transition to an alternative treatment.\*\*NOTE: Fentanyl transdermal 37.5mcg/hr, 62.5mcg/hr, and 87.5mcg/hr are non-preferred.

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets

Diagnosis	Criteria for Quantity Limit Reviews*
Guideline Type	Quantity Limit

**Approval Criteria**

1 - The requested dose cannot be achieved by moving to a higher strength of the product

**AND**

2 - The requested dose is within the Food and Drug Administration (FDA) maximum dose per day, where an FDA maximum dose per day exists (see Table 1 in the Background section)

Notes	*Note: Authorization will be issued for <ul style="list-style-type: none"><li>• Cancer pain/hospice/end-of-life related pain: 12 months</li><li>• All Tramadol ER requests: 6 months</li><li>• Non-cancer pain/non-hospice/non-end-of-life related pain: 6 months</li></ul>
-------	---

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets

Diagnosis	Opioid Naïve (Not having filled an opioid in the past 120 days)*
Guideline Type	Morphine Milligram Equivalents (MME)** MME 50.00 exceeded; PA Required for dosage above 50 MEDD

**Approval Criteria**

1 - Opioid naïve members may receive greater than 50 morphine milligram equivalent (MME) based on the following:

1.1 If the request is for 50 MME to 90 MME, ONE of the following (NOTE: If the request exceeds 90 MME please skip this section and proceed to the Exceeding the 90 MME Cumulative Threshold Reviews section):

1.1.1 Diagnosis of ONE of the following:

- Cancer
- End of life pain (including hospice care)
- Palliative care

- Sickle cell anemia

**OR**

**1.1.2** Patient is currently exceeding 50 MME and prescriber attests patient has been on a short-acting opioid in the past 120 days

**OR**

**1.1.3** Document ALL of the following:

- The diagnosis associated with the need for pain management with opioid
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression
- The prescriber has acknowledged that they have completed an addiction risk and risk of overdose assessment
- Prescriber attests the member requires more than 50 MME per day to adequately control pain

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets

Diagnosis	Doses Exceeding the Cumulative MME of 90 mg - Cancer/Hospice/End-of-Life/Palliative Care/Skilled Nursing Facility/Traumatic Injury Related Pain*
Approval Length	12 month(s)
Guideline Type	Morphine Milligram Equivalent (MME)** (MME 90.00 exceeded; PA REQUIRED; Dosage Above MEDD Limit)

**Approval Criteria**

**1** - Doses exceeding the cumulative morphine milligram equivalent (MME) of 90 milligrams

will be approved up to the requested amount for ALL opioid products if the patient has one of the following conditions:

- Active oncology diagnosis
- Hospice care
- End-of-life care (other than hospice)
- Palliative care
- Skilled nursing facility care
- Traumatic injury, including burns and excluding post-surgical procedure

**AND**

**2** - Provider attests patient has been prescribed naloxone (may also be verified via paid pharmacy claims)

Notes	*Note: Authorization will be issued for 12 months for one of the above conditions. The authorization should be entered for an MME of 9999 so as to prevent future disruptions in therapy if the patient's dose is increased.
-------	--

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets

Diagnosis	Doses- Exceeding the Cumulative MME of 90 mg - Non-cancer/non-hospice/non-end-of-life/non-palliative care/non-skilled Nursing Facility/Traumatic Injury Related Pain*
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Morphine Milligram Equivalent (MME)** MME 90.00 exceeded; PA REQUIRED; Dosage Above MEDD Limit

**Approval Criteria**

**1** - Prescriber attests to ALL of the following:

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment

- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- if used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression

**AND**

**2 - BOTH of the following:**

**2.1** Patient has tried and failed non-opioid pain medication (document drug name and date of trial)

**AND**

**2.2** Opioid medication doses of less than 90 morphine milligram equivalent (MME) have been tried and did not adequately control pain (document drug regimen or MME and dates of therapy)

**AND**

**3 - Provider attests patient has been prescribed naloxone (may also be verified via paid pharmacy claims)**

Notes	*Note: If the member has been established on the requested MME dose for at least 30 days and does not meet the medical necessity authorization criteria requirements, a denial should be issued and a maximum 30 -day authorization may be authorized one time for the requested MME dose. **Note: Authorization will be issued for 6 months for non-cancer/non-hospice/non-end-of-life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain up to the current requested MME plus 90 MME.
-------	---

Product Name:fentanyl transdermal; generic morphine sulfate ER tablets; Brand Oxycontin, generic tramadol ER tablets; generic hydromorphone ER; Brand hydrocodone ER capsules; Brand Hysingla ER; generic hydrocodone ER tablets; Brand Methadose; generic methadone tablets/concentrate/soln/tablets for oral susp; Brand morphine sulfate ER capsules; Brand MS Contin; Nucynta ER; Brand oxymorphone ER; Brand oxycodone ER; Brand Conzip; Brand tramadol ER 24HR biphasic capsules; Brand tramadol ER biphasic release tablets

Diagnosis	Doses Exceeding the Cumulative MME of 90 mg - Non-cancer/non-hospice/non-end-of-life/non-palliative Nursing Facility/Traumatic Injury Related Pain* care/non-skilled
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Morphine Milligram Equivalent (MME)** MME 90.00 exceeded; PA REQUIRED; Dosage Above MEDD Limit

**Approval Criteria**

1 - Prescriber attests to ALL of the following:

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment
- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- if used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression

**AND**

2 - Identify rationale for not tapering and discontinuing opioid (Document rationale)

**AND**

3 - Patient demonstrates meaningful improvement in pain and function (Document improvement in function or pain score improvement)

**AND**

4 - Provider attests patient has been prescribed naloxone (may also be verified via paid pharmacy claims)

Notes	<p>*Note: If the member has been established on the requested MME dose for at least 30 days and does not meet the medical necessity authorization criteria requirements, a denial should be issued and a maximum 30 -day authorization may be authorized one time for the requested MME dose. **Note: Authorization will be issued for 6 months for non-cancer/non-hospice/non-end-of-life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain up to the current requested MME plus 90 MME.</p>
-------	--

**2 . Background**

Benefit/Coverage/Program Information	
<p><b>Table 1. CDC Recommended Long-Acting Opioid Maximum Milligram Morphine Equivalents per Day*</b></p>	
Active Ingredient	FDA Label Max Daily Doses
Morphine	None
Hydromorphone	None
Fentanyl transdermal, mcg/hr	None
Hydrocodone	None
Methadone	None
Tapentadol	500mg ER
Oxymorphone	None
<p>*Doses are not considered equianalgesic and table does not represent a dose conversion chart.</p>	

Max MME is the maximum dose per day based on morphine milligram equivalents allowed without consultation or prescription by a pain specialist. Max MME is based upon the CDC guidelines and adjusted for currently available product strengths. Fentanyl is dosed in mcg/hr rather than mg/day

### 3 . Revision History

Date	Notes
1/31/2025	Added "pain is chronic" bullet to 1st NP/non-cancer initial auth bucket to align with guideline. Moved all targets to non-cancer reauth, removed PDL/NPD reauth buckets. Confirmed all targets added to appropriate sections

Lonhala and Yupelri - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-161649
<b>Guideline Name</b>	Lonhala and Yupelri - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Lonhala Magnair, Yupleri	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderate to severe chronic obstructive pulmonary disease (COPD  <b>AND</b>	

**2 - ONE of the following:**

**2.1 History of failure, contraindication or intolerance to ALL of the following:**

- Brand Spiriva (tiotropium)
- Tudorza (aclidinium)
- Incruse Ellipta (umeclidinium) (\*May require PA)

**OR**

**2.2 BOTH of the following:**

**2.2.1 Patient is unable to use a metered-dose, dry powder or slow mist inhaler (e.g. Spiriva Handihaler) to control his/her COPD due to ONE of the following**

- Cognitive or physical impairment limiting coordination of handheld devices (e.g., cognitive decline, arthritis in the hands) (Document impairment)
- Patient is unable to generate adequate inspiratory force (e.g., peak inspiratory flow rate (PIFR) resistance is less than 60 Liters per minute)

**AND**

**2.2.2 History of failure, contraindication or intolerance to ipratropium nebulized solution (generic Atrovent)**

Product Name:Lonhala Magnair, Yupleri	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	

**2 - ONE of the following:**

**2.1 History of failure, contraindication or intolerance to ALL of the following:**

- Brand Spiriva (tiotropium)
- Tudorza (aclidinium)
- Incruse Ellipta (umeclidinium) (\*May require PA)

**OR**

**2.2 BOTH of the following:**

**2.2.1 Patient is unable to use a metered-dose, dry powder or slow mist inhaler (e.g. Spiriva Handihaler) to control his/her COPD due to ONE of the following**

- Cognitive or physical impairment limiting coordination of handheld devices (e.g., cognitive decline, arthritis in the hands) (Document impairment)
- Patient is unable to generate adequate inspiratory force (e.g., peak inspiratory flow rate (PIFR) resistance is less than 60 Liters per minute)

**AND**

**2.2.2 History of failure, contraindication or intolerance to ipratropium nebulized solution (generic Atrovent)**

## **2 . Revision History**

Date	Notes
12/4/2024	Added additional prerequisites to embedded step, added step to reauth.

Lucemyra (lofexidine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-154516
<b>Guideline Name</b>	Lucemyra (lofexidine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Brand Lucemyra, generic lofexidine	
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - For symptoms of abrupt opioid withdrawal  <p style="text-align: center;"><b>AND</b></p> 2 - Opioids have been discontinued	

**AND**

**3** - History of failure, contraindication, or intolerance to clonidine as verified by recent clonidine claims history in the past 180 days

**AND**

**4** - Prescriber must verify patient has been screened for hepatic and renal impairment and that dosing is appropriate for the patient's degree of hepatic and renal function

**AND**

**5** - Prescriber must verify patient's vital signs have been monitored and that the patient is capable of and has been instructed on self-monitoring for hypotension, orthostasis, bradycardia, and associated symptoms

**AND**

**6** - Patient does not have severe coronary insufficiency, a recent myocardial infarction, cerebrovascular disease, chronic renal failure, or marked bradycardia

**AND**

**7** - Patient does not have congenital long QT syndrome

**AND**

**8** - For Brand Lucemyra requests ONLY, history of failure or intolerance to generic lofexidine

## **2 . Revision History**

Date	Notes
------	-------

9/24/2024	Added step through generic lofexidine, removed bypass for step through clonidine. Updated guideline name
-----------	--

Lumizyme -Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99471
<b>Guideline Name</b>	Lumizyme -Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Lumizyme	
Diagnosis	Pompe disease
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Pompe disease (acid alpha-glucosidase [GAA] deficiency)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Lupkynis (voclosporin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-153491
<b>Guideline Name</b>	Lupkynis (voclosporin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Lupkynis	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of active lupus nephritis  <b>AND</b>	

**2** - Provider attests to ONE of the following:

- Diagnosis is biopsy proven
- Biopsy is contraindicated in the patient

**AND**

**3** - Prescribed in combination with a background immunosuppressive therapy regimen (e.g., mycophenolate mofetil and corticosteroids)

**AND**

**4** - Patient is NOT receiving Lupkynis in combination with either of the following:

- Cyclophosphamide
- Benlysta (belimumab)

**AND**

**5** - Prescribed by ONE of the following:

- Nephrologist
- Rheumatologist

Product Name:Lupkynis	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Lupkynis therapy	

**AND**

**2** - Prescribed in combination with a background immunosuppressive therapy regimen (e.g., mycophenolate mofetil and corticosteroids)

**AND**

**3** - Patient is NOT receiving Lupkynis in combination with either of the following:

- Cyclophosphamide
- Benlysta (belimumab)

**AND**

**4** - Prescribed by ONE of the following:

- Nephrologist
- Rheumatologist

**AND**

**5** - ONE of the following:

**5.1** Patient has been on Lupkynis therapy for less than 12 months

**OR**

**5.2** BOTH of the following:

**5.2.1** Patient has completed 12 or more months of Lupkynis therapy

**AND**

**5.2.2** The provider attests that the benefit of continuation of therapy exceeds the risk in light of the patient's treatment response and risk of worsening nephrotoxicity

**2 . Revision History**

Date	Notes
9/26/2024	New program

Lyfgenia (lovotibeglogene autotemcel)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-211206
<b>Guideline Name</b>	Lyfgenia (lovotibeglogene autotemcel)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Lyfgenia	
Approval Length	1 Time Authorization in Lifetime*
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of sickle cell disease (SCD) confirmed by patient genotype $\beta S/\beta S$ , $\beta S/\beta 0$ , or $\beta S/\beta +$  <b>AND</b>	

**2** - Patient is 12 years of age or older

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to hydroxyurea (defined as being unable to take hydroxyurea per health care professional judgement) at any point in the past

**AND**

**4** - Provider attests that patient is clinically stable and eligible to undergo hematopoietic stem cell transplant (HSCT)

**AND**

**5** - Submission of medical records (e.g., chart notes) documenting one of the following:

**5.1** Patient has a history of at least 4 vaso-occlusive events (VOEs) in the past 24 months defined by one of following scenarios:

- an episode of acute pain with no medically determined cause other than vaso-occlusion, lasting more than 2 hours
- acute chest syndrome (ACS)
- acute hepatic sequestration
- acute splenic sequestration
- VOE requiring a hospitalization or multiple visits to an emergency department/urgent care over 72 hours and receiving intravenous medications at each visit
- priapism requiring any level of medical attention

**OR**

**5.2** Patient is currently receiving chronic transfusion therapy for recurrent Vaso-Occlusive Events (VOEs)

**AND**

**6** - Prescribed by a provider at a SCD treatment center with expertise in gene therapy

**AND**

**7** - Prescribed by or in consultation with a board-certified hematologist with SCD expertise

**AND**

**8** - Patient has never received any previous sickle cell gene therapy treatment in their lifetime (i.e., Casgevy, Lyfgenia)

Notes	*Per prescribing information, Lyfgenia is for one-time, single dose intravenous use only.
-------	---

## 2 . Revision History

Date	Notes
3/20/2025	Criteria updated.

Lyrice

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-105529
<b>Guideline Name</b>	Lyrice
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

### 1 . Criteria

Product Name:Brand Lyrice	
Diagnosis	Seizure Disorder
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of seizure disorder  <b>AND</b>	

**2** - History of failure, contraindication, or intolerance to generic pregabalin immediate-release capsules or generic pregabalin solution

Product Name: Brand Lyrica

Diagnosis	Neuropathic Pain Associated with Spinal Cord Injury
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of neuropathic pain associated with spinal cord injury

**AND**

**2** - One of the following:

- History of failure to generic pregabalin immediate-release capsules or solution at a minimum dose of 300mg daily for 4 weeks
- Contraindication or intolerance to generic pregabalin immediate-release capsules or solution

Product Name: Brand Lyrica

Diagnosis	Fibromyalgia
-----------	--------------

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of fibromyalgia

**AND**

**2** - One of the following:

- History of failure to generic pregabalin immediate-release capsules or solution at a minimum dose of 300mg daily for 4 weeks
- Contraindication or intolerance to generic pregabalin immediate-release capsules or solution

Product Name:Brand Lyrica	
Diagnosis	Diabetic peripheral neuropathy (DPN)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of diabetic peripheral neuropathy (DPN)	
<b>AND</b>	
2 - One of the following:	
<ul style="list-style-type: none"><li>• History of failure to generic pregabalin immediate-release capsules or solution at a minimum dose of 300mg daily for 4 weeks</li><li>• Contraindication or intolerance to generic pregabalin immediate-release capsules or solution</li></ul>	

Product Name:Brand Lyrica	
Diagnosis	Post herpetic neuralgia (PHN)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of post herpetic neuralgia (PHN)

**AND**

2 - One of the following:

- History of failure to generic pregabalin immediate-release capsules or solution at a minimum dose of 300mg daily for 4 weeks
- Contraindication or intolerance to generic pregabalin immediate-release capsules or solution

Product Name:Lyrice CR	
Diagnosis	Diabetic peripheral neuropathy (DPN)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of diabetic peripheral neuropathy (DPN)	
<b>AND</b>	
2 - History of failure, contraindication, or intolerance to gabapentin (generic Neurontin) at a minimum dose of 1800 milligrams daily for 4 weeks	
<b>AND</b>	
3 - History of failure, contraindication, or intolerance to treatment with ONE of the following:	
<ul style="list-style-type: none"><li>• Tricyclic antidepressant at the maximum tolerated dose for 6 to 8 weeks, or intolerance to a tricyclic antidepressant</li><li>• Serotonin and norepinephrine reuptake inhibitor (SNRI) antidepressant (i.e. duloxetine, venlafaxine)</li></ul>	

**AND**

**4** - History of failure, contraindication, or intolerance to generic pregabalin immediate-release capsules or generic pregabalin solution

Product Name:Lyrice CR

Diagnosis	Post herpetic neuralgia (PHN)
-----------	-------------------------------

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of post herpetic neuralgia (PHN)

**AND**

**2** - History of failure, contraindication, or intolerance to gabapentin (generic Neurontin) at a minimum dose of 1800 milligrams daily for 4 weeks

**AND**

**3** - History of failure, contraindication, or intolerance to a tricyclic antidepressant at the maximum tolerated dose for 6 to 8 weeks

**AND**

**4** - History of failure, contraindication, or intolerance to generic pregabalin immediate-release capsules or generic pregabalin solution

**2 . Revision History**

Date	Notes
------	-------

3/31/2022	Added step through generic for seizure indication. Updated all indications to allow for any manufacturer of generic immediate-release capsules or solution.
-----------	---

Lysteda

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99473
<b>Guideline Name</b>	Lysteda
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Lysteda, generic tranexamic acid	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cyclic heavy menstrual bleeding	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Makena- AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114911
<b>Guideline Name</b>	Makena- AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/5/2022
-----------------	-----------

## 1 . Criteria

Product Name:Brand Makena*, generic hydroxyprogesterone caproate*	
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Current singleton pregnancy  <b>AND</b>  2 - History of a prior spontaneous preterm birth of a singleton pregnancy	

**AND**

**3** - Treatment is initiated between 16 weeks, 0 days of gestation and 20 weeks, 6 days of gestation

**AND**

**4** - Administration is to continue weekly until week 37 (through 36 weeks, 6 days) of gestation or delivery, whichever occurs first

**AND**

**5** - Applies to generic hydroxyprogesterone caproate ONLY: patient has a history of failure, contraindication or intolerance to Brand Makena

Notes	*NOTE: Approval duration is up to 21 weeks; approval duration should take into account gestation week when Makena will be started and only authorized up to week 37.
-------	--

## 2 . Revision History

Date	Notes
10/4/2022	Updated gestational days for drug initiation to align w PI

Marinol, Syndros

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-108625
<b>Guideline Name</b>	Marinol, Syndros
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/23/2022
-----------------	-----------

### 1 . Criteria

Product Name:Brand Marinol, Syndros	
Diagnosis	Chemotherapy-induced nausea and vomiting
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Patient is receiving cancer chemotherapy  <b>AND</b>	

**2 - ONE of the following:**

**2.1** History of failure, contraindication, or intolerance to formulary generic dronabinol

**OR**

**2.2** Patient is unable to swallow capsules

**AND**

**3 -** History of failure, contraindication, or intolerance to a 5HT-3 (5-hydroxytryptamine) receptor antagonist [eg, Anzemet (dolasetron), Kytril (granisetron), or Zofran (ondansetron)]

**AND**

**4 -** History of failure, contraindication, or intolerance to **ONE** of the following:

- Ativan (lorazepam)
- Compazine (prochlorperazine)
- Decadron (dexamethasone)
- Haldol (haloperidol)
- Phenergan (promethazine)
- Reglan (metoclopramide)
- Zyprexa (olanzapine)

Product Name:Generic Dronabinol	
Diagnosis	Chemotherapy-induced nausea and vomiting
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient is receiving cancer chemotherapy	

**AND**

**2** - History of failure, contraindication, or intolerance to a 5HT-3 (5-hydroxytryptamine) receptor antagonist [eg, Anzemet (dolasetron), Kytril (granisetron), or Zofran (ondansetron)]

**AND**

**3** - History of failure, contraindication, or intolerance to **ONE** of the following:

- Ativan (lorazepam)
- Compazine (prochlorperazine)
- Decadron (dexamethasone)
- Haldol (haloperidol)
- Phenergan (promethazine)
- Reglan (metoclopramide)
- Zyprexa (olanzapine)

Product Name: Brand Marinol, Syndros	
Diagnosis	Anorexia in Patients with AIDS
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of anorexia with weight loss in patients with AIDS (acquired immunodeficiency syndrome)</p> <p><b>AND</b></p> <p><b>2</b> - Patient is on antiretroviral therapy</p> <p><b>AND</b></p>	

**3 - ONE of the following:**

**3.1** Patient is 65 years of age or greater

**OR**

**3.2 BOTH of the following:**

- Patient is less than 65 years of age
- History of failure, contraindication, or intolerance to Megace (megestrol)

**AND**

**4 - ONE of the following:**

**4.1** History of failure, contraindication, or intolerance to formulary generic dronabinol

**OR**

**4.2** Patient is unable to swallow capsules

Product Name:Generic dronabinol	
Diagnosis	Anorexia in Patients with AIDS
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of anorexia with weight loss in patients with AIDS (acquired immunodeficiency syndrome)	
<b>AND</b>	
2 - Patient is on antiretroviral therapy	

**AND**

**3** - ONE of the following:

**3.1** Patient is 65 years of age or greater

**OR**

**3.2** BOTH of the following:

- Patient is less than 65 years of age
- History of failure, contraindication, or intolerance to Megace (megestrol)

## **2 . Revision History**

Date	Notes
6/23/2022	Removed cesamet from guideline name. Added Brand Marinol as NP target

Mepron

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99474
<b>Guideline Name</b>	Mepron
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Mepron, generic atovaquone	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following:  1.1 BOTH of the following:  1.1.1 The patient has a diagnosis (e.g. human immunodeficiency virus [HIV]) warranting Pneumocystis jirovecii pneumonia (PCP) infection prophylaxis	

**AND**

**1.1.2** The patient has a documented intolerance or contraindication to trimethoprim-sulfamethoxazole (TMP-SMX) and dapsone

**OR**

**1.2** BOTH of the following:

**1.2.1** The patient has a diagnosis of mild to moderate pneumonia caused by *P. jirovecii*

**AND**

**1.2.2** The patient has a documented intolerance, contraindication, or history of treatment failure to TMP-SMX

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Metformin products - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-115355
<b>Guideline Name</b>	Metformin products - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/13/2022
-----------------	------------

### 1 . Criteria

Product Name:generic metformin 625 mg immediate-release tablets	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of greater than or equal to 12 week trial of preferred metformin immediate-release products	

Product Name:generic metformin extended-release (generic for Fortamet and generic for Glumetza)
---

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ALL of the following:</p> <p>1.1 History of greater than or equal to 12 week trial of metformin extended-release (generic Glucophage XR)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 ONE of the following:</p> <p>1.2.1 Submission of medical records (e.g. chart notes, laboratory values) documenting an inadequate response to metformin extended-release (generic Glucophage XR), in diabetic patients, as evidenced by the hemoglobin A1c level being above the patient's goal</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2.2 Submission of medical records (e.g. chart notes, laboratory values) documenting an intolerance to metformin extended-release (generic Glucophage XR) which is unable to be resolved with attempts to minimize the adverse effects where appropriate (e.g. dose reduction)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.3 History of greater than or equal to 12 week trial of metformin immediate-release</p> <p style="text-align: center;"><b>AND</b></p> <p>1.4 One of the following:</p> <p>1.4.1 Submission of medical records (e.g. chart notes, laboratory values) documenting an inadequate response to metformin immediate-release, in diabetic patients, as evidenced by the hemoglobin A1c level being above the patient's goal</p>	

**OR**

**1.4.2** Submission of medical records (e.g. chart notes, laboratory values) documenting an intolerance to metformin immediate-release which is unable to be resolved with attempts to minimize the adverse effects where appropriate (e.g. dose reduction)

Product Name: Brand Glumetza, Brand Fortamet

Approval Length | 12 month(s)

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - ALL of the following:

**1.1** History of greater than or equal to 12 week trial of metformin extended-release (generic Glucophage XR)

**AND**

**1.2** ONE of the following:

**1.2.1** Submission of medical records (e.g. chart notes, laboratory values) documenting an inadequate response to metformin extended-release (generic Glucophage XR), in diabetic patients, as evidenced by the hemoglobin A1c level being above the patient's goal

**OR**

**1.2.2** Submission of medical records (e.g. chart notes, laboratory values) documenting an intolerance to metformin extended-release (generic Glucophage XR) which is unable to be resolved with attempts to minimize the adverse effects where appropriate (e.g. dose reduction)

**AND**

**1.3** History of greater than or equal to 12 week trial of metformin extended-release (generic Fortamet)

**AND**

**1.4** One of the following:

**1.4.1** Submission of medical records (e.g. chart notes, laboratory values) documenting an inadequate response to metformin extended-release (generic Fortamet), in diabetic patients, as evidenced by the hemoglobin A1c level being above the patient's goal

**OR**

**1.4.2** Submission of medical records (e.g. chart notes, laboratory values) documenting an intolerance to metformin extended-release (generic Fortamet) which is unable to be resolved with attempts to minimize the adverse effects where appropriate (e.g. dose reduction)

**AND**

**1.5** History of greater than or equal to 12 week trial of metformin immediate-release

**AND**

**1.6** One of the following:

**1.6.1** Submission of medical records (e.g. chart notes, laboratory values) documenting an inadequate response to metformin immediate-release, in diabetic patients, as evidenced by the hemoglobin A1c level being above the patient's goal

**OR**

**1.6.2** Submission of medical records (e.g. chart notes, laboratory values) documenting an intolerance to metformin immediate-release which is unable to be resolved with attempts to minimize the adverse effects where appropriate (e.g. dose reduction)

**AND**

1.7 Submission of article(s) published in the peer-reviewed medical literature showing that the requested drug is likely to be more efficacious to this patient than metformin extended-release (generic Glucophage XR)

## 2 . Revision History

Date	Notes
10/13/2022	Removed Brand Glucophage XR as target

Migranal

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-133813
<b>Guideline Name</b>	Migranal
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2023
-----------------	-----------

### 1 . Criteria

Product Name:Brand Migranal, Generic dihydroergotamine mesylate	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of migraine headaches with or without aura  <b>AND</b>  2 - History of failure, contraindication, or intolerance to TWO preferred 5-HT1 (5-	

hydroxytryptamine-1) receptor agonist (triptan) alternatives [eg, Imitrex (sumatriptan), Maxalt or Maxalt-MLT (rizatriptan)]

## 2 . Revision History

Date	Notes
9/26/2023	Removed QL section, no QLs in place.

Miplyffa (arimoclomol)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-240207
<b>Guideline Name</b>	Miplyffa (arimoclomol)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/15/2025
-----------------	-----------

## 1 . Criteria

Product Name:Miplyffa	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes documenting a diagnosis of Niemann-Pick disease type C (NPC))	

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming diagnosis by one of the following:

**2.1** Genetically confirmed (deoxyribonucleic acid [DNA] sequence analysis) by mutations in both alleles of NPC1 or NPC2

**OR**

**2.2** Mutation in only one allele of NPC1 or NPC2 plus either positive filipin staining or elevated cholestane triol/oxysterols (>2 x upper limit of normal)

**AND**

**3** - Patient has at least one neurological symptom of the disease (e.g., hearing loss, vertical supranuclear gaze palsy, ataxia, dementia, dystonia, seizures, dysarthria, or dysphagia)

**AND**

**4** - Patient is 2 years of age or older

**AND**

**5** - Requested drug will be used in combination with miglustat

**AND**

**6** - Requested drug will NOT be used in combination with Aqneursa (levacetylleucine). \*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.

**AND**

7 - Prescribed by or in consultation with a specialist knowledgeable in the treatment of Niemann-Pick disease type C

Notes

\*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.

Product Name:Miplyffa

Approval Length

12 month(s)

Therapy Stage

Reauthorization

Guideline Type

Prior Authorization

**Approval Criteria**

1 - Patient demonstrates positive clinical response to therapy (e.g., slowing of disease progression, improvement in neurological symptoms of the disease)

**AND**

2 - Requested drug will be used in combination with miglustat

**AND**

3 - Requested drug will NOT be used in combination with Aqneursa (levacetylleucine). \*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.

Notes

\*The use of both Miplyffa and Aqneursa concomitantly is not covered due to lack of evidence to support the use of both products at the same time and the lack of a Miplyffa and Aqneursa head-to-head study.

**2 . Revision History**

Date

Notes

4/15/2025	Added verbiage regarding concomitant use with Aqneursa, no change to clinical criteria
-----------	--

Monurol

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-108624
<b>Guideline Name</b>	Monurol
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/23/2022
-----------------	-----------

### 1 . Criteria

Product Name: Monurol	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The provider has submitted labs showing the culture and sensitivity is positive for Monural and negative to Ciprofloxacin or Nitrofurantoin  <b>OR</b>	

**2** - Trial and failure, contraindication, or intolerance to ONE of the following:

- Ciprofloxacin
- Nitrofurantoin

## **2 . Revision History**

Date	Notes
6/23/2022	Added product name to criteria section, no change to criteria

Mozobil

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99625
<b>Guideline Name</b>	Mozobil
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Mozobil	
Approval Length	4 Days*
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following: <ul style="list-style-type: none"><li>Patients with non-Hodgkin's lymphoma (NHL) who will be undergoing autologous hematopoietic stem cell (HSC) transplantation</li><li>Patients with multiple myeloma (MM) who will be undergoing autologous HSC transplantation</li></ul>	

**AND**

**2** - Used in combination with granulocyte-colony stimulating factor (G-CSF) [e.g., Zarxio (filgrastim)]

**AND**

**3** - Prescribed by, or in consultation with, a hematologist/oncologist

Notes

\*Authorization will be issued for 1 course of therapy (up to four days of therapy).

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

MS Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157637
<b>Guideline Name</b>	MS Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name: PREFERRED: Avonex, Brand Copaxone, generic dalfampridine, generic dimethyl fumarate capsules, generic fingolimod, Kesimpta, Ocrevus, Ocrevus Zunovo, Rebif, generic teriflunomide	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of multiple sclerosis (MS)	

Product Name:NON-PREFERRED: Brand Ampyra, Brand Aubagio, Bafiertam, Betaseron, Briumvi, Extavia, Brand Gilenya, generic glatiramer acetate, Glatopa, Mavenclad, Mayzent, Plegridy, Ponvory, Tascenso ODT, Brand Tecfidera capsules, Brand Tecfidera starter packs, generic dimethyl fumarate starter packs, Vumerity

Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of multiple sclerosis (MS)

**AND**

2 - One of the following:

2.1 Patient has a history of failure, contraindication, or intolerance to a trial of at least TWO of the preferred alternatives \* (May require PA) (Verified via pharmacy paid claims or submission of medical records)

- Avonex
- Brand Copaxone
- generic dalfampridine
- generic dimethyl fumarate
- generic fingolimod
- Kesimpta
- Ocrevus/Ocrevus Zunovo
- Rebif
- generic teriflunomide
- Tysabri

**OR**

2.2 Patient is currently established on requested medication as documented by claims history or medical records (document drug, date, and duration of therapy)

Notes	* NOTE: Preferred Drug May Require PA
-------	---------------------------------------

Product Name:PREFERRED: Avonex, Brand Copaxone, generic dalfampridine, generic dimethyl fumarate, generic fingolimod, Kesimpta, Ocrevus, Ocrevus Zunovo, Rebif, generic

teriflunomide; NON-PREFERRED: Brand Ampyra, Brand Aubagio, Bafiertam, Betaseron, Briumvi, Extavia, Brand Gilenya, generic glatiramer acetate, Glatopa, Mavenclad, Mayzent, Plegridy, Ponvory, Tascenso ODT, Brand Tecfidera capsules, Brand Tecfidera starter packs, generic dimethyl fumarate starter packs, Vumerity	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression)</p>	

**2 . Revision History**

Date	Notes
10/25/2024	Added Ocrevus Zunovo as preferred target

Multaq

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99476
<b>Guideline Name</b>	Multaq
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name: Multaq	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 All of the following:</p> <p>1.1.1 Diagnosis of ONE of the following:</p> <ul style="list-style-type: none"><li>Paroxysmal Atrial Fibrillation (AF)</li></ul>	

- Persistent AF defined as AF less than 6 months duration

**AND**

**1.1.2** ONE of the following:

- Patient is in sinus rhythm
- Patient is planned to undergo cardioversion to sinus rhythm

**AND**

**1.1.3** Patient does not have New York Heart Association (NYHA) Class IV heart failure

**AND**

**1.1.4** Patient does not have symptomatic heart failure with recent decompensation requiring hospitalization

**OR**

**1.2** For continuation of current therapy

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Myalept

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99626
<b>Guideline Name</b>	Myalept
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Myalept	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of ONE of the following: <ul style="list-style-type: none"><li>Congenital generalized lipodystrophy associated with leptin deficiency</li><li>Acquired generalized lipodystrophy associated with leptin deficiency</li></ul>	

**AND**

**2** - Used as an adjunct to diet modification

**AND**

**3** - Prescribed by an endocrinologist

**AND**

**4** - Documentation demonstrates that patient has at least **ONE** of the following:

**4.1** Diabetes mellitus or insulin resistance with persistent hyperglycemia ( hemoglobin A1C greater than 7.0%) despite **BOTH** of the following:

- Dietary intervention
- Optimized insulin therapy at maximum tolerated doses

**OR**

**4.2** Persistent hypertriglyceridemia (triglycerides greater than 250 milligrams per deciliter) despite **BOTH** of the following:

- Dietary intervention
- Optimized therapy with at least two triglyceride-lowering agents from different classes (e.g., fibrates, statins) at maximum tolerated doses

Product Name: Myalept	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to Myalept therapy

**AND**

2 - Used as an adjunct to diet modification

**AND**

3 - Prescribed by an endocrinologist

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Myfembree (relugolix, estradiol, and norethindrone acetate), Oriahnn (elagolix, estradiol, and norethindrone acetate capsules; elagolix capsules)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-118559
<b>Guideline Name</b>	Myfembree (relugolix, estradiol, and norethindrone acetate), Oriahnn (elagolix, estradiol, and norethindrone acetate capsules; elagolix capsules)
<b>Formulary</b>	<ul style="list-style-type: none"> <li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li> </ul>

### Guideline Note:

Effective Date:	1/1/2023
-----------------	----------

## 1 . Criteria

Product Name: Oriahnn, Myfembree	
Diagnosis	Heavy Menstrual Bleeding Associated With Uterine Leiomyomas (Fibroids)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of heavy menstrual bleeding associated with uterine leiomyomas (fibroids)

**AND**

2 - Patient is premenopausal

**AND**

3 - One of the following:

3.1 History of inadequate control of bleeding following a trial of at least 3 months, or history of intolerance or contraindication to one of the following:

- Combination (estrogen/progestin) contraceptive
- Progestins
- Tranexamic acid

**OR**

3.2 Patient has had a previous interventional therapy to reduce bleeding

**AND**

4 - Treatment duration of therapy has not exceeded a total of 24 months

Product Name: Oriahnn, Myfembree	
Diagnosis	Heavy Menstrual Bleeding Associated With Uterine Leiomyomas (Fibroids)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Patient has improvement in bleeding associated with uterine leiomyomas (fibroids) (e.g., significant/sustained reduction in menstrual blood loss per cycle, improved quality of life, etc.)

**AND**

2 - Treatment duration of therapy has not exceeded a total of 24 months

Product Name: Myfembree	
Diagnosis	Pain Associated With Endometriosis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of moderate to severe pain associated with endometriosis

**AND**

2 - Patient is premenopausal

**AND**

3 - ONE of the following:

3.1 History of inadequate pain control response following a trial of 30 days, or history of intolerance or contraindication to one of the following:

- Danazol
- Combination (estrogen/progestin) contraceptive
- Progestins

**OR**

**3.2** Patient has had surgical ablation to prevent recurrence

**AND**

**4** - Treatment duration of Myfembree has not exceeded a total of 24 months

Product Name:Myfembree	
Diagnosis	Pain Associated With Endometriosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has improvement in pain associated with endometriosis (e.g., improvement in dysmenorrhea and nonmenstrual pelvic pain)	
<b>AND</b>	
2 - Treatment duration of Myfembree has not exceeded a total of 24 months	

## 2 . Revision History

Date	Notes
12/19/2022	New Program

Mytesi

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99477
<b>Guideline Name</b>	Mytesi
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Mytesi	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome (AIDS) associated diarrhea	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk copy C&S Arizona standard to Medicaid Arizona

Nadolol

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-109903
<b>Guideline Name</b>	Nadolol
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Nadolol	
Diagnosis	PA required for patients 18 years of age or older
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, contraindication, or intolerance to 3 of the following: <ul style="list-style-type: none"><li>atenolol</li><li>atenolol/chlorthalidone</li><li>bisoprolol fumarate</li></ul>	

- bisoprolol/hydrochlorothiazide
- carvedilol
- labetalol HCl
- metoprolol succinate
- metoprolol tartrate
- metoprolol/hydrochlorothiazide
- propranolol HCl
- propranolol/hydrochlorothiazide
- sotalol HCl

## 2 . Revision History

Date	Notes
7/28/2022	Updated indication verbiage, no change to clinical criteria.

Namzaric (memantine/donepezil)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164377
<b>Guideline Name</b>	Namzaric (memantine/donepezil)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Namzaric, generic memantine/donepezil	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - BOTH of the following:  1.1 History of BOTH of the following:  1.1.1 Memantine (generic Namenda)	

**AND**

**1.1.2** Donepezil (generic Aricept)

**AND**

**1.2** Patient is stabilized on 10mg of donepezil once daily

## **2 . Revision History**

Date	Notes
1/30/2025	Added GPIs for new generic memantine/donepezil as NP target.

Natpara

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99627
<b>Guideline Name</b>	Natpara
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Natpara	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ALL of the following:  1.1 Diagnosis of hypocalcemia resulting from chronic hypoparathyroidism	

**AND**

**1.2** 25-hydroxy vitamin D level is above the lower limit of the normal laboratory reference range

**AND**

**1.3** Patient is currently on active vitamin D (calcitriol) therapy

**AND**

**1.4** Total serum calcium level (albumin corrected) is above 7.5 milligrams per deciliter

**AND**

**2** - ONE of the following:

**2.1** Patient is currently on calcium supplementation of 1-2 grams per day of elemental calcium in divided doses

**OR**

**2.2** Patient has a contraindication to calcium supplementation

**AND**

**3** - Prescribed by ONE of the following:

- Endocrinologist
- Nephrologist

Product Name: Natpara	
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Total serum calcium level (albumin corrected) within the lower half of the normal range (approximately 8 to 9 milligrams per deciliter)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient continues to take concomitant calcium supplementation that is sufficient to meet daily requirements</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by ONE of the following:</p> <ul style="list-style-type: none"> <li>• Endocrinologist</li> <li>• Nephrologist</li> </ul>	

**2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Nemluvio (nemolizumab-ilto)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-193199
<b>Guideline Name</b>	Nemluvio (nemolizumab-ilto)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Nemluvio	
Diagnosis	Prurigo Nodularis (PN)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming diagnosis of prurigo nodularis (PN)	

**AND**

**2** - Patient has at least 20 nodular lesions

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist
- Immunologist

**AND**

**4** - Submission of medical records (e.g., chart notes) or paid claims confirming trial and failure, contraindication, or intolerance to BOTH of the following:\*

- One medium or higher potency topical corticosteroid
- Dupixent (dupilumab) (PA may be required)

Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication
-------	--

Product Name:Nemluvio	
Diagnosis	Prurigo Nodularis (PN)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> <b>1</b> - Submission of medical records (e.g., chart notes) documenting that patient demonstrates a positive clinical response to therapy as evidenced by at least ONE of the following: <ul style="list-style-type: none"><li>• Reduction in the number of nodular lesions from baseline</li></ul>	

- Improvement in symptoms (e.g., pruritus, inflammation) from baseline

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist
- Immunologist

Product Name:Nemluvio	
Diagnosis	Atopic Dermatitis (AD)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming diagnosis of moderate to severe atopic dermatitis (AD)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - One of the following:</p> <ul style="list-style-type: none"> <li>• Involvement of at least 10% body surface area (BSA)</li> <li>• SCORing Atopic Dermatitis (SCORAD) index value of at least 25</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Patient is 12 years of age or older</p> <p style="text-align: center;"><b>AND</b></p>	

**4** - Will be used in combination with topical corticosteroids and/or calcineurin inhibitors (e.g., tacrolimus ointment)

**AND**

**5** - Submission of medical records (e.g., chart notes) or paid claims confirming history of failure, contraindication, or intolerance to ALL of the following:\*

- One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]
- Eucrisa (crisaborole)
- Adbry (tralokinumab-ldrm)
- Opzelura (ruxolitinib)

**AND**

**6** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist
- Immunologist

Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and/or contraindication to medication
-------	--

Product Name:Nemluvio	
Diagnosis	Atopic Dermatitis (AD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes) documenting that patient demonstrates a positive clinical response to therapy as evidenced by at least ONE of the following:	
<ul style="list-style-type: none"><li>• Reduction in body surface area involvement from baseline</li></ul>	

- Reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline

**AND**

**2 - Prescribed by or in consultation with one of the following:**

- Dermatologist
- Allergist
- Immunologist

## 2 . Background

Benefit/Coverage/Program Information			
Table 1: Relative potencies of topical corticosteroids			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05

	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream	0.1
	Triamcinolone acetonide	Cream, ointment	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

### 3 . Revision History

Date	Notes
2/25/2025	Added criteria for new AD indication

Nexiclon XR (clonidine ER)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-125301
<b>Guideline Name</b>	Nexiclon XR (clonidine ER)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Nexiclon XR, Brand Clonidine ER (Nexiclon XR ABA)	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Requested medication is being used for treatment of hypertension  <b>AND</b>	

**2** - Trial and failure, contraindication, or intolerance to one of the following (verified via paid pharmacy claims or submitted chart notes):

- generic clonidine oral tablet
- generic clonidine topical patch

## **2 . Revision History**

Date	Notes
5/25/2023	Added Nexiclon XR ABA as NP target. Updated dx criterion.

Nexletol (bempedoic acid) and Nexlizet (bempedoic acid-ezetimibe)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147690
<b>Guideline Name</b>	Nexletol (bempedoic acid) and Nexlizet (bempedoic acid-ezetimibe)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Nexletol, Nexlizet	
Diagnosis	Heterozygous familial hypercholesterolemia (HeFH) or primary hyperlipidemia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following diagnoses: <ul style="list-style-type: none"><li>Heterozygous familial hypercholesterolemia (HeFH)</li></ul>	

- Primary Hyperlipidemia

**AND**

**2** - One of the following:

- Patient has been receiving at least 12 consecutive weeks of highest tolerable dose of statin therapy
- Patient is statin intolerant as evidenced by an inability to tolerate at least two statins, with at least one started at the lowest starting daily dose, due to intolerable symptoms or clinically significant biomarker changes of liver function or muscle function (e.g., creatine kinase)
- Patient has an FDA labeled contraindication to all statins

**AND**

**3** - One of the following LDL-C values while on maximally tolerated statin therapy within the last 120 days:

- LDL-C greater than or equal to 55 mg/dL with ASCVD
- LDL-C greater than or equal to 100 mg/dL without ASCVD

**AND**

**4** - One of the following:

**4.1** For Nexletol, ONE of the following:

- Patient has been receiving at least 12 consecutive weeks of generic ezetimibe therapy as adjunct to maximally tolerated statin therapy
- Patient has a history of contraindication or intolerance to ezetimibe

**OR**

**4.2** For Nexlizet, patient has been receiving at least 12 consecutive weeks of generic ezetimibe therapy as adjunct to maximally tolerated statin therapy

Product Name: Nexletol, Nexlizet

Diagnosis	Heterozygous familial hypercholesterolemia (HeFH) or primary hyperlipidemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy as evidenced by a reduction in LDL-C levels from baseline while on therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <ul style="list-style-type: none"> <li>• Patient continues to receive other lipid-lowering therapy (e.g., statins, ezetimibe) at the maximally tolerated dose</li> <li>• Patient has a documented inability to take other lipid-lowering therapy (e.g., statins, ezetimibe)</li> </ul>	

Product Name: Nexletol, Nexlizet	
Diagnosis	Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Established cardiovascular disease (CVD) (e.g., coronary artery disease, symptomatic peripheral arterial disease, cerebrovascular atherosclerotic disease)</li> </ul>	

- A high risk for a CVD event but without established CVD [e.g., diabetes mellitus (type 1 or type 2) in females over 65 years of age or males over 60 years of age]

**AND**

**2 - One of the following:**

- Patient is statin intolerant as evidenced by an inability to tolerate at least two statins, with at least one started at the lowest starting daily dose, due to intolerable symptoms or clinically significant biomarker changes of liver function or muscle function (e.g., creatine kinase)
- Patient has an FDA labeled contraindication to all statins

**AND**

**3 - One of the following LDL-C values within the last 120 days:**

- LDL-C greater than or equal to 55 mg/dL with ASCVD
- LDL-C greater than or equal to 100 mg/dL without ASCVD

**AND**

**4 - One of the following:**

**4.1 For Nexletol, ONE of the following:**

- Patient has been receiving at least 12 consecutive weeks of generic ezetimibe therapy
- Patient has a history of contraindication or intolerance to ezetimibe

**OR**

**4.2 For Nexlizet, patient has been receiving at least 12 consecutive weeks of generic ezetimibe therapy**

Product Name:Nexletol, Nexlizet	
Diagnosis	Established cardiovascular disease (CVD) or high risk for a CVD event but without established CVD
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
5/23/2024	Updated criteria due to indication updates in prescribing information. Added criteria for the new FDA approved indication.

Niktimvo (axatilimab-csfr)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-226188
<b>Guideline Name</b>	Niktimvo (axatilimab-csfr)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Niktimvo	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of chronic graft versus host disease (cGVHD)	

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure or inadequate response to at least two other systemic therapies [e.g., corticosteroids (e.g., prednisone, methylprednisolone), mycophenolate]

**AND**

**3** - Patient weighs at least 40kg

Product Name:Niktimvo	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting patient does not show evidence of progressive disease while on therapy	

## 2 . Revision History

Date	Notes
3/26/2025	New program

Nityr- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99628
<b>Guideline Name</b>	Nityr- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Nityr	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of hereditary tyrosinemia type 1  <b>AND</b>	

**2** - Prescriber provides a reason or special circumstance the patient cannot use Orfadin (nitisinone) capsules or suspension

Product Name:Nityr

Approval Length 12 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Patient shows evidence of positive clinical response (e.g. decrease in urinary/plasma succinylacetone and alpha-1-microglobulin levels) while on Nityr therapy

**2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Nocdurna

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99505
<b>Guideline Name</b>	Nocdurna
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Nocdurna	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of nocturia due to nocturnal polyuria (as defined by nighttime urine production that exceeds one-third of the 24-hour urine production)	

**AND**

**2** - Patient wakes at least twice per night on a reoccurring basis to void

**AND**

**3** - Documented serum sodium level is currently within normal limits of the normal laboratory reference range and has been within normal limits over the previous six months

**AND**

**4** - The patient has been evaluated for other medical causes and has either not responded to, tolerated, or has a contraindication to treatments for identifiable medical causes [e.g., overactive bladder, benign prostatic hyperplasia/lower urinary tract symptoms (BPH/LUTS), elevated post-void residual urine, and heart failure]

**AND**

**5** - Prescriber attests that the risks have been assessed and benefits outweigh the risks

Product Name:Nocdurna	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Documentation of positive clinical response to therapy	
<b>AND</b>	
<b>2</b> - Patient has routine monitoring for serum sodium levels	

**AND**

**3** - Prescriber attests that the risks of hyponatremia have been assessed and benefits outweigh the risks

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk copy from C&S Medicaid to Arizona Medicaid for 7/1 eff

Non-Preferred Drugs - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-223223
<b>Guideline Name</b>	Non-Preferred Drugs - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Non-Preferred Drugs	
Approval Length	12 months*
Guideline Type	Administrative
<b>Approval Criteria</b>  1 - ALL of the following:  1.1 ONE of the following: <ul style="list-style-type: none"><li>If there are at least three preferred alternatives, history of trial per member's pharmacy claims resulting in a therapeutic failure, contraindication, or intolerance to at least THREE preferred alternatives (Prior trials of formulary/preferred drug list (PDL)</li></ul>	

alternatives must sufficiently demonstrate that the formulary/PDL alternatives are either ineffective or inappropriate at the time of the request)\*

- If there are fewer than three preferred alternatives, the patient must have a history of trial per member's pharmacy claims resulting in a therapeutic failure, contraindication, or intolerance to ALL of the preferred products (Prior trials of formulary/preferred drug list (PDL) alternatives must sufficiently demonstrate that the formulary/PDL alternatives are either ineffective or inappropriate at the time of the request)\*
- There are no preferred formulary alternatives for the requested drug\*

**AND**

**1.2** If the request is for a multi-source brand medication (i.e., MSC O) ONE of the following:

**1.2.1** BOTH of the following:

- The brand is being requested because of an adverse reaction, allergy or sensitivity to the generic and the prescriber must attest to submitting the FDA MedWatch Form for allergic reactions to the medications
- The patient has tried three generic formulations (if available) of the requested brand product

**OR**

**1.2.2** ONE of the following:

- The brand is being requested due to a therapeutic failure with the generic (please provide reason for therapeutic failure)
- The brand is being requested because transition to the generic could result in destabilization of the patient (rationale must be provided)
- Special clinical circumstances exist that preclude the use of the generic equivalent of the multi-source brand medication for the patient (rationale must be provided)

**AND**

**1.3** ONE of the following:

**1.3.1** The requested drug must be used for a Food and Drug Administration (FDA)-approved indication

**OR**

**1.3.2** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- The requested drug must be used for a Food and Drug Administration (FDA)-approved indication
- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmacoeconomic studies
- Other drug reference resources

**AND**

**1.4** The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plan's program\*\*

**OR**

**2** - If the requested medication is a behavioral health medication, ONE of the following:

- The patient has been receiving treatment with the requested non-preferred behavioral health medication and is new to the plan (enrollment effective date within the past 90 days)
- The patient is currently receiving treatment with the requested non-preferred behavioral health medication in the hospital and must continue upon discharge

Notes

\*Anti-infectives: Approve for the requested time frame, or if duration is not specified approve the request for 30 days.

\*Controlled Substances shall be approved for the requested time. If there is not a requested time period and it is not clear in the directions, approve for one time only.

\*Other medications: Approved for the requested time frame, or if duration is not specified, approve for 12 months.

	<p>* For Non-Preferred Generics (i.e. MSC=Y) approvals: Please approve at MSC=Y only.</p> <p>For preferred alternatives, use the non-preferred alternatives grid to identify appropriate alternatives: <a href="https://uhgazure.sharepoint.com/sites/CST/CSDM/Shared%20Documents/Forms/AllItems.aspx?FolderCTID=0x01200027C80175A8369D44AC45A99A99328B80&amp;View=%7B4B6D25AD%2D6A95%2D496D%2D9937%2D65CECD43AFE7%7D&amp;viewid=c2ad0afa%2D814c%2D499e%2Dbf25%2D3411fac9171f&amp;id=%2Fsites%2FCST%2FCSDM%2FShared%20Documents%2FAZM%2FN%20Alt%20Tables">https://uhgazure.sharepoint.com/sites/CST/CSDM/Shared%20Documents/Forms/AllItems.aspx?FolderCTID=0x01200027C80175A8369D44AC45A99A99328B80&amp;View=%7B4B6D25AD%2D6A95%2D496D%2D9937%2D65CECD43AFE7%7D&amp;viewid=c2ad0afa%2D814c%2D499e%2Dbf25%2D3411fac9171f&amp;id=%2Fsites%2FCST%2FCSDM%2FShared%20Documents%2FAZM%2FN%20Alt%20Tables</a></p> <p>**Medications used to treat obesity/weight loss, cosmetic (e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, or sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs (Cialis/Tadalafil) are covered for clinical diagnoses other than ED.</p>
--	---

**2 . Revision History**

Date	Notes
3/26/2025	Updated note section regarding cosmetic/excluded use.

Non-Preferred Prenatal Vitamins

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-99528
<b>Guideline Name</b>	Non-Preferred Prenatal Vitamins
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Non-Preferred Prenatal Vitamins	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, contraindication, or intolerance to ALL of the following preferred products:*	
Notes	*Please refer to the background table for the alternatives

## 2 . Background

Benefit/Coverage/Program Information			
Preferred Products:			
GPI-14	Product ID	Product Label	GPI-14 Description
785120000003 15	7331710500 9	PRENATVITE TA B RX	*PRENATAL MULTIVITAMINS & MINERALS W/IRON & FA TAB 0.8 MG***
785120100003 30	6954302679 0	PNV TABS TAB 29-1MG	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	6025801930 9	PRENATABS RX TAB	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	4293707051 0	PRENATAL+FE T AB 29-1MG	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	4293707051 6	PRENATAL+FE T AB 29-1MG	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	4293707051 8	PRENATAL+FE T AB 29-1MG	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	5865701339 0	THRIVITE RX TAB 29-1MG	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	7118600192 4	VIL-RX TAB 29-1MG	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 30	1381105169 0	VOL-TAB RX TAB	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 29-1 MG***
785120100003 52	1381100271 0	ELITE-OB TAB	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 50-1.25 MG***
785120100003 52	6802500101 0	OB COMPLETE TAB	*PRENATAL VIT W/ IRON CARBONYL-FA TAB 50-1.25 MG***
785120150003 24	5865701700 1	M-NATAL PLUS TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	1283008000 1	M-VIT TAB 27- 1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***

785120150003 24	7089802200 1	NEONATAL TAB COMPLTE	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	7089801150 1	NEONATAL PLS TAB 27-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	7583400500 1	NIVA-PLUS TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	0081393160 1	O-CAL FA TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	7139962460 9	ONE VITE TAB 1MG PLUS	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	3932801061 0	PRENATAL TAB 27-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	3932801065 0	PRENATAL TAB 27-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	6304401500 1	PRENATAL VIT TAB LOW IRON	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	6304401500 5	PRENATAL VIT TAB LOW IRON	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	6954302581 0	PREPLUS TAB 27-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	6954302585 0	PREPLUS TAB 27-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	6711201010 0	TRICARE TAB PRENATAL	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	1713908003 0	VITATHELY TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	1381105191 0	VOL-PLUS TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	1381105195 0	VOL-PLUS TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***
785120150003 24	6936702670 1	WESTAB PLUS TAB 27- 1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 27-1 MG***

785120150003 29	6025801920 1	TRINATE TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 28-1 MG***
785120150003 29	1381105141 0	VOL-NATE TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 28-1 MG***
785120150003 32	1026722700 1	CO-NATAL FA TAB 29-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 29-1 MG***
785120150003 32	7331782860 1	NEONATAL TAB COMPLETE	*PRENATAL VIT W/ FE FUMARATE-FA TAB 29-1 MG***
785120150003 32	6954302591 0	PRETAB TAB 29-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 29-1 MG***
785120150003 60	1381100071 0	TRINATAL RX TAB 1	*PRENATAL VIT W/ FE FUMARATE-FA TAB 60-1 MG***
785120150003 60	5199105660 1	VINATE ONE TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 60-1 MG***
785120150003 66	5860708112 0	MYNATAL PLUS TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 65-1 MG***
785120150003 66	5860701056 5	MYNATAL-Z TAB	*PRENATAL VIT W/ FE FUMARATE-FA TAB 65-1 MG***
785120150003 66	0064200791 2	VITAFOL-OB TAB 65-1MG	*PRENATAL VIT W/ FE FUMARATE-FA TAB 65-1 MG***
785120150005 30	1381100149 0	COMPLETENATE CHW	*PRENATAL VIT W/ FE FUMARATE-FA CHEW TAB 29-1 MG***
785120150005 30	4293707071 0	PRENATAL 19 CHW 29-1MG	*PRENATAL VIT W/ FE FUMARATE-FA CHEW TAB 29-1 MG***
785120150005 30	4293707071 6	PRENATAL 19 CHW 29-1MG	*PRENATAL VIT W/ FE FUMARATE-FA CHEW TAB 29-1 MG***
785120150005 30	4293707071 8	PRENATAL 19 CHW 29-1MG	*PRENATAL VIT W/ FE FUMARATE-FA CHEW TAB 29-1 MG***

785120150005 30	6025801970 1	PRENATAL 19 CHW TAB	*PRENATAL VIT W/ FE FUMARATE-FA CHEW TAB 29-1 MG***
785120150005 30	1392501170 1	SE-NATAL 19 CHW	*PRENATAL VIT W/ FE FUMARATE-FA CHEW TAB 29-1 MG***
785120160001 30	1381100493 0	ULTIMATECARE CAP ONE	*PRENATAL VIT W/ FE CBN-FE ASP GLYC-FA-OMEGA 3 CAP 27- 1MG***
7851201800011 6	2335901053 0	C-NATE DHA CAP 28-1- 200	*PRENATAL VIT W/ FE FUM-FA- OMEGA 3 CAP 28-1-200 MG***
7851201800011 6	2335902003 0	RELNATE DHA CAP	*PRENATAL VIT W/ FE FUM-FA- OMEGA 3 CAP 28-1-200 MG***
7851201800011 6	6954303703 0	VIRT-NATE CAP DHA	*PRENATAL VIT W/ FE FUM-FA- OMEGA 3 CAP 28-1-200 MG***
7851201800011 6	6466100803 0	VIVA DHA CAP	*PRENATAL VIT W/ FE FUM-FA- OMEGA 3 CAP 28-1-200 MG***
785120220003 20	6954302419 0	VIRT-PN TAB	*PRENATAL VIT W/ FE FUM- METHYLFOLATE-FA TAB 27-0.6- 0.4 MG***
785120460003 30	5549501250 1	ATABEX OB TAB 29-1MG	*PRENATAL VIT W/ FE BISGLYCINATE CHELATE-FA TAB 29-1 MG***
785120460003 30	5199101780 1	VINATE II TAB	*PRENATAL VIT W/ FE BISGLYCINATE CHELATE-FA TAB 29-1 MG***
785120510003 27	0017808589 0	CITRANATAL TA B RX	*PRENATAL W/O A W/ FE CARBONYL-FE GLUC-DSS-FA TAB 27-1MG***
785120580001 50	5274706203 0	CONCEPT OB CAP	*PRENATAL W/O A W/FE FUM-FE POLY-FA CAP 130-92.4-1 MG***
785120580001 50	1381105353 0	FOLIVANE- OB CAP	*PRENATAL W/O A W/FE FUM-FE POLY-FA CAP 130-92.4-1 MG***

785120600003 25	5199101550 1	VINATE M TAB	*PRENATAL VIT W/ SEL-FE FUMARATE-FA TAB 27-1 MG***
785120700003 30	4293707061 0	PRENATAL 19 TAB 29-1MG	*PRENATAL VIT W/ DSS-FE FUMARATE-FA TAB 29-1 MG***
785120700003 30	4293707061 6	PRENATAL 19 TAB 29-1MG	*PRENATAL VIT W/ DSS-FE FUMARATE-FA TAB 29-1 MG***
785120700003 30	4293707061 8	PRENATAL 19 TAB 29-1MG	*PRENATAL VIT W/ DSS-FE FUMARATE-FA TAB 29-1 MG***
785120700003 30	1392501160 1	SE-NATAL 19 TAB	*PRENATAL VIT W/ DSS-FE FUMARATE-FA TAB 29-1 MG***
785120910001 35	5274706213 0	CONCEPT DHA CAP	*PRENATAL W/FE FUM-FE POLY - FA-OMEGA 3 CAP 53.5-38-1 MG***
785120910001 35	5865701213 0	DOTHELLE DHA CAP	*PRENATAL W/FE FUM-FE POLY - FA-OMEGA 3 CAP 53.5-38-1 MG***
785120910001 35	1381105363 0	TARON-C DHA CAP	*PRENATAL W/FE FUM-FE POLY - FA-OMEGA 3 CAP 53.5-38-1 MG***
785120910001 35	7643903313 0	VIRT-C DHA CAP	*PRENATAL W/FE FUM-FE POLY - FA-OMEGA 3 CAP 53.5-38-1 MG***
785160200063 30	0064200763 0	VITAFOL-OB PAK +DHA	*PRENATAL MV W/FE FUM-FA TAB 65-1 MG & DHA CAP 250 MG PACK *
785160320001 30	0064200703 0	VITAFOL- ONE CAP	*PRENATAL MV W/ FE POLYSAC CMLX-FA-DHA CAP 29-1-200 MG***
785160320063 25	0064200753 0	SELECT- OB+ PAK DHA	*PRENATAL MV W/FE POLY-FA CHW 29-1 MG & DHA CAP 250 MG PAK *

### 3 . Revision History

Date	Notes
5/18/2021	Arizona Medicaid 7.1 Implementation

Northera

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99629
<b>Guideline Name</b>	Northera
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Northera	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) as defined by ONE of the following when an upright position is assumed or when using a head-up tilt-table testing at an angle of at least 60 degrees: <ul style="list-style-type: none"><li>At least a 20 millimeters of mercury (mm Hg) fall in systolic pressure</li></ul>	

- At least a 10 mm Hg fall in diastolic pressure

**AND**

**2** - nOH caused by ONE of the following:

- Primary autonomic failure (e.g., Parkinson's disease, multiple system atrophy, and pure autonomic failure)
- Dopamine beta-hydroxylase deficiency
- Non-diabetic autonomic neuropathy

**AND**

**3** - Diagnostic evaluation has excluded other causes associated with orthostatic hypotension (e.g., congestive heart failure, fluid restriction, malignancy)

**AND**

**4** - The patient has tried at least TWO of the following non-pharmacologic interventions:

- Discontinuation of drugs which can cause orthostatic hypotension [e.g., diuretics, antihypertensive medications (primarily sympathetic blockers), anti-anginal drugs (nitrates), alpha-adrenergic antagonists, and antidepressants]
- Raising the head of the bed 10 to 20 degrees
- Compression garments to the lower extremities or abdomen
- Physical maneuvers to improve venous return (e.g., regular modest-intensity exercise)
- Increased salt and water intake, if appropriate
- Avoiding precipitating factors (e.g., overexertion in hot weather, arising too quickly from supine to sitting or standing)

**AND**

**5** - No previous diagnosis of supine hypertension

**AND**

**6** - Prescribed by, or in consultation with, ONE of the following specialists:

- Cardiologist

- Neurologist
- Nephrologist

**AND**

**7** - History of failure (after a trial of at least 30 days), contraindication or intolerance to BOTH of the following medications:

- Florinef (fludrocortisone)
- ProAmatine (midodrine)

Product Name:Northera	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Documentation of positive clinical response to Northera therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Physiological countermeasures for neurogenic orthostatic hypotension (nOH) continue to be employed</p>	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Nourianz

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99481
<b>Guideline Name</b>	Nourianz
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Nourianz	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Parkinson's disease  <b>AND</b>	

**2** - Used as adjunctive treatment to levodopa/carbidopa in patients experiencing “off” episodes

**AND**

**3** - History of failure, contraindication, or intolerance to TWO anti-Parkinson’s disease therapies from the following adjunctive pharmacotherapy classes (trial must be from two different classes):

- Dopamine agonists (e.g., pramipexole, ropinirole)
- Catechol-O-methyl transferase (COMT) inhibitors (e.g., entacapone)
- Monoamine oxidase (MAO) B inhibitors (e.g., rasagiline, selegiline)

Product Name:Nourianz	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Nourianz therapy	
<b>AND</b>	
2 - Patient will continue to receive treatment with a carbidopa/levodopa-containing medication	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Nucala (mepolizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-141161
<b>Guideline Name</b>	Nucala (mepolizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/7/2024
-----------------	----------

## 1 . Criteria

Product Name:Nucala	
Diagnosis	Severe Asthma
Approval Length	6 Months [G]
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of severe asthma	

**AND**

**2** - Asthma is an eosinophilic phenotype as defined by one of the following:

- Baseline (pre-treatment) peripheral blood eosinophil level is greater than or equal to 150 cells/microliter
- Peripheral blood eosinophil levels were greater than or equal to 300 cells/microliter within the past 12 months

**AND**

**3** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting one of the following:

**3.1** Patient has had at least two or more asthma exacerbations requiring systemic corticosteroids (e.g., prednisone) within the past 12 months

**OR**

**3.2** Prior asthma-related hospitalization within the past 12 months

**AND**

**4** - Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications (verified via paid pharmacy claims):

**4.1** Both of the following:

- High-dose inhaled corticosteroid (ICS) (e.g., greater than 500 mcg fluticasone propionate equivalent/day)
- Additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium)

**OR**

**4.2** One maximally-dosed combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Symbicort [budesonide/formoterol], Breo Ellipta [fluticasone/vilanterol])

**AND**

**5** - Age greater than or equal to 6 years

**AND**

**6** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Allergist/Immunologist

Product Name:Nucala	
Diagnosis	Severe Asthma
Approval Length	12 Months
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications) [C]</p> <p><b>AND</b></p> <p><b>2</b> - Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications (verified via paid pharmacy claims)</p> <p><b>AND</b></p>	

**3** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Allergist/Immunologist

Product Name:Nucala	
Diagnosis	Chronic rhinosinusitis with nasal polyps (CRSwNP)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient is 18 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of documentation (e.g., chart notes) confirming ONE of the following:</p> <p>2.1 ALL of the following:</p> <p>2.1.1 Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) defined by ALL of the following:</p> <p>2.1.1.1 TWO or more of the following symptoms for greater than or equal to 12 weeks duration:</p> <ul style="list-style-type: none"><li>• Mucopurulent discharge</li><li>• Nasal obstruction and congestion</li><li>• Decreased or absent sense of smell</li><li>• Facial pressure or pain</li></ul> <p style="text-align: center;"><b>AND</b></p> <p>2.1.1.2 ONE of the following:</p>	

- Evidence of inflammation on paranasal sinus examination or computed tomography (CT)
- Evidence of purulence coming from paranasal sinuses or ostiomeatal complex

**AND**

**2.1.1.3** The presence of nasal polyps

**AND**

**2.1.2** ONE of the following:

- Patient has required prior sino-nasal surgery
- Patient has required systemic corticosteroids in the previous 2 years

**AND**

**2.1.3** Patient has been unable to obtain symptom relief after trial of ALL of the following agents/classes of agents:

- Nasal saline irrigations
- Intranasal corticosteroids (e.g. fluticasone, mometasone, triamcinolone, etc.)
- Antileukotriene agents (e.g. montelukast, zafirlukast, zileuton)

**OR**

**2.2** ALL of the following:

**2.2.1** Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP)

**AND**

**2.2.2** Patient is currently on Nucala therapy

**AND**

**3** - Patient will receive Nucala as add-on maintenance therapy in combination with intranasal corticosteroids

**AND**

**4** - Patient is NOT receiving Nucala in combination with another biologic medication [e.g., Xolair (omalizumab), Dupixent (dupilumab)]

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Otolaryngologist
- Allergist
- Pulmonologist

Product Name:Nucala	
Diagnosis	Chronic rhinosinusitis with nasal polyps (CRSwNP)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of documentation (e.g., chart notes) confirming positive clinical response to Nucala therapy	
<b>AND</b>	
2 - Patient will continue to receive Nucala as add-on maintenance therapy in combination with intranasal corticosteroids	

**AND**

**3** - Patient is NOT receiving Nucala in combination with another biologic medication [e.g., Xolair (omalizumab), Dupixent (dupilumab)]

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Otolaryngologist
- Allergist
- Pulmonologist

Product Name:Nucala	
Diagnosis	Eosinophilic Granulomatosis with Polyangiitis (EGPA)
Approval Length	12 Months
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA)	
<b>AND</b>	
<b>2</b> - Patient's disease has relapsed or is refractory to standard of care therapy (i.e., corticosteroid treatment with or without immunosuppressive therapy)	
<b>AND</b>	
<b>3</b> - Patient is currently receiving corticosteroid therapy (e.g., prednisolone, prednisone)	

**AND**

4 - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Rheumatologist
- Allergist/Immunologist

Product Name:Nucala	
Diagnosis	Eosinophilic Granulomatosis with Polyangiitis (EGPA)
Approval Length	12 Months
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (e.g., increase in remission time)	

Product Name:Nucala	
Diagnosis	Hypereosinophilic Syndrome (HES)
Approval Length	12 Months
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of hypereosinophilic syndrome (HES)	

**AND**

**2** - Patient has been diagnosed for at least 6 months

**AND**

**3** - Verification that other non-hematologic secondary causes have been ruled out (e.g., drug hypersensitivity, parasitic helminth infection, HIV infection, non-hematologic malignancy)

**AND**

**4** - Patient is Fip1-like1-platelet-derived growth factor receptor alpha (FIP1L1-PDGFR $\alpha$ )-negative

**AND**

**5** - Patient has uncontrolled HES defined as both of the following:

- History of 2 or more flares within the past 12 months [I]
- Pre-treatment blood eosinophil count greater than or equal to 1000 cells/microliter

**AND**

**6** - Trial and failure, contraindication, or intolerance to one of the following:

- Corticosteroid therapy (e.g., prednisone)
- Cytotoxic/immunosuppressive therapy (e.g., hydroxyurea, cyclosporine, imatinib)

**AND**

**7** - Prescribed by or in consultation with one of the following:

- Allergist/Immunologist
- Hematologist

Product Name:Nucala	
Diagnosis	Hypereosinophilic Syndrome (HES)
Approval Length	12 Months
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (e.g., reduction in flares, decreased blood eosinophil count, reduction in corticosteroid dose)</p>	

## 2 . Background

<b>Clinical Practice Guidelines</b>			
<p><b>The Global Initiative for Asthma Global Strategy for Asthma Management and Prevention: Table 1. Low, medium and high daily doses of inhaled corticosteroids in adolescents and adults 12 years and older [6]</b></p>			
Inhaled corticosteroid	Total Daily ICS Dose (mcg)		
	Low	Medium	High
Beclometasone dipropionate (pMDI, standard particle, HFA)	200-500	> 500-1000	> 1000
Beclometasone dipropionate (pMDI, extrafine particle*, HFA)	100-200	> 200-400	> 400
Budesonide (DPI)	200-400	> 400-800	> 800
Ciclesonide (pMDI, extrafine particle*, HFA)	80-160	> 160-320	> 320
Fluticasone furoate (DPI)	100		200

Fluticasone propionate (DPI)	100-250	> 250-500	> 500
Fluticasone propionate (pMDI, standard particle, HFA)	100-250	> 250-500	> 500
Mometasone furoate (DPI)	200		400
Mometasone furoate (pMDI, standard particle, HFA)	200-400		> 400
<p>DPI: dry powder inhaler; HFA: hydrofluoroalkane propellant; ICS: inhaled corticosteroid; N/A: not applicable; pMDI: pressurized metered dose inhaler (non-chlorofluorocarbon formulations); ICS by pMDI should be preferably used with a spacer *See product information.</p> <p><b><i>This is not a table of equivalence</i></b>, but instead, suggested total daily doses for the 'low', 'medium' and 'high' dose ICS options for adults/adolescents, based on available studies and product information. Data on comparative potency are not readily available and therefore this table does NOT imply potency equivalence. Doses may be country -specific depending on local availability, regulatory labelling and clinical guidelines.</p> <p>For new preparations, including generic ICS, the manufacturer's information should be reviewed carefully; products containing the same molecule may not be clinically equivalent.</p>			

### 3 . Revision History

Date	Notes
2/6/2024	Updated CRSwNP criteria to align with Dupixent criteria.

Nuedexta

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99482
<b>Guideline Name</b>	Nuedexta
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Nuedexta	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of pseudobulbar affect (PBA)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Nuplazid

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99483
<b>Guideline Name</b>	Nuplazid
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Nuplazid	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Parkinson's disease  <b>AND</b>	

2 - Patient is currently experiencing hallucinations and delusions associated with Parkinson's disease psychosis (i.e., hallucination and delusion symptoms started after Parkinson's disease diagnosis)

Product Name:Nuplazid	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Nuplazid therapy</p>	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Nuzyra

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99522
<b>Guideline Name</b>	Nuzyra
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Nuzyra	
Diagnosis	Community-Acquired Bacterial Pneumonia
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 For continuation of therapy upon hospital discharge	

**OR**

**1.2** As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

**1.3** ALL of the following:

**1.3.1** Diagnosis of community-acquired bacterial pneumonia (CABP)

**AND**

**1.3.2** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Nuzyra

**AND**

**1.3.3** History of failure, contraindication, or intolerance to THREE of the following antibiotics or antibiotic regimens:

- Amoxicillin
- A macrolide
- Doxycycline
- A fluoroquinolone
- Combination therapy with amoxicillin/clavulanate or cephalosporin AND a macrolide or doxycycline

Product Name:Nuzyra	
Diagnosis	Acute Bacterial Skin and Skin Structure Infections
Approval Length	14 Day(s)
Guideline Type	Prior Authorization

## **Approval Criteria**

1 - ONE of the following:

1.1 For continuation of therapy upon hospital discharge

**OR**

1.2 As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

1.3 ALL of the following:

1.3.1 ONE of the following diagnoses:

1.3.1.1 BOTH of the following:

- Acute bacterial skin and skin structure infections
- Infection caused by methicillin-resistant *Staphylococcus aureus* (MRSA) documented by culture and sensitivity report

**OR**

1.3.1.2 BOTH of the following:

- Empirical treatment of patients with acute bacterial skin and skin structure infections
- Presence of MRSA infection is likely

**AND**

1.3.2 History of failure, contraindication, or intolerance to linezolid (generic Zyvox)

**AND**

1.3.3 History of failure, contraindication, or intolerance to ONE of the following antibiotics:

- Sulfamethoxazole-trimethoprim (SMZ-TMP)

- A tetracycline
- Clindamycin

**OR**

**1.4 ALL of the following:**

**1.4.1** Diagnosis of acute bacterial skin and skin structure infections

**AND**

**1.4.2** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Nuzyra

**AND**

**1.4.3** History of failure, contraindication, or intolerance to THREE of the following antibiotics:

- A penicillin
- A cephalosporin
- A tetracycline
- Sulfamethoxazole-trimethoprim (SMZ-TMP)
- Clindamycin

Product Name:Nuzyra	
Diagnosis	Off-Label Uses*
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - ONE of the following:</b></p> <p><b>1.1</b> For continuation of therapy upon hospital discharge</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

**1.3** The medication is being prescribed by or in consultation with an infectious disease specialist.

Notes

\*Note: Authorization duration based on provider treatment durations, not to exceed 6 months.

## 2 . Revision History

Date	Notes
5/13/2021	Arizona Medicaid 7.1 Implementation

OAB - Overactive Bladder Agents - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147213
<b>Guideline Name</b>	OAB - Overactive Bladder Agents - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name: Brand Enablex, generic darifenacin ER, Brand Ditropan XL, Flavoxate, Gelnique, Gemtesa, Brand Myrbetriq, generic mirabegron, generic oxybutynin 2.5mg IR tablet, generic oxybutynin oral solution, Oxytrol (Rx) patch, trospium IR/ER, Brand Vesicare, generic solifenacin	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The patient has a history of failure, contraindication, or intolerance to a trial of THREE preferred products  <ul style="list-style-type: none"><li>oxybutynin (generic Ditropan) 5 mg tablet</li></ul>	

- oxybutynin ER (generic Ditropan XL)
- Brand Detrol
- Brand Detrol LA
- Brand Toviaz

**AND**

**2** - For oxybutynin solution requests ONLY, patient must have intolerance to oxybutynin syrup

## **2 . Revision History**

Date	Notes
5/24/2024	Added generic mirabegron as NP target

Ocaliva

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99630
<b>Guideline Name</b>	Ocaliva
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Ocaliva	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of primary biliary cholangitis (aka primary biliary cirrhosis)  <b>AND</b>	

**2 - ONE of the following:**

**2.1 BOTH of the following:**

**2.1.1** Patient has failed to achieve an alkaline phosphatase (ALP) level of less than 1.67 times the upper limit of normal after at least 12 consecutive months of treatment with ursodeoxycholic acid(e.g., Urso, ursodiol)

**AND**

**2.1.2** Used in combination with ursodeoxycholic acid (e.g., Urso, ursodiol)

**OR**

**2.2** History of contraindication or intolerance to ursodeoxycholic acid (e.g., Urso, ursodiol)

**AND**

**3 - Prescribed by ONE of the following:**

- Hepatologist
- Gastroenterologist

Product Name:Ocaliva	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., laboratory values) documenting a reduction in alkaline phosphatase (ALP) level from pre-treatment baseline (i.e., prior to Ocaliva therapy) while on Ocaliva therapy	

**AND**

**2** - Prescribed by ONE of the following:

- Hepatologist
- Gastroenterologist

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Octreotide Products - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242218
<b>Guideline Name</b>	Octreotide Products - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Sandostatin, Generic octreotide, Sandostatin LAR, Bynfezia	
Diagnosis	Acromegaly
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of acromegaly	

**AND**

**2** - One of the following:

**2.1** Inadequate response to one of the following:

- Surgery
- Pituitary irradiation

**OR**

**2.2** Not a candidate for surgical resection or pituitary irradiation

**AND**

**3** - Trial and failure, contraindication, or intolerance to a dopamine agonist (e.g., bromocriptine or cabergoline) at maximally tolerated doses

**AND**

**4** - One of the following:

**4.1** Patient has had a trial of short-acting generic octreotide and responded to and tolerated therapy (Applies to Sandostatin LAR only)

**OR**

**4.2** Trial and failure, or intolerance to generic octreotide (Applies to Brand Sandostatin and Bynfezia only)

Product Name: Mycapssa	
Diagnosis	Acromegaly
Approval Length	12 month(s)
Therapy Stage	Initial Authorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of acromegaly</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p>2.1 Inadequate response to one of the following:</p> <ul style="list-style-type: none"> <li>• Surgery</li> <li>• Pituitary irradiation</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Not a candidate for surgical resection or pituitary irradiation</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient has responded to and tolerated treatment with generic octreotide or lanreotide</p>	

Product Name: Brand Sandostatin, Generic octreotide, Sandostatin LAR, Bynfezia, Mycapssa	
Diagnosis	Acromegaly
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., reduction or normalization of IGF-1/GH level for same age and sex, reduction in tumor size)</p>	

Product Name: Brand Sandostatin, Generic octreotide, Sandostatin LAR, Bynfezia	
Diagnosis	Carcinoid Tumors, for Symptomatic Treatment of Diarrhea or Flushing
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of metastatic carcinoid tumor requiring symptomatic treatment of severe diarrhea or flushing episodes</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p>2.1 Patient has had a trial of short-acting generic octreotide and responded to and tolerated therapy (Applies to Sandostatin LAR only)</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Trial and failure, or intolerance to generic octreotide (Applies to Brand Sandostatin and Bynfezia only)</p>	

Product Name: Brand Sandostatin, Generic octreotide, Sandostatin LAR, Bynfezia	
Diagnosis	Carcinoid Tumors, for Symptomatic Treatment of Diarrhea or Flushing
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of an improvement in the number of diarrhea or flushing episodes</p>	

Product Name: Brand Sandostatin, Generic octreotide, Sandostatin LAR, Bynfezia	
Diagnosis	Vasoactive Intestinal Peptide Tumors, for Symptomatic Treatment of Diarrhea
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of vasoactive intestinal peptide tumor requiring treatment of profuse watery diarrhea</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p>2.1 Patient has had a trial of short-acting generic octreotide and responded to and tolerated therapy (Applies to Sandostatin LAR only)</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Trial and failure, or intolerance to generic octreotide (Applies to Brand Sandostatin and Bynfezia only)</p>	

Product Name: Brand Sandostatin, Generic octreotide, Sandostatin LAR, Bynfezia	
Diagnosis	Vasoactive Intestinal Peptide Tumors, for Symptomatic Treatment of Diarrhea
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of an improvement in the number of diarrhea episodes</p>	

## 2 . Revision History

Date	Notes
4/24/2025	Added octreotide 10mg kit

Ohtuvayre (ensifentrine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-156840
<b>Guideline Name</b>	Ohtuvayre (ensifentrine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Ohtuvayre	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of chronic obstructive pulmonary disease  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) documenting BOTH of the following:

- Post-bronchodilator FEV1 % predicted greater than or equal to 30% and less than 80%
- Post-bronchodilator forced expiratory volume (FEV1) / forced vital capacity (FVC) ratio less than 0.7

**AND**

**3** - ONE of the following:

**3.1** BOTH of the following:

**3.1.1** Patient is on a stabilized dose and receiving concomitant therapy with ONE of the following (document name):

- A LABA/LAMA (e.g., Anoro Ellipta, Bevespi Aerosphere, Stiolto Respimat)
- A long-acting antimuscarinic agent [LAMA (e.g., Spiriva Respimat/HandiHaler)]
- A long-acting beta-agonist [LABA (e.g., Serevent Diskus)]
- An ICS/LABA/LAMA (i.e., Breztri Aerosphere, Trelegy Ellipta)

**AND**

**3.1.2** ONE of the following:

**3.1.2.1** ALL of the following:

- FEV1 less than 50% of predicted
- History of chronic bronchitis
- History of failure, contraindication, or intolerance to a selective phosphodiesterase 4 (PDE4) inhibitor [i.e., roflumilast (Daliresp)]

**OR**

**3.1.2.2** FEV1 is less than 80% of predicted but greater than or equal to 50% of predicted

**OR**

**3.2** BOTH of the following:

**3.2.1** Patient has a failure, contraindication, or intolerance to ALL of the following (document name and date tried):

- A long-acting beta-agonist [LABA (e.g., Serevent Diskus)]
- A long-acting antimuscarinic agent [LAMA (e.g., Spiriva Respimat/HandiHaler)]
- A LABA/LAMA (e.g., Anoro Ellipta, Bevespi Aerosphere, Stiolto Respimat),
- An ICS/LABA/LAMA (i.e., Breztri Aerosphere, Trelegy Ellipta)

**AND**

**3.2.2** ONE of the following:

**3.2.2.1** ALL of the following:

- FEV1 less than 50% of predicted
- History of chronic bronchitis
- History of failure, contraindication, or intolerance to a selective phosphodiesterase 4 (PDE4) inhibitor [i.e., roflumilast (Daliresp)]

**OR**

**3.2.2.2** FEV1 is less than 80% of predicted but greater than or equal to 50% of predicted

**OR**

**3.3** ALL of the following:

**3.3.1** Patient is unable to use a metered-dose, dry powder or slow mist inhaler (e.g. Spiriva Respimat) to control their COPD due to ONE of the following:

- Cognitive or physical impairment limiting coordination of handheld devices (e.g., cognitive decline, arthritis in the hands) (Document impairment)
- Patient is unable to generate adequate inspiratory force (e.g., peak inspiratory flow rate (PIFR) resistance is <60 L/min)

**AND**

**3.3.2** Patient requires the use of BOTH of the following (document date)

- A nebulized LABA [i.e., arformoterol (generic Brovana), formoterol (generic Perforomist)]
- A nebulized long-acting antimuscarinic agent [LAMA (i.e.,Yupelri)]

**AND**

**3.3.3 ONE** of the following:

**3.3.3.1 ALL** of the following:

- FEV1 less than 50% of predicted
- History of chronic bronchitis
- History of failure, contraindication, or intolerance to a selective phosphodiesterase 4 (PDE4) inhibitors [i.e., roflumilast (Daliresp)]

**OR**

**3.3.3.2** FEV1 is less than 80% of predicted but greater than or equal to 50% of predicted

**AND**

**4** - Patient experiences dyspnea during everyday activities (e.g., short of breath when walking up a slight hill)

**AND**

**5** - Prescribed by or in consultation with a Pulmonologist

Product Name:Ohtuvayre	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to Ohtuvayre therapy demonstrated by BOTH of the following:

- Improved COPD symptoms (e.g., dyspnea)
- Improved FEV1

## 2 . Revision History

Date	Notes
10/1/2024	Updated reauth guideline type to Prior Authorization

Ojjaara (momelotinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-136958
<b>Guideline Name</b>	Ojjaara (momelotinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Ojjaara	
Diagnosis	Myelofibrosis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting all of the following: 1.1 Diagnosis of one of the following:	

- Primary myelofibrosis
- Post-polycythemia vera myelofibrosis
- Post-essential thrombocythemia myelofibrosis

**AND**

**1.2** Disease is intermediate or high risk

**AND**

**1.3** Patient has anemia

Product Name:Ojjaara	
Diagnosis	Myelofibrosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., symptom improvement, spleen volume reduction)</p>	

## 2 . Revision History

Date	Notes
11/27/2023	New Program

Olumiant (baricitinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300291
<b>Guideline Name</b>	Olumiant (baricitinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Olumiant	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**1.2** History of failure to a 3 month trial of one non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Oencia (abatacept)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred tocilizumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name: Olumiant	
Diagnosis	Rheumatoid Arthritis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a rheumatologist	

Product Name: Olumiant	
Diagnosis	Coronavirus disease 2019 (COVID-19)
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of COVID-19</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is hospitalized (Olumiant is only FDA approved when used for COVID-19 patients in an inpatient setting)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient requires one of the following:</p> <ul style="list-style-type: none"> <li>• Supplemental oxygen</li> <li>• Non-invasive mechanical ventilation</li> <li>• Invasive mechanical ventilation</li> <li>• Extracorporeal membrane oxygenation (ECMO)</li> </ul>	
Notes	NOTE: Olumiant is only FDA approved when used for COVID-19 patients in an inpatient setting

Product Name: Olumiant	
Diagnosis	Alopecia Areata
Approval Length	N/A - Requests for non-approvable diagnoses should not be approved
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Requests for coverage for diagnosis of Alopecia Areata are not authorized and will not be approved

Notes	Approval Length: N/A - Requests for Alopecia Areata should not be approved. Deny as a benefit exclusion.
-------	--

## 2 . Revision History

Date	Notes
7/3/2025	Updated criteria for RA (preferred agents/embedded steps, removed CoT)

Omvoh (mirikizumab-mrkz)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300292
<b>Guideline Name</b>	Omvoh (mirikizumab-mrkz)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Omvoh IV	
Diagnosis	Crohn's disease (CD)
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's disease (CD)  <b>AND</b>	

**2 - One of the following:**

- Frequent diarrhea and abdominal pain
- At least 10% weight loss
- Complications such as obstruction, fever, abdominal mass
- Abnormal lab values (e.g., C-reactive protein [CRP])
- CD Activity Index (CAI) greater than 220

**AND**

**3 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:**

- 6-mercaptopurine
- Azathioprine
- Corticosteroids (e.g., prednisone)
- Methotrexate

**AND**

**4 - One of the following:**

**4.1 Paid claims or submission of medical records (e.g., chart notes) confirming history of failure, contraindication, or intolerance to ALL of the following:**

- A preferred adalimumab biosimilar
- infliximab
- A preferred ustekinumab biosimilar

**OR**

**4.2 Paid claims or submission of medical records (e.g., chart notes) confirming continuation of prior therapy, defined as no more than a 45-day gap in therapy**

**AND**

**5 - Will be administered as an intravenous induction dose**

**AND**

**6** - Prescribed by or in consultation with a gastroenterologist

Product Name: Omvoh SC

Diagnosis	Crohn's disease (CD)
-----------	----------------------

Approval Length	6 month(s)
-----------------	------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's disease (CD)

**AND**

**2** - Will be used as a maintenance dose following the intravenous induction doses

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name: Omvoh IV

Diagnosis	Ulcerative Colitis (UC)
-----------	-------------------------

Approval Length	3 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis

**AND**

**2** - One of the following:

- Greater than 6 stools per day
- Frequent blood in the stools
- Frequent urgency
- Presence of ulcers
- Abnormal lab values (e.g., hemoglobin, erythrocyte sedimentation rate, C-reactive protein)
- Dependent on, or refractory to, corticosteroids

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

- 6-mercaptopurine
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)
- Azathioprine
- Corticosteroids (e.g., prednisone)

**AND**

**4** - One of the following:

**4.1** Paid claims or submission of medical records (e.g., chart notes) confirming history of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**OR**

**4.2** Paid claims or submission of medical records (e.g., chart notes) confirming continuation of prior therapy, defined as no more than a 45-day gap in therapy

**AND**

**5** - Will be administered as an intravenous induction dose

**AND**

**6** - Prescribed by or in consultation with a gastroenterologist

Product Name: Omvoh SC

Diagnosis	Ulcerative Colitis (UC)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis

**AND**

**2** - Will be used as a maintenance dose following the intravenous induction doses

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name: Omvoh SC

Diagnosis	Crohn's Disease (CD), Ulcerative Colitis (UC)
-----------	---

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:</p> <ul style="list-style-type: none"> <li>• Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li> <li>• Reversal of high fecal output state</li> </ul>	

## 2 . Revision History

Date	Notes
7/3/2025	Updated steps for UC/CD (adding step through Yesintek)

Opfolda (miglustat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-139340
<b>Guideline Name</b>	Opfolda (miglustat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Opfolda	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting all of the following:  1.1 Diagnosis of late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency)	

**AND**

**1.2** Disease is confirmed by one of the following:

- Absence or deficiency (less than 40% of the lab specific normal mean) of GAA enzyme activity in lymphocytes, fibroblasts, or muscle tissues as confirmed by an enzymatic assay
- Molecular genetic testing confirms mutations in the GAA gene

**AND**

**1.3** Presence of clinical signs and symptoms of the disease (e.g., respiratory distress, skeletal muscle weakness, etc.)

**AND**

**1.4** Medication is used in combination with Pombiliti (cipaglucosidase alfa-atga)

**AND**

**1.5** Patient weight is greater than or equal to 40 kg

**AND**

**2** - Opfolda is not substituted with other miglustat products (i.e., Zavesca, Yargesa)

Product Name:Opfolda	
Approval Length	24 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Patient demonstrates positive clinical response to therapy (e.g., improvement in FVC, improvement in 6-minute walk distance [6MWD])

**AND**

2 - Medication is used in combination with Pombiliti (cipaglicosidase alfa-atga)

**AND**

3 - Opfolda is not substituted with other miglustat products (i.e., Zavesca, Yargesa)

## 2 . Revision History

Date	Notes
1/23/2024	New program

Opzelura (ruxolitinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-156490
<b>Guideline Name</b>	Opzelura (ruxolitinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Opzelura	
Diagnosis	Atopic Dermatitis
Approval Length	12 Week(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of mild to moderate atopic dermatitis	

**AND**

**2** - One of the following:

- Greater than or equal to 3% body surface area (BSA) involvement
- Involvement of sensitive body areas (e.g., face, hands, feet, scalp, groin)

**AND**

**3** - Patient is 12 years of age or older

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Allergist/Immunologist

**AND**

**5** - Trial and failure of a minimum 30-day supply of non-pharmacologic topical therapies (e.g., moisturizers)

**AND**

**6** - Trial and failure of a minimum 30-day supply (14-day supply for topical corticosteroids), contraindication, or intolerance to at least TWO of the following:

- Medium or higher potency topical corticosteroid
- Elidel (pimecrolimus) cream\*
- Tacrolimus ointment
- Eucrisa (crisaborole) ointment\*

**AND**

7 - Patient is not receiving Opzelura in combination with a potent immunosuppressant (e.g., azathioprine or cyclosporine)

**AND**

8 - Opzelura will only be used for short-term and/or non-continuous chronic treatment

Notes

\*Product may require step therapy

Product Name:Opzelura

Diagnosis Atopic Dermatitis

Approval Length 6 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

### Approval Criteria

1 - Documentation of a positive clinical response to therapy as evidenced by at least ONE of the following:

- Reduction in body surface area involvement from baseline
- Reduction in pruritus severity from baseline
- Improvement in quality of life from baseline

**AND**

2 - Patient is not receiving Opzelura in combination with a potent immunosuppressant (e.g., azathioprine or cyclosporine)

**AND**

3 - Opzelura will only be used for short-term and/or non-continuous chronic treatment

Product Name:Opzelura

Diagnosis	Nonsegmental Vitiligo
Approval Length	N/A - Requests for non-approvable diagnoses should not be approved
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Requests for coverage for diagnosis of nonsegmental vitiligo are not authorized and will not be approved	
Notes	Approval Length: N/A - Requests for nonsegmental vitiligo should not be approved. Deny as a benefit exclusion.

## 2 . Background

Clinical Practice Guidelines			
Table 1. Relative potencies of topical corticosteroids [2]			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment, gel	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream, lotion	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05

	Diflorasone diacetate	Cream	0.05
	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment, lotion	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream, lotion	0.1
	Triamcinolone acetonide	Cream, ointment, lotion	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

### 3 . Revision History

Date	Notes
9/30/2024	Opzelura now Preferred, no change to clinical criteria.

Oral Oncology Agents

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242214
<b>Guideline Name</b>	Oral Oncology Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Oral Oncology Drugs: Brand Gleevec, generic imatinib, Imkeldi, Brand Revlimid, generic lenalidomide	
Diagnosis	Cancer Indications
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The drug is being used as indicated by National Comprehensive Cancer Network (NCCN) guidelines with a Category of Evidence and Consensus of 1, 2A, or 2B	

**AND**

**2** - If the request is for the non-preferred Brand (Brand Gleevec, Imkeldi, or Brand Revlimid), patient must have tried and failed the preferred generic counterpart (generic imatinib or generic lenalidomide)

Product Name: Oral Oncology Drugs: Abirtega, Akeega, Alecensa, Alunbrig, Augtyro, Avyakit, Balversa, Bosulif capsules, Bosulif tablets, Braftovi, Brukinsa, Cabometyx, Calquence, Caprelsa, Cometriq, Copiktra, Cotellic, Danziten, Daurismo, Erivedge, Erleada, etoposide capsules, Farydak, Fruzaqla, Gavreto, Gilotrif, Gomekli, Hycamtin capsules, Ibrance, Iclusig, Idhifa, Imbruvica, Inlyta, Inrebic, Brand Iressa, generic gefitinib, Itovebi, Iwilfin, Jakafi, Jaypirca, Jylamvo, Kisqali, Kisqali-Femara Co-pack, Koselugo, Krazati, Lazcluze, Lenvima, Lonsurf, Lorbrena, Lumakras, Lynparza, Lytgobi, Mekinist, Mektovi, Nerlynx, Brand Nexavar, generic sorafenib, Ninlaro, Nubeqa, Odomzo, Ogsiveo, Ojemda, Orserdu, Pemazyre, Piqray, Pomalyst, Brand Purixan, generic mercaptopurine suspension, Qinlock, Retevmo, Revuforj, Rezlidhia, Romvimza, Rozlytrek, Rubraca, Rydapt, Scemblix, Brand Sprycel, generic dasatinib, Stivarga, Tabrecta, Tafinlar, Tagrisso, Talzena, Brand Tarceva, generic erlotinib, Tassigna, Tazverik, temozolomide capsules, Tepmetko, Tibsovo, Truqap, Tukysa, Turalio, Brand Tykerb, generic lapatinib, Vanflyta, Venclexta, Verzenio, Vitrakvi, Vizimpro, Voranigo, Brand Votrient, generic pazopanib, Welireg, Xalkori, Xatmep, Brand Xeloda, generic capecitabine, Xospata, Xpovio, Xtandi, Yonsa, Zejula, Zelboraf, Zolinza, Zydelig, Zykadia, Brand Zytiga, generic abiraterone

Diagnosis	Cancer Indications
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - The drug is being used as indicated by National Comprehensive Cancer Network (NCCN) guidelines with a Category of Evidence and Consensus of 1, 2A, or 2B

**2 . Revision History**

Date	Notes
4/24/2025	Added pazopanib (gen Votrient), revuforj, and Abirtega.

Orencia (abatacept)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-136981
<b>Guideline Name</b>	Orencia (abatacept)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Orencia IV or Orencia SC	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active rheumatoid arthritis	

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Trial and failure, contraindication, or intolerance to ONE nonbiologic disease-modifying antirheumatic drug (DMARD) (e.g., methotrexate [Rheumatrex/Trexall], Arava [leflunomide], Azulfidine [sulfasalazine])

Product Name: Oencia IV or Oencia SC	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

Product Name: Oencia IV or Oencia SC	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis	

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Trial and failure, contraindication, or intolerance to ONE of the following nonbiologic disease modifying anti-rheumatic drugs (DMARDs):

- leflunomide (Arava)
- methotrexate (Rheumatrex/Trexall)

Product Name:Orencia IV or Orencia SC	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

Product Name:Orencia IV or Orencia SC	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of active psoriatic arthritis (PsA)

**AND**

2 - Patient is 2 years of age or older

**AND**

3 - Prescribed by or in consultation with one of the following:

- Dermatologist
- Rheumatologist

Product Name: Orenzia IV or Orenzia SC

Diagnosis	Psoriatic Arthritis (PsA)
-----------	---------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

Product Name: Orenzia IV

Diagnosis	Prophylaxis for Acute Graft versus Host Disease (aGVHD)
-----------	---

Approval Length	2 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Used for prophylaxis of acute graft versus host disease (aGVHD)

**AND**

**2** - Patient is 2 years of age or older

**AND**

**3** - Patient will receive hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated donor

**AND**

**4** - Recommended antiviral prophylactic treatment for Epstein-Barr Virus (EBV) reactivation (e.g., acyclovir) will be administered prior to Orencia and continued for six months after HSCT

**AND**

**5** - Used in combination with both of the following:

- calcineurin inhibitor (e.g., cyclosporine, tacrolimus)
- methotrexate

## 2 . Revision History

Date	Notes
11/28/2023	Added age criterion to PsA indication

Orfadin (nitisinone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-128930
<b>Guideline Name</b>	Orfadin (nitisinone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Brand Orfadin, generic nitisinone	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of hereditary tyrosinemia type 1	

## 2 . Revision History

Date	Notes
7/25/2023	Added GPI for nitisinone. Updated guideline name.

Orilissa

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99485
<b>Guideline Name</b>	Orilissa
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Orilissa 150 mg	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderate to severe pain associated with endometriosis  <b>AND</b>	

2 - Patient is premenopausal

**AND**

3 - History of trial and failure (e.g., inadequate pain relief), contraindication or intolerance after a three month trial of TWO analgesics (e.g., ibuprofen, meloxicam, naproxen)

**AND**

4 - History of trial and failure, contraindication, or intolerance after a three month trial to ONE of the following:

- Hormonal contraceptives
- Progestins [e.g., norethindrone (generic Aygestin)]

**AND**

5 - Prescribed by or in consultation with ONE of the following:

- Obstetrics/Gynecologist (OB/GYN)
- Reproductive endocrinologist

Product Name:Orilissa 150 mg	
Approval Length	6 months*
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	

2 - Impact to bone mineral density has been considered

**AND**

3 - Treatment duration has not exceeded a total of 24 months\*\*

Notes

\*NOTE: Authorization for Orilissa 150 mg will be issued for 6 months up to a maximum of 24 months.

\*\*NOTE: Orilissa 150 mg once daily is indicated for a maximum of 24 months.

Product Name:Orilissa 200 mg

Approval Length

6 months\*

Guideline Type

Prior Authorization

**Approval Criteria**

1 - Diagnosis of moderate to severe pain associated with endometriosis

**AND**

2 - Patient is premenopausal

**AND**

3 - History of trial and failure (e.g., inadequate pain relief), contraindication or intolerance after a three month trial of TWO analgesics (e.g., ibuprofen, meloxicam, naproxen)

**AND**

4 - History of trial and failure, contraindication, or intolerance after a three month trial to ONE of the following:

- Hormonal contraceptives

- Progestins [e.g., norethindrone (generic Aygestin)]

**AND**

**5** - Prescribed by or in consultation with ONE of the following:

- Obstetrics/Gynecologist (OB/GYN)
- Reproductive endocrinologist

Notes

\*NOTE: Orilissa 200 mg twice daily is indicated for a maximum of 6 months.

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Orkambi - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116122
<b>Guideline Name</b>	Orkambi - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Orkambi	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystic fibrosis (CF)  <b>AND</b>	

**2** - Submission of laboratory results confirming that patient is homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene

**AND**

**3** - Patient is 1 year of age or older

**AND**

**4** - Prescribed by, or in consultation with, a specialist affiliated with a CF care center

Product Name:Orkambi	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Provider attests that the patient has achieved a clinically meaningful response while on Orkambi therapy to ONE of the following:</p> <ul style="list-style-type: none"><li>• Lung function as demonstrated by percent predicted expiratory volume in 1 second (ppFEV1)</li><li>• Body mass index (BMI)</li><li>• Pulmonary exacerbations</li><li>• Quality of life as demonstrated by Cystic Fibrosis Questionnaire-Revised (CFQ-R) respiratory domain score</li></ul> <p><b>AND</b></p> <p><b>2</b> - Prescribed by, or in consultation with, a specialist affiliated with a cystic fibrosis (CF) care center</p>	

## 2 . Revision History

Date	Notes
10/27/2022	Updated age requirement, added new GPI

Osphena - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99486
<b>Guideline Name</b>	Osphena - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Osphena	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Treatment of moderate to severe vaginal dryness, a symptom of vulvar and vaginal atrophy (VVA), due to menopause*	

**AND**

**2** - History of failure, contraindication, or intolerance to BOTH of the following:

- Estradiol vaginal cream
- Estradiol vaginal tablet

Notes

\*Treatment of dyspareunia is a benefit exclusion.

Product Name: Osphena

Approval Length 12 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Otezla (apremilast)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152645
<b>Guideline Name</b>	Otezla (apremilast)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Otezla	
Diagnosis	Psoriatic Arthritis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of active psoriatic arthritis	

**AND**

**2** - History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial)\*

**AND**

**3** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Notes

\*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial

Product Name: Otezla

Diagnosis Behcet's Disease

Approval Length 12 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Diagnosis of Behcet's Disease

**AND**

**2** - Patient has active oral ulcers

**AND**

**3** - History of failure, contraindication, or intolerance to one non-biologic (e.g., corticosteroids,

colchicine) within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial)\*

**AND**

**4** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Notes

\*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial

Product Name: Otezla

Diagnosis	Psoriatic Arthritis, Behcet's Disease
-----------	---------------------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Documentation of positive clinical response to Otezla therapy

**AND**

**2** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Product Name: Otezla

Diagnosis	Plaque Psoriasis
-----------	------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - Diagnosis of plaque psoriasis</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>2 - Both of the following:</b></p> <ul style="list-style-type: none"> <li>• Patient is 6 years of age or older</li> <li>• Patient weighs at least 20 kg</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3 - Both of the following:</b></p> <p><b>3.1</b> History of failure to one of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial):*</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., betamethasone, clobetasol, desonide)</li> <li>• Vitamin D analogs (e.g., calcitriol, calcipotriene)</li> <li>• Tazarotene</li> <li>• Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)</li> <li>• Anthralin</li> <li>• Coal tar</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>3.2</b> History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced (document drug, date, and duration of trial)*</p> <p style="text-align: center;"><b>AND</b></p> <p><b>4 - Prescribed by or in consultation with a dermatologist</b></p>	

Notes	*Note: Claims history may be used in conjunction as documentation of drug, date, and duration of trial
-------	--

Product Name:Otezla	
Diagnosis	Plaque Psoriasis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Otezla therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a dermatologist</p>	

## 2 . Revision History

Date	Notes
8/27/2024	Added new GPIs

Oxervate

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99634
<b>Guideline Name</b>	Oxervate
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Oxervate	
Diagnosis	Neurotrophic keratitis
Approval Length	8 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Stage 2 or 3 neurotrophic keratitis  <b>AND</b>	

**2** - History of failure to at least one OTC ocular artificial tear product (e.g., Systane® Ultra, Akwa® Tears, Refresh Optive®, Soothe® XP)

**AND**

**3** - Prescribed by or in consultation with ONE of the following:

- Ophthalmologist
- Optometrist

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Palforzia [Peanut (Arachis hypogaea)]

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-224202
<b>Guideline Name</b>	Palforzia [Peanut (Arachis hypogaea)]
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Palforzia	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis and clinical history of peanut allergy as documented by BOTH of the following:  1.1 A serum peanut-specific IgE level of greater than or equal to 0.35 kUA/L (kilo units of allergen per liter)	

**AND**

**1.2** A meal wheal diameter that is at least 3mm (millimeters) larger than the negative control on skin-prick testing for peanut

**AND**

**2** - ONE of the following:

**2.1** BOTH of the following:

- Patient is 1 to 17 years of age
- Patient is in the initial dose escalation phase of therapy

**OR**

**2.2** BOTH of the following:

- Patient is 1 year of age and older
- Patient is in the up-dosing or maintenance phase of therapy

**AND**

**3** - Used in conjunction with a peanut-avoidant diet

**AND**

**4** - Patient does not have one of the following:

- History of eosinophilic esophagitis (EoE) or eosinophilic gastrointestinal disease
- History of severe or life-threatening episode(s) of anaphylaxis or anaphylactic shock within the past 2 months
- Severe or poorly controlled asthma

**AND**

**5** - Prescribed by or in consultation with an allergist or immunologist

**AND**

**6** - Prescriber is certified/enrolled in the Palforzia REMS (Risk Evaluation and Mitigation Strategy) Program

Product Name:Palforzia

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to Palforzia therapy

**AND**

**2** - Used in conjunction with a peanut-avoidant diet

**AND**

**3** - Prescribed by or in consultation with an allergist or immunologist

**AND**

**4** - Prescriber is certified/enrolled in the Palforzia REMS (Risk Evaluation and Mitigation Strategy) Program

**2 . Revision History**

Date	Notes
3/26/2025	Added new GPIs.

Palynziq

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99636
<b>Guideline Name</b>	Palynziq
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Palynziq	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of phenylketonuria (PKU)  <b>AND</b>	

**2** - Patient is actively on a phenylalanine-restricted diet

**AND**

**3** - Physician attestation that the patient will not be receiving Palynziq in combination with Kuvan (sapropterin dihydrochloride)

**AND**

**4** - Submission of medical records (e.g. chart notes, laboratory values) documenting that the patient has a blood phenylalanine concentration greater than 600 micromoles per liter

Product Name:Palynziq	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Patient is actively on a phenylalanine-restricted diet	
<b>AND</b>	
<b>2</b> - ONE of the following:	
<b>2.1</b> Submission of medical records (e.g. chart notes, laboratory values) documenting that the patient has a blood phenylalanine concentration less than 600 micromoles per liter	
<b>OR</b>	
<b>2.2</b> Submission of medical records (e.g. chart notes, laboratory values) documenting that the patient has achieved a 20% reduction in blood phenylalanine concentration from pre-treatment baseline	

**OR**

**2.3** BOTH of the following:

**2.3.1** Patient is in initial titration/maintenance phase of dosing regimen (week 1-33)

**AND**

**2.3.2** Patient will receive maximum labeled dosage of 40 milligrams (mg) once daily if response has not been obtained after 24 weeks of 20 mg once daily maintenance dosing

**AND**

**3** - Submission of medical records (e.g. chart notes, laboratory values) documenting that the patient is not receiving Palynziq in combination with Kuvan (sapropterin dihydrochloride) [Prescription claim history that does not show any concomitant Kuvan claim within 60 days of reauthorization request may be used as documentation]

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Panretin

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99511
<b>Guideline Name</b>	Panretin
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name:Panretin	
Diagnosis	AIDS-related Kaposi's Sarcoma (KS)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of acquired immunodeficiency syndrome (AIDS)-related Kaposi's Sarcoma (KS)	

**AND**

**2** - Patient is not receiving systemic anti-KS treatment

Product Name:Panretin	
Diagnosis	NCCN Recommended Regimen
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Panretin will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.	

Product Name:Panretin	
Diagnosis	NCCN Recommended Regimen
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Documentation of positive clinical response to Panretin therapy	

## 2 . Revision History

Date	Notes
------	-------

4/8/2021	7/1 Implementation
----------	--------------------

Pediculicides - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-105258
<b>Guideline Name</b>	Pediculicides - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

### 1 . Criteria

Product Name:Sklice, Brand Natroba, generic spinosad susp	
Diagnosis	Head lice
Approval Length	30 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of topical treatment of head lice infestations  <b>AND</b>	

**2 - For Brand Natroba requests ONLY: Trial and failure to generic spinosad suspension (verified via paid pharmacy claims or submission of medical records/chart notes)**

## **2 . Revision History**

Date	Notes
3/28/2022	Added step through generic for Brand Natroba.

Pedmark (sodium thiosulfate injection, solution)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-120432
<b>Guideline Name</b>	Pedmark (sodium thiosulfate injection, solution)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Pedmark	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting diagnosis of solid tumors  <b>AND</b>  2 - Disease is BOTH of the following:	

- Localized
- Non-Metastatic

**AND**

**3** - Used for the prevention of ototoxicity due to cisplatin-based chemotherapy

**AND**

**4** - Patient is 1 month of age or older

**AND**

**5** - Prescribed by or in consultation with an oncologist

**AND**

**6** - History of failure, or intolerance to generic sodium thiosulfate

## 2 . Revision History

Date	Notes
1/24/2023	New program

Piasky (crovalimab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157795
<b>Guideline Name</b>	Piasky (crovalimab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Piasky	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of paroxysmal nocturnal hemoglobinuria (PNH)	

**AND**

**2** - Patient is 13 years of age or older

**AND**

**3** - Patient weighs at least 40 kg

**AND**

**4** - Submission of medical records (e.g., chart notes) or paid claims confirming trial and failure, contraindication, or intolerance to one of the following:

- Soliris (eculizumab)
- Ultomiris (ravulizumab)

**AND**

**5** - Prescribed by or in consultation with a hematologist/oncologist

Product Name:Piasky	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions)	
<b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) or paid claims confirming trial and failure, contraindication, or intolerance to one of the following:

- Soliris (eculizumab)
- Ultomiris (ravulizumab)

## **2 . Revision History**

Date	Notes
10/25/2024	New program

Pombiliti (cipaglucoasidase alfa-atga)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-139338
<b>Guideline Name</b>	Pombiliti (cipaglucoasidase alfa-atga)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Pombiliti	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting all of the following:  1.1 Diagnosis of late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency)	

**AND**

**1.2** Disease is confirmed by one of the following:

- Absence or deficiency (less than 40% of the lab specific normal mean) of GAA enzyme activity in lymphocytes, fibroblasts, or muscle tissues as confirmed by an enzymatic assay
- Molecular genetic testing confirms mutations in the GAA gene

**AND**

**1.3** Presence of clinical signs and symptoms of the disease (e.g., respiratory distress, skeletal muscle weakness, etc.)

**AND**

**1.4** Medication is used in combination with Opfolda (miglustat)

**AND**

**1.5** Patient weight is greater than or equal to 40 kg

**AND**

**2** - Not to be used in combination with other miglustat products (i.e., Zavesca, Yargesa)

Product Name:Pombiliti	
Approval Length	24 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Patient demonstrates positive clinical response to therapy (e.g., improvement in FVC, improvement in 6-minute walk distance [6MWD])

**AND**

2 - Medication is used in combination with Opfolda (miglustat)

**AND**

3 - Not to be used in combination with other miglustat products (i.e., Zavesca, Yargesa)

## 2 . Revision History

Date	Notes
1/23/2024	New program

Pradaxa Pellet Packs

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-133837
<b>Guideline Name</b>	Pradaxa Pellet Packs
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Pradaxa Pellet Packs	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Patient is 8 years of age or younger</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** ALL of the following:

**1.2.1** Patient is between 9 and 12 years of age

**AND**

**1.2.2** Requested medication is being used for one of the following diagnoses:

- Treatment of venous thromboembolic events (VTE) in patients who have been treated with a parenteral anticoagulant for at least 5 days
- To reduce the risk of recurrence of VTE in patients who have been previously treated

**AND**

**1.2.3** One of the following:

**1.2.3.1** Trial and failure, contraindication, or intolerance to Brand Pradaxa capsules (verified via paid pharmacy claims or submitted chart notes)

**OR**

**1.2.3.2** Patient is unable to swallow oral tablets/capsules

## **2 . Revision History**

Date	Notes
9/28/2023	New program

Praluent

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-129084
<b>Guideline Name</b>	Praluent
<b>Formulary</b>	<ul style="list-style-type: none"><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Praluent	
Diagnosis	Primary Hyperlipidemia [Including Heterozygous Familial Hypercholesterolemia (HeFH), Atherosclerotic Cardiovascular Disease (ASCVD), and Secondary Prevention of Cardiovascular Events in Patients with ASCVD]
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1 - Diagnosis of ONE of the following:**

**1.1 Heterozygous familial hypercholesterolemia (HeFH) as confirmed by ONE of the following\*:**

**1.1.1 BOTH of the following:**

**1.1.1.1 Pre-treatment low density lipoprotein cholesterol (LDL-C) of ONE of the following:**

- Greater than 190 milligrams per deciliter (mg/dL)
- Greater than 155 mg/dL if less than 16 years of age

**AND**

**1.1.1.2 ONE of the following:**

- Family history of myocardial infarction in first-degree relative less than 60 years of age
- Family history of myocardial infarction in second-degree relative less than 50 years of age
- Family history of LDL-C greater than 190 mg/dL in first- or second-degree relative
- Family history of heterozygous or homozygous familial hypercholesterolemia in first- or second-degree relative
- Family history of tendinous xanthomata and/or arcus cornealis in first- or second degree relative

**OR**

**1.1.2 BOTH of the following:**

**1.1.2.1 Pre-treatment LDL-C of ONE of the following:**

- Greater than 190 mg/dL
- Greater than 155 mg/dL if less than 16 years of age

**AND**

**1.1.2.2 Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following:**

- Functional mutation in LDL (low density lipoprotein), apoB (apolipoprotein B), or PCSK9 (proprotein convertase subtilisin/kexin type 9) gene\*
- Tendinous xanthomata

- Arcus cornealis before age 45

**OR**

**1.2** Atherosclerotic cardiovascular disease (ASCVD) as confirmed by ONE of the following:

- Acute coronary syndromes
- History of myocardial infarction
- Stable or unstable angina
- Coronary or other arterial revascularization
- Stroke
- Transient ischemic attack
- Peripheral arterial disease presumed to be of atherosclerotic origin

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following [prescription claims history may be used in conjunction as documentation of medication use, dose, and duration]:

**2.1** Patient has been receiving at least 12 consecutive weeks of high-intensity statin therapy [i.e. atorvastatin 40-80 milligrams (mg), rosuvastatin 20-40mg] and will continue to receive high intensity statin at maximally tolerated dose

**OR**

**2.2** BOTH of the following:

**2.2.1** Patient is unable to tolerate high-intensity statin as evidenced by one of the following intolerable and persistent (i.e. more than 2 weeks) symptoms:

- Myalgia (muscle symptoms without creatine kinase [CK] elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

**AND**

**2.2.2** ONE of the following:

**2.2.2.1** Patient has been receiving at least 12 consecutive weeks of moderate-intensity statin [i.e. atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin greater than or equal to

20 mg, pravastatin greater than or equal to 40 mg, lovastatin 40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 40 mg twice daily or Livalo (pitavastatin) greater than or equal to 2 mg] and will continue to receive a moderate-intensity statin at maximally tolerated dose

**OR**

**2.2.2.2** Patient has been receiving at least 12 consecutive weeks of low-intensity statin [i.e. simvastatin 10 mg, pravastatin 10-20 mg, lovastatin 20 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1 mg] therapy and will continue to receive a low-intensity statin at maximally tolerated dose

**OR**

**2.3** Patient is unable to tolerate low or moderate-, and high-intensity statins as evidenced by ONE of the following:

**2.3.1** ONE of the following intolerable and persistent (i.e. more than 2 weeks) symptoms for low or moderate-, and high-intensity statins:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

**OR**

**2.3.2** Patient has a labeled contraindication to all statins as documented in medical records

**OR**

**2.3.3** Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations greater than 10 times ULN

**AND**

**3** - ONE of the following:

**3.1** Submission of medical records (e.g., laboratory values) documenting ONE of the following LDL-C values while on maximally tolerated lipid lowering therapy for a minimum of at least 12 weeks within the last 120 days:

- LDL-C greater than or equal to 100 mg/dL with ASCVD
- LDL-C greater than or equal to 130 mg/dL without ASCVD

**OR**

**3.2 BOTH of the following:**

**3.2.1** Submission of medical records (e.g., laboratory values) documenting ONE of the following LDL-C values while on maximally tolerated lipid lowering therapy for a minimum of at least 12 weeks within the last 120 days:

- LDL-C between 55 mg/dL and 99 mg/dL with ASCVD
- LDL-C between 100 mg/dL and 129 mg/dL without ASCVD

**AND**

**3.2.2** Submission of medical records (e.g., laboratory values) documenting ONE of the following [prescription claims history may be used in conjunction as documentation of medication use, dose, and duration]:

**3.2.2.1** Patient has been receiving at least 12 consecutive weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy

**OR**

**3.2.2.2** Patient has a history of contraindication or intolerance to ezetimibe

**AND**

**4** - Used as an adjunct to a low-fat diet and exercise

**AND**

**5** - Prescribed by ONE of the following:

- Cardiologist
- Endocrinologist

<ul style="list-style-type: none"> <li>Lipid specialist</li> </ul>	
<p><b>AND</b></p>	
<p><b>6</b> - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Repatha (evolocumab))</p>	
Notes	*Note: Results of prior genetic testing can be submitted as confirmation of diagnosis of HeFH.

Product Name: Praluent	
Diagnosis	Primary Hyperlipidemia [Including Heterozygous Familial Hypercholesterolemia (HeFH), Atherosclerotic Cardiovascular Disease (ASCVD), and Secondary Prevention of Cardiovascular Events in Patients with ASCVD]
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient continues to receive statin at maximally tolerated dose (unless patient has documented inability to take statins)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is continuing a low-fat diet and exercise regimen</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by ONE of the following:</p> <ul style="list-style-type: none"> <li>Cardiologist</li> <li>Endocrinologist</li> <li>Lipid specialist</li> </ul>	

**AND**

**4** - Submission of medical records (e.g. chart notes, laboratory values) documenting low density lipoprotein cholesterol (LDL-C) reduction while on Praluent therapy

**AND**

**5** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Repatha (evolcumab))

Product Name:Praluent	
Diagnosis	Homozygous Familial Hypercholesterolemia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of homozygous familial hypercholesterolemia (HoFH) as confirmed by submission of medical records (e.g., chart notes, laboratory values) documenting BOTH of the following:*</p> <p><b>1.1</b> ONE of the following:</p> <ul style="list-style-type: none"><li>• Pre-treatment LDL-C (low-density lipoprotein cholesterol) greater than 500 mg/dL (milligrams per deciliter)</li><li>• Treated LDL-C greater than 300 mg/dL</li></ul> <p><b>AND</b></p> <p><b>1.2</b> ONE of the following:</p> <ul style="list-style-type: none"><li>• Xanthoma before 10 years of age</li><li>• Evidence of heterozygous familial hypercholesterolemia (HeFH) in both parents</li></ul>	

**AND**

**2** - Used as an adjunct to a low-fat diet and exercise

**AND**

**3** - Patient is receiving other lipid-lowering therapy (e.g., statin, ezetimibe, LDL [low-density lipoprotein] apheresis)

**AND**

**4** - Prescribed by ONE of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

**AND**

**5** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Repatha (evolcumab))

Notes	*Results of prior genetic testing can be submitted as confirmation of diagnosis of HoFH.
-------	--

Product Name:Praluent	
Diagnosis	Homozygous Familial Hypercholesterolemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Patient continues to receive other lipid-lowering therapy (e.g., statin, LDL apheresis)	

**AND**

**2** - Patient is continuing a low-fat diet and exercise regimen

**AND**

**3** - Prescribed by ONE of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

**AND**

**4** - Submission of medical records (e.g. chart notes, laboratory values) documenting low density lipoprotein cholesterol (LDL-C) reduction while on Praluent therapy

**AND**

**5** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Repatha (evolocumab))

## 2 . Revision History

Date	Notes
9/1/2023	Update to account for 2022 ACC recommendations of a lower LDL threshold of 55mg/dl for patients with ASCVD at very high risk.

Preferred Drugs- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99538
<b>Guideline Name</b>	Preferred Drugs- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Diagnosis	Prior Authorization Administrative Guideline for Preferred Drugs Without Drug-Specific Criteria
Approval Length	12 month(s)
Guideline Type	Administrative
<b>Approval Criteria</b> 1 - ALL of the following:  1.1 ONE of the following:	

**1.1.1** The requested drug must be used for a Food and Drug Administration (FDA)-approved indication

**OR**

**1.1.2** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology

**AND**

**1.2** The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plans' program

**AND**

**1.3** If the patient is less than FDA minimum age, the prescriber attests they are aware of FDA labeling and feels the treatment with the requested product is medically necessary.  
(Document rationale for use)

Notes	Medications used solely for anti-obesity/weight loss, cosmetic (e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs (Cialis/Tadalafil) are covered for clinical diagnoses other than ED.
-------	---

## 2 . Revision History

Date	Notes
6/2/2021	Arizona Medicaid 7.1 Implementation

Presbyopia Agents

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164139
<b>Guideline Name</b>	Presbyopia Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Qlosi, Vuity	
Approval Length	1 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of presbyopia  <b>AND</b>	

**2** - History of failure or intolerance to a generic pilocarpine ophthalmic preparation

**AND**

**3** - Prescribed by or in consultation with **ONE** of the following:

- Ophthalmologist
- Optometrist

**AND**

**4** - Provider confirms valid clinical rationale, which excludes lifestyle choice, as to why patient is unable to use corrective lenses (e.g., eyeglasses or contact lenses)

Product Name:Qlosi, Vuity	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Patient demonstrates positive clinical response to therapy ( e.g., improvement in near vision in low light conditions without lost of distance vision)	
<b>AND</b>	
<b>2</b> - Prescribed by or in consultation with <b>ONE</b> of the following:	
<ul style="list-style-type: none"><li>• Ophthalmologist</li><li>• Optometrist</li></ul>	

## **2 . References**

1. Vuity Prescribing Information. Abbvie Inc. North Chicago, IL. March 2023.
2. Qlosi Prescribing Information. Orasis Pharmaceuticals, Inc. Ponte Vedra, FL. October 2023.
3. ClinicalTrials.gov. Available at: Study Details | An Evaluation of the Efficacy and Safety of CSF-1 in the Temporary Correction of Presbyopia (NEAR-1) | ClinicalTrials.gov. Accessed January 12, 2025.
4. ClinicalTrials.gov. Available at: Study Details | An Evaluation of the Efficacy and Safety of CSF-1 in the Temporary Correction of Presbyopia (NEAR-2) | ClinicalTrials.gov. Accessed January 12, 2025.
5. Mckenzie, K.. A Look at a New Presbyopia Treatment - American Academy of Ophthalmology. Accessed January 12, 2025.
6. Orasis Pharmaceuticals Announces FDA Approval of QLOSI™ (pilocarpine hydrochloride ophthalmic solution) 0.4% for the Treatment of Presbyopia. October 18, 2023. Accessed January 12, 2025
7. Another Attempt at Presbyopia. December 15, 2023. Available at: Another Attempt at Presbyopia | PatientPoint. Accessed January 12, 2025.

### 3 . Revision History

Date	Notes
1/31/2025	New program

Pretomanid

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99488
<b>Guideline Name</b>	Pretomanid
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Pretomanid	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Diagnosis of pulmonary extensively drug resistant (XDR) tuberculosis (TB)  <p style="text-align: center;"><b>OR</b></p>	

**1.2** Treatment-intolerant or nonresponsive multidrug-resistant (MDR) tuberculosis (TB)

**AND**

**2** - Pretomanid will be used in combination with bedaquiline and linezolid

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Prevymis (letermovir)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164378
<b>Guideline Name</b>	Prevymis (letermovir)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Prevymis	
Diagnosis	CMV Prophylaxis in Hematopoietic Stem Cell Transplant (HSCT) Recipients
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - BOTH of the following: <ul style="list-style-type: none"><li>Patient is 6 months of age or older</li></ul>	

- Patient weighs at least 6 kg

**AND**

**2 - BOTH** of the following:

- Patient is a recipient of an allogeneic hematopoietic stem cell transplant
- Patient is cytomegalovirus (CMV) seropositive (R+)

**AND**

**3 - Provider attests that Prevymsis will be initiated between Day 0 and Day 28 post-transplantation (before or after engraftment) and is being prescribed as prophylaxis and not treatment of CMV infection**

Product Name:Prevymsis	
Diagnosis	CMV Prophylaxis in Kidney Transplant Recipients
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - BOTH</b> of the following:</p> <ul style="list-style-type: none"> <li>• Patient is 12 years of age or older</li> <li>• Patient weighs at least 40 kg</li> </ul> <p><b>AND</b></p> <p><b>2 - BOTH</b> of the following:</p> <ul style="list-style-type: none"> <li>• Patient is a recipient of a kidney transplant</li> <li>• Patient is cytomegalovirus (CMV) seronegative (Donor CMV seropositive/Recipient CMV seronegative [D+/R-])</li> </ul>	

**AND**

**3** - Provider attests that Prevymsis will be initiated between Day 0 and Day 7 post-transplantation; and is being prescribed as prophylaxis and not treatment of CMV infection

## **2 . Revision History**

Date	Notes
1/30/2025	Added age/weight criteria due to expanded approval for both indications.

Primary Hyperoxaluria (PH1) Agents [Oxlumo (lumasiran), Rivfloza (nedosiran)]

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242219
<b>Guideline Name</b>	Primary Hyperoxaluria (PH1) Agents [Oxlumo (lumasiran), Rivfloza (nedosiran)]
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Oxlumo	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of primary hyperoxaluria type 1 (PH1)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting diagnosis has been confirmed by both of the following:

**2.1** One of the following:

- Elevated urinary oxalate excretion
- Elevated plasma oxalate concentration
- Spot urinary oxalate to creatinine molar ratio greater than normal for age

**AND**

**2.2** One of the following:

- Genetic testing demonstrating a mutation in the alanine:glyoxylate aminotransferase (AGXT) gene
- Liver biopsy demonstrating absence or reduced alanine:glyoxylate aminotransferase (AGT) activity

**AND**

**3** - Patient has not received a liver transplant

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Hepatologist
- Nephrologist
- Urologist
- Geneticist
- Specialist with expertise in the treatment of PH1

Product Name: Rivfloza	
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of primary hyperoxaluria type 1 (PH1)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records (e.g., chart notes) documenting diagnosis has been confirmed by both of the following:</p> <p>2.1 One of the following:</p> <ul style="list-style-type: none"> <li>• Elevated urinary oxalate excretion</li> <li>• Elevated plasma oxalate concentration</li> <li>• Spot urinary oxalate to creatinine molar ratio greater than normal for age</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2.2 One of the following:</p> <ul style="list-style-type: none"> <li>• Genetic testing demonstrating a mutation in the alanine:glyoxylate aminotransferase (AGXT) gene</li> <li>• Liver biopsy demonstrating absence or reduced alanine:glyoxylate aminotransferase (AGT) activity</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is 2 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Patient has preserved kidney function ( e.g., eGFR greater than or equal to 30mL/min/1.73m<sup>2</sup>)</p>	

**AND**

**5** - Patient has not received a liver transplant

**AND**

**6** - Prescribed by or in consultation with one of the following:

- Hepatologist
- Nephrologist
- Urologist
- Geneticist
- Specialist with expertise in the treatment of PH1

Product Name: Oxlumo, Rivfloza

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy (e.g., decreased urinary oxalate excretion, decreased plasma oxalate concentration)

**AND**

**2** - Patient has not received a liver transplant

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Hepatologist

- Nephrologist
- Urologist
- Geneticist
- Specialist with expertise in the treatment of PH1

## 2 . Revision History

Date	Notes
4/24/2025	Updated age criterion for Rivfloza due to expanded indication

Procysbi

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99725
<b>Guideline Name</b>	Procysbi
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Procysbi	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of nephropathic cystinosis  <b>AND</b>	

2 - Patient is 1 year of age or older

**AND**

3 - History of failure or intolerance to Cystagon (immediate-release cysteamine bitartrate)\*

Notes

\*Note: AZM generally does not consider frequency of dosing and/or lack of compliance to dosing regimens, an indication of medical necessity

Product Name: Procysbi

Approval Length 12 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to Procysbi therapy

**2 . Revision History**

Date	Notes
5/14/2021	Arizona Medicaid 7.1 Implementation

Progesterone - Non-Oral

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99489
<b>Guideline Name</b>	Progesterone - Non-Oral
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Crinone, Endometrin	
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Treatment is for non-infertility use (e.g., secondary amenorrhea, reduce the risk of recurrent spontaneous preterm birth)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Proton Pump Inhibitors (PPIs)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242221
<b>Guideline Name</b>	Proton Pump Inhibitors (PPIs)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:generic lansoprazole ODT, Brand Nexium granules/suspension packets, generic esomeprazole granules/suspension packets, Brand Protonix Pak, generic pantoprazole suspension packet	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following: 1.1 Patient is less than 18 years of age	

**OR**

**1.2** All of the following:

**1.2.1** Patient is 18 years of age or older

**AND**

**1.2.2** Patient is unable to swallow solid oral dosage forms (e.g., tablets) in the past 30 days

**AND**

**1.2.3** History of failure to lansoprazole ODT (DOES NOT APPLY TO REQUESTS FOR LANSOPRAZOLE ODT)

**AND**

**2** - For BRAND NEXIUM granules/suspension packets requests ONLY: history of failure or intolerance to generic esomeprazole granules/suspension packets

## **2 . Revision History**

Date	Notes
4/24/2025	Added step through generic for Brand Nexium requests

Provigil, Nuvigil

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99490
<b>Guideline Name</b>	Provigil, Nuvigil
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Provigil, generic modafinil, Brand Nuvigil, generic armodafinil	
Diagnosis	Narcolepsy, Obstructive Sleep Apnea, Shift Work Disorder, Idiopathic Hypersomnia (off label)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following diagnoses: <ul style="list-style-type: none"><li>Narcolepsy</li><li>Excessive sleepiness due to obstructive sleep apnea</li></ul>	

- Excessive sleepiness due to shift work disorder (circadian rhythm sleep disorder, shift work type)
- Idiopathic hypersomnia

**AND**

**2** - If the request is for modafinil, the patient has a history of failure, contraindication, or intolerance to armodafinil

Product Name: Brand Provigil, generic modafinil, Brand Nuvigil, generic armodafinil	
Diagnosis	Fatigue due to Multiple Sclerosis (off-label)
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of multiple sclerosis (MS)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is experiencing fatigue</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - If the request is for modafinil, the patient has a history of failure, contraindication, or intolerance to armodafinil</p>	

Product Name: Brand Provigil, generic modafinil, Brand Nuvigil, generic armodafinil	
Diagnosis	Adjunctive Therapy for the Treatment of Major Depressive Disorder or Bipolar Depression (off-label)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Treatment-resistant depression, defined as BOTH of the following:

**1.1** Diagnosis of ONE of the following:

- Major depressive disorder (MDD)
- Bipolar depression

**AND**

**1.2** History of failure, contraindication, or intolerance to at least TWO antidepressants from different classes (e.g., SSRIs [selective serotonin reuptake inhibitors], SNRIs [serotonin-norepinephine reuptake inhibitors], bupropion)

**AND**

**2** - Used as adjunctive therapy

**AND**

**3** - If the request is for modafinil, the patient has a history of failure, contraindication, or intolerance to armodafinil

Product Name: Brand Provigil, generic modafinil, Brand Nuvigil, generic armodafinil	
Diagnosis	Adjunctive Therapy for the Treatment of Major Depressive Disorder or Bipolar Depression (off-label)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Documentation of positive clinical response to therapy

**AND**

2 - Used as adjunctive therapy

**AND**

3 - If the request is for modafinil, the patient has a history of failure, contraindication, or intolerance to armodafinil

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Pulmonary Arterial Hypertension (PAH) Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-208208
<b>Guideline Name</b>	Pulmonary Arterial Hypertension (PAH) Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	4/1/2025
-----------------	----------

### 1 . Criteria

Product Name: PREFERRED DRUGS: Alyq, generic tadalafil, generic ambrisentan, Orenitram, generic sildenafil suspension, generic sildenafil tablets, Tracleer tablet for oral suspension; NON-PREFERRED DRUGS: Brand Adcirca, Adempas, Brand Flolan, Brand Veletri, generic eprosteno, Brand Letairis, Opsumit, Opsynvi, Brand Remodulin, generic trepostinil, Brand Revatio tablets, Brand Revatio injection, generic sildenafil injection, Tadliq suspension, Brand Tracleer tablet, generic bosentan, Tyvaso DPI, Tyvaso inhalation solution, Upravi, Ventavis	
Diagnosis	Pulmonary Arterial Hypertension
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Diagnosis of pulmonary arterial hypertension

**AND**

**2** - Pulmonary arterial hypertension is symptomatic

**AND**

**3** - One of the following:

**3.1** Diagnosis of pulmonary arterial hypertension was confirmed by right heart catheterization

**OR**

**3.2** Patient is currently on any therapy for the diagnosis of pulmonary arterial hypertension

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Cardiologist

**AND**

**5** - If the patient is requesting a non preferred product, patient has a history of failure, contraindication or intolerance to BOTH of the following:

- A preferred Endothelin Receptor Antagonist (ERA) (e.g., generic ambrisentan, Tracleer tablet for oral suspension))
- A preferred Phosphodiesterase 5 inhibitor (PDE5i) [e.g., Alyq or tadalafil, generic sildenafil tablet (generic for Revatio tablet), generic sildenafil suspension]

**AND**

**6** - If the request is for Brand Adcirca, patient must have tried and failed generic tadalafil or Alyq

**AND**

**7** - If the request is for Opsynvi, patient must have tried and failed both of the following as separate products:

- generic tadalafil
- Opsumit (may require PA)

Product Name:Adempas tablet	
Diagnosis	Chronic Thromboembolic Pulmonary Hypertension (CTEPH)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - One of the following:</p> <p><b>1.1</b> Both of the following:</p> <p><b>1.1.1</b> Diagnosis of inoperable or persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH)</p> <p><b>AND</b></p> <p><b>1.1.2</b> CTEPH is symptomatic</p> <p><b>OR</b></p> <p><b>1.2</b> Patient is currently on any therapy for the diagnosis of CTEPH</p>	

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Cardiologist

Product Name: PREFERRED DRUGS: Alyq, generic tadalafil, generic ambrisentan, Orenitram, generic sildenafil suspension, generic sildenafil tablets, Tracleer tablet for oral suspension; NON-PREFERRED DRUGS: Brand Adcirca, Adempas, Brand Flolan, Brand Veletri, generic eprosteno, Brand Letairis, Opsumit, Opsynvi, Brand Remodulin, generic trepostinil, Brand Revatio tablets, Brand Revatio injection, generic sildenafil injection, Tadliq suspension, Brand Tracleer tablet, generic bosentan, Tyvaso DPI, Tyvaso inhalation solution, Upravi, Ventavis

Diagnosis	All indications listed above
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response to therapy

Product Name: Winrevair Injection

Diagnosis	Pulmonary Arterial Hypertension
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of pulmonary arterial hypertension

**AND**

**2** - Pulmonary arterial hypertension is symptomatic

**AND**

**3** - Patient is currently on at least two therapies indicated for the treatment of pulmonary arterial hypertension from the following different mechanisms of action, unless there is a contraindication or intolerance:

- Endothelin receptor antagonists (i.e., Bosentan, ambrisentan or macitentan)
- Phosphodiesterase 5 inhibitors (i.e., Tadalafil or sildenafil)

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Cardiologist

Product Name:Winrevair Injection	
Diagnosis	Pulmonary Arterial Hypertension
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	

## 2 . Revision History

Date	Notes
3/26/2025	P&T changes: Tracleer (tab for oral susp) to preferred, generic bosentan to NP.

Pulmozyme

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99638
<b>Guideline Name</b>	Pulmozyme
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Pulmozyme	
Diagnosis	Cystic Fibrosis
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of Cystic Fibrosis	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Pyrukynd (mitapivat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-107467
<b>Guideline Name</b>	Pyrukynd (mitapivat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Pyrukynd	
Diagnosis	Hemolytic Anemia
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ALL of the following:  1.1 Diagnosis of hemolytic anemia confirmed by the presence of chronic hemolysis (e.g.,	

increased indirect bilirubin, elevated lactated dehydrogenase [LDH], decreased haptoglobin, increased reticulocyte count)

**AND**

**1.2** Diagnosis of pyruvate kinase deficiency confirmed by molecular testing of ALL the following mutations on the PKLR gene:

- Presence of at least 2 variant alleles in the pyruvate kinase liver and red blood cell (PKLR) gene, of which at least 1 was a missense variant
- Patient is not homozygous for the c.1436G>A (p.R479H) variant
- Patient does not have 2 non-missense variants (without the presence of another missense variant) in the PKLR gene

**AND**

**1.3** Hemoglobin is less than or equal to 10g/dL

**AND**

**1.4** Patient has symptomatic anemia or is transfusion dependent

**AND**

**1.5** Exclusion of other causes of hemolytic anemias (e. g., infections, toxins, drugs)

**AND**

**2** - Prescribed by or in consultation with a hematologist

Product Name:Pyrukynd	
Diagnosis	Hemolytic Anemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy [e.g., hemoglobin greater than or equal to 1.5g/dL from baseline, reduction in transfusions of greater than or equal to 33% in the number of red blood cell units transfused during the fixed dose period compared with the patient's historical transfusion burden, improvement in markers of hemolysis from baseline (e.g., bilirubin, lactated dehydrogenase [LDH], haptoglobin, reticulocyte count)]

**AND**

2 - Prescribed by or in consultation with a hematologist

Notes	If the member does not meet the medical necessity reauthorization criteria requirements, a denial should be issued and a 1-month authorization should be issued one time for Pyrukynd gradual therapy discontinuation.
-------	--

## 2 . Revision History

Date	Notes
5/24/2022	New Program

Qalsody (tofersen)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-128982
<b>Guideline Name</b>	Qalsody (tofersen)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Qalsody	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting ALL of the following: 1.1 Diagnosis amyotrophic lateral sclerosis (ALS)	

**AND**

**1.2** Molecular genetic testing confirms mutation in the SOD1 gene

**AND**

**1.3** Patient's baseline functional ability has been documented prior to initiating treatment (e.g., speech, walking, climbing stairs, etc.)

**AND**

**1.4** Patient has a percent (%) slow vital capacity (%SVC) greater than or equal to 50% at the start of treatment [A]

**AND**

**1.5** Patient does not require permanent noninvasive ventilation or invasive ventilation

**AND**

**2** - Prescribed by or in consultation with a neurologist with expertise in the diagnosis of ALS

Product Name:Qalsody	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting slowed disease progression from baseline	

**AND**

**2** - Prescribed by or in consultation with a neurologist with expertise in the diagnosis of ALS

## **2 . Endnotes**

- A. Those in the faster-progressing subgroup, which the primary and key secondary endpoints were formally tested, were required to have a slow vital capacity (SVC) greater than or equal to 65% of predicted value for sex, age, and height (from the sitting position) at screening. [2]

## **3 . Revision History**

Date	Notes
7/26/2023	New program

Qutenza (capsaicin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-129071
<b>Guideline Name</b>	Qutenza (capsaicin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Qutenza	
Diagnosis	Neuropathic pain associated with postherpetic neuralgia (PHN)
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of neuropathic pain associated with postherpetic neuralgia (PHN)	

**AND**

**2** - Submission of medical records (e.g., chart notes, paid claims history) documenting trial and failure, contraindication, or intolerance to ALL of the following:

- gabapentin
- pregabalin
- minimum 60-day trial of a tricyclic antidepressant (e.g., amitriptyline, nortriptyline, desipramine)
- generic lidocaine 5% patch
- topical capsaicin cream

Product Name: Qutenza	
Diagnosis	Neuropathic pain associated with diabetic peripheral neuropathy (DPN) of the feet
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming diagnosis of neuropathic pain associated with diabetic peripheral neuropathy (DPN) of the feet</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Submission of medical records (e.g., chart notes, paid claims history) documenting trial and failure, contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"><li>• gabapentin</li><li>• pregabalin</li><li>• minimum 60-day trial of a tricyclic antidepressant (e.g., amitriptyline, nortriptyline, desipramine)</li><li>• generic lidocaine 5% patch</li><li>• topical capsaicin cream</li><li>• duloxetine</li></ul>	

Product Name:Qutenza	
Diagnosis	All indications
Approval Length	3 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - It has been at least 3 months since the last application/administration [B]</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient experienced pain relief with a prior course of therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is experiencing a return of neuropathic pain</p>	

## 2 . Revision History

Date	Notes
7/28/2023	New program

Radicava (edaravone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-163769
<b>Guideline Name</b>	Radicava (edaravone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Radicava IV, Radicava ORS, generic edaravone	
Diagnosis	Amyotrophic Lateral Sclerosis (ALS)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of documentation (e.g., chart notes) confirming diagnosis of “definite” or “probable” amyotrophic lateral sclerosis (ALS) per the revised EL Escorial and Airlie House diagnostic criteria	

**AND**

**2** - Prescribed by or in consultation with a neurologist with expertise in the diagnosis of ALS

**AND**

**3** - Patient has scores greater than or equal to 2 in all items of the ALS Functional Rating Scale-Revised (ALSFRS-R) criteria at the start of treatment

**AND**

**4** - Patient has a percent (%) forced vital capacity (%FVC) greater than or equal to 80% at the start of treatment

Product Name:Radicava IV, Radicava ORS, generic edaravone	
Diagnosis	Amyotrophic Lateral Sclerosis (ALS)
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of documentation (e.g., chart notes) confirming positive clinical response to therapy (e.g., slowing in the decline of functional abilities)	
<b>AND</b>	
<b>2</b> - Patient is not dependent on invasive ventilation or tracheostomy	

## 2 . Revision History

Date	Notes
1/31/2025	Added generic edaravone as target

Ranolazine products

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-110773
<b>Guideline Name</b>	Ranolazine products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/15/2022
-----------------	-----------

## 1 . Criteria

Product Name:Brand Ranexa, generic ranolazine	
Approval Length	12 month(s)
Guideline Type	Step Therapy
<b>Approval Criteria</b>  1 - History of ONE of the following standard anti-angina treatments:  1.1 One beta-blocker [e.g. Lopressor (metoprolol), Inderal (propranolol)]  <b>OR</b>	

**1.2** One calcium channel blocker [e.g. Procardia XL (nifedipine ER), Cardizem LA/Cardizem CD (diltiazemER)]

**OR**

**1.3** One long acting nitrate therapy [e.g. Imdur (isosorbide mononitrate), Isordil (isosorbide dinitrate), Nitro-Time/Nitro-Dur/Nitro-Bid (nitroglycerin ER)]

**AND**

**2** - For Brand Ranexa requests ONLY: Trial and failure to generic ranolazine (verified via paid pharmacy claims or submission of medical records/chart notes)

Product Name:Aspruzyo Sprinkle

Approval Length | 12 month(s)

Guideline Type | Step Therapy

**Approval Criteria**

**1** - History of ONE of the following standard anti-angina treatments:

**1.1** One beta-blocker [e.g. Lopressor (metoprolol), Inderal (propranolol)]

**OR**

**1.2** One calcium channel blocker [e.g. Procardia XL (nifedipine ER), Cardizem LA/Cardizem CD (diltiazemER)]

**OR**

**1.3** One long acting nitrate therapy [e.g. Imdur (isosorbide mononitrate), Isordil (isosorbide dinitrate), Nitro-Time/Nitro-Dur/Nitro-Bid (nitroglycerin ER)]

**AND**

**2** - One of the following:

**2.1** Trial and failure to generic ranolazine (verified via paid pharmacy claims or submission of medical records/chart notes)

**OR**

**2.2** One of the following:

- Patient is 8 years of age or younger
- Patient is unable to swallow the oral tablet (solid formulation) due to swallowing difficulties

## **2 . Revision History**

Date	Notes
8/4/2022	Added Aspruzyo Sprinkle as target. Updated guideline name to Ranolazine Products

Rayos

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99523
<b>Guideline Name</b>	Rayos
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Rayos	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following:  1.1 The requested drug must be used for a Food and Drug Administration (FDA)-approved indication	

**OR**

**1.2** The use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- American Hospital Formulary Service Drug Information
- National Comprehensive Cancer Network Drugs and Biologics Compendium
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**AND**

**2** - The drug is being prescribed for a medically accepted indication that is recognized as a covered benefit by the applicable health plan's program\*

**AND**

**3** - Submission of medical records (e.g. chart notes, laboratory values) or claims history documenting an intolerance to generic prednisone tablets which is unable to be resolved with attempts to minimize the adverse effects where appropriate

**AND**

**4** - History of failure, contraindication, or intolerance to TWO the following:

- Dexamethasone tablet, oral solution
- Hydrocortisone tablet
- Methylprednisolone tablet
- Prednisolone tablet, oral solution

Notes

\*Note: Medications used solely for anti-obesity/weight loss, cosmetic (e.g., alopecia, actinic keratosis, vitiligo), erectile dysfunction, and sexual dysfunction purposes are NOT medically accepted indications and are NOT recognized as a covered benefit. Erectile dysfunction drugs (Cialis/Tadalafil) are covered for clinical diagnoses other than ED.

## 2 . Revision History

Date	Notes
5/14/2021	Arizona Medicaid 7.1 Implementation

Reblozyl (luspatercept-aamt)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135344
<b>Guideline Name</b>	Reblozyl (luspatercept-aamt)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Reblozyl	
Diagnosis	Beta Thalassemia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Both of the following:	

1.1.1 Diagnosis of beta thalassemia major

**AND**

1.1.2 Patient requires regular red blood cell (RBC) transfusions

**OR**

1.2 Diagnosis of transfusion-dependent beta thalassemia

**AND**

2 - Prescribed by or in consultation with one of the following:

- Hematologist
- Oncologist

Product Name:Reblozyl	
Diagnosis	Beta Thalassemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of a positive clinical response to therapy (e.g., reduction in RBC transfusion burden)	

Product Name:Reblozyl	
Diagnosis	Myelodysplastic Syndromes, Myelodysplastic/Myeloproliferative Neoplasm (MDS-RS, MDS/MPN-RS-T)
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following diagnoses:</p> <p>1.1 Very low-to intermediate-risk myelodysplastic syndrome with ring sideroblasts (MDS-RS)</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 Myelodysplastic or myeloproliferative neoplasm with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has failed an erythropoiesis stimulating agent [e.g., Epogen (epoetin alfa), Aranesp (darbepoetin)]</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient requires transfusions of 2 or more red blood cell (RBC) units over 8 weeks</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Hematologist</li> <li>• Oncologist</li> </ul>	

Product Name:Reblozyl	
Diagnosis	Myelodysplastic Syndromes
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of very low- to intermediate-risk myelodysplastic syndromes (MDS)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient does not have previous erythropoiesis stimulating agent use (ESA-naïve)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient requires transfusions of 2 or more red blood cell (RBC) units over 8 weeks</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Hematologist</li> <li>• Oncologist</li> </ul>	

Product Name:Reblozyl	
Diagnosis	Myelodysplastic Syndromes, Myelodysplastic/Myeloproliferative Neoplasm
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Documentation of a positive clinical response to therapy (e.g., RBC transfusion independence, improvement in hemoglobin levels)

## 2 . Revision History

Date	Notes
10/23/2023	Added criteria for new indication of treatment of anemia without previous erythropoiesis stimulating agent use (ESA-naïve).

Recorlev (levoketoconazole)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-102891
<b>Guideline Name</b>	Recorlev (levoketoconazole)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/4/2022
-----------------	----------

## 1 . Criteria

Product Name:Recorlev	
Diagnosis	Cushing's Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Both of the following: <b>1.1</b> Diagnosis of Cushing's disease	

**AND**

**1.2 ONE** of the following:

- Patient is not a candidate for pituitary surgery
- Pituitary surgery has not been curative

Product Name:Recorlev	
Diagnosis	Cushing's Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive response to therapy	

Product Name:Recorlev	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Recorlev will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.	

Product Name:Recorlev
-----------------------

Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
2/3/2022	New Program (mirrors Isturisa PA criteria)

Rectiv

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99492
<b>Guideline Name</b>	Rectiv
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Rectiv	
Diagnosis	Pain Associated with Chronic Anal Fissures
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderate to severe pain associated with chronic anal fissures	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Regranex

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-102898
<b>Guideline Name</b>	Regranex
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/3/2022
-----------------	----------

## 1 . Criteria

Product Name:Regranex	
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Patient has a lower extremity diabetic neuropathic ulcer	

## 2 . Revision History

Date	Notes
2/3/2022	Removed t/f Santyl prerequisite

Relyvrio (sodium phenylbutyrate and taurursodiol)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-120433
<b>Guideline Name</b>	Relyvrio (sodium phenylbutyrate and taurursodiol)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Relyvrio	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting diagnosis of amyotrophic lateral sclerosis (ALS)	

**AND**

**2** - Diagnosis of ALS is further supported by neurogenic changes in electromyography (EMG)

**AND**

**3** - Patient has had ALS symptoms for less than or equal to 18 months

**AND**

**4** - Patient has a percent (%) forced vital capacity (% FVC) or slow vital capacity (% SVC) greater than or equal to 60% at the start of treatment

**AND**

**5** - Patient does not require permanent noninvasive ventilation or invasive ventilation

**AND**

**6** - Prescribed by or in consultation with a neurologist with expertise in the diagnosis of ALS

Product Name:Relyvrio	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting slowed disease progression from baseline	

**AND**

**2** - Prescribed by or in consultation with a neurologist with expertise in the diagnosis of ALS

## **2 . Revision History**

Date	Notes
1/24/2023	New program

Repatha

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-129083
<b>Guideline Name</b>	Repatha
<b>Formulary</b>	<ul style="list-style-type: none"><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Repatha	
Diagnosis	Heterozygous familial hypercholesterolemia (HeFH), Atherosclerotic cardiovascular disease (ASCVD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following diagnoses:	

**1.1** Heterozygous familial hypercholesterolemia (HeFH) as confirmed by ONE of the following\*:

**1.1.1** BOTH of the following:

**1.1.1.1** Pre-treatment LDL-C (low-density lipoprotein cholesterol) greater than 190 milligrams per deciliter (mg/dL) (greater than 155 mg/dL if less than 16 years of age)

**AND**

**1.1.1.2** ONE of the following:

- Family history of myocardial infarction in first degree relative less than 60 years of age
- Family history of myocardial infarction in second degree relative less than 50 years of age
- Family history of LDL-C greater than 190 mg/dL in first or second degree relative
- Family history of heterozygous or homozygous familial hypercholesterolemia in first or second degree relative
- Family history of tendinous xanthomata and or arcus cornealis in first or second degree relative

**OR**

**1.1.2** BOTH of the following:

**1.1.2.1** Pre-treatment LDL-C greater than 190 mg/dL (greater than 155 mg/dL if less than 16 years of age)

**AND**

**1.1.2.2** Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following:

- Functional mutation in LDL (low-density lipoprotein), apoB (Apolipoprotein B), or PCSK9 (Proprotein convertase subtilisin/kexin type 9) gene\*
- Tendinous xanthomata
- Arcus cornealis before age 45

**OR**

**1.2** Atherosclerotic cardiovascular disease (ASCVD) as confirmed by ONE of the following:

- Acute coronary syndromes
- History of myocardial infarction
- Stable or unstable angina
- Coronary or other arterial revascularization
- Stroke
- Transient ischemic attack
- Peripheral arterial disease presumed to be of atherosclerotic origin

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following (prescription claims history may be used in conjunction as documentation of medication use, dose, and duration):

**2.1** Patient has been receiving at least 12 consecutive weeks of high-intensity statin therapy (i.e. atorvastatin 40-80 mg, rosuvastatin 20-40 mg) and will continue to receive high-intensity statin at maximally tolerated dose

**OR**

**2.2** BOTH of the following:

**2.2.1** Patient is unable to tolerate high-intensity statin as evidenced by ONE of the following intolerable and persistent (i.e. more than 2 weeks) symptoms:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

**AND**

**2.2.2** ONE of the following:

**2.2.2.1** Patient has been receiving at least 12 consecutive weeks of moderate-intensity statin [i.e. atorvastatin 10-20 mg, rosuvastatin 5-10 mg, simvastatin greater than or equal to 20 mg, pravastatin greater than or equal to 40 mg, lovastatin 40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 40 mg twice daily or Livalo (pitavastatin) greater than or equal to 2 mg] and will continue to receive a moderate-intensity statin at maximally tolerated dose

**OR**

**2.2.2.2** Patient has been receiving at least 12 consecutive weeks of low-intensity statin [i.e. simvastatin 10 mg, pravastatin 10-20 mg, lovastatin 20 mg, fluvastatin 20-40 mg, or Livalo (pitavastatin) 1 mg] therapy and will continue to receive a low-intensity statin at maximally tolerated dose

**OR**

**2.3** Patient is unable to tolerate low or moderate, and high intensity statins as evidenced by ONE of the following:

**2.3.1** ONE of the following intolerable and persistent (i.e. more than 2 weeks) symptoms for low or moderate, and high intensity statins:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations less than 10 times upper limit of normal [ULN])

**OR**

**2.3.2** Patient has a labeled contraindication to all statins as documented in medical records

**OR**

**2.3.3** Patient has experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations greater than 10 times ULN

**AND**

**3** - ONE of the following:

**3.1** Submission of medical records (e.g., laboratory values) documenting ONE of the following LDL-C values while on maximally tolerated lipid lowering therapy for a minimum of at least 12 weeks within the last 120 days:

- LDL-C greater than or equal to 100 mg/dL with ASCVD

- LDL-C greater than or equal to 130 mg/dL without ASCVD

**OR**

**3.2** BOTH of the following:

**3.2.1** Submission of medical records (e.g., laboratory values) documenting ONE of the following LDL-C values while on maximally tolerated lipid lowering therapy for a minimum of at least 12 weeks within the last 120 days:

- LDL-C between 55 mg/dL and 99 mg/dL with ASCVD
- LDL-C between 100 mg/dL and 129 mg/dL without ASCVD

**AND**

**3.2.2** Submission of medical records (e.g., chart notes, laboratory values) documenting ONE of the following [prescription claims history may be used in conjunction as documentation of medication use, dose, and duration]:

- Patient has been receiving at least 12 consecutive weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy
- Patient has a history of contraindication or intolerance to ezetimibe

**AND**

**4** - Used as an adjunct to a low-fat diet and exercise

**AND**

**5** - Prescribed by ONE of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

**AND**

**6** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Praluent (alirocumab))

Notes	*Results of prior genetic testing can be submitted as confirmation of diagnosis of HeFH .
-------	---

Product Name:Repatha	
Diagnosis	Heterozygous familial hypercholesterolemia (HeFH), Atherosclerotic cardiovascular disease (ASCVD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient continues to receive statin at maximally tolerated dose (unless patient has documented inability to take statins)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is continuing a low-fat diet and exercise regimen</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by ONE of the following:</p> <ul style="list-style-type: none"> <li>• Cardiologist</li> <li>• Endocrinologist</li> <li>• Lipid specialist</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Submission of medical records (e.g. chart notes, laboratory values) documenting LDL-C (low-density lipoprotein cholesterol) reduction while on Repatha therapy</p> <p style="text-align: center;"><b>AND</b></p>	

**5** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Praluent (alirocumab))

Product Name: Repatha

Diagnosis	Homozygous Familial Hypercholesterolemia (HoFH)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

### Approval Criteria

**1** - Diagnosis of homozygous familial hypercholesterolemia (HoFH) as confirmed by submission of medical records (e.g., chart notes, laboratory values) documenting BOTH of the following:\*

**1.1** ONE of the following:

- Pre-treatment LDL-C (low-density lipoprotein cholesterol) greater than 500 mg/dL (milligrams per deciliter)
- Treated LDL-C greater than 300 mg/dL

**AND**

**1.2** ONE of the following:

- Xanthoma before 10 years of age
- Evidence of heterozygous familial hypercholesterolemia (HeFH) in both parents

**AND**

**2** - Used as an adjunct to a low-fat diet and exercise

**AND**

**3** - Patient is receiving other lipid-lowering therapy (e.g., statin, ezetimibe, LDL [low-density lipoprotein] apheresis)

**AND**

**4** - Prescribed by ONE of the following:

- Cardiologist
- Endocrinologist
- Lipid specialist

**AND**

**5** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Praluent (alirocumab))

Notes

\*Results of prior genetic testing can be submitted as confirmation of diagnosis of HoFH.

Product Name: Repatha

Diagnosis | Homozygous Familial Hypercholesterolemia

Approval Length | 12 month(s)

Therapy Stage | Reauthorization

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Patient is continuing a low-fat diet and exercise regimen

**AND**

**2** - Patient continues to receive other lipid-lowering therapy (e.g., statin, LDL apheresis)

**AND**

**3** - Submission of medical records (e.g. chart notes, laboratory values) documenting LDL-C (low-density lipoprotein cholesterol) reduction while on Repatha therapy

**AND**

**4** - Prescribed by ONE of the following:

- Cardiologist
- Endocrinologist
- Lipid Specialist

**AND**

**5** - Not used in combination with another proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., Praluent (alirocumab))

## 2 . Revision History

Date	Notes
9/1/2023	Update to account for 2022 ACC recommendations of a lower LDL threshold of 55mg/dl for patients with ASCVD at very high risk.

Respiratory Syncytial Virus (RSV) Vaccines

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-161591
<b>Guideline Name</b>	Respiratory Syncytial Virus (RSV) Vaccines
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	1/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Abrysvo, Arexvy, mResvia	
Approval Length	14 days (1 injection per 2 years)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Vaccine is being used for the prevention of lower respiratory tract disease (LRTD) caused by respiratory syncytial virus (RSV)</p> <p style="text-align: center;"><b>AND</b></p>	

**2** - Patient has not received an RSV vaccine (i.e., Abrysvo, Arexvy, mResvia) in the previous 2 years

**AND**

**3** - One of the following:

**3.1** One of the following:

- For ABRYSSVO requests: Age greater than or equal to 60 years\*
- For AREXVY requests: Age greater than or equal to 50 years\*
- For MRESVIA requests: Age greater than or equal to 60 years\*

**OR**

**3.2** One of the following (applies to ABRYSSVO only):

**3.2.1** Both of the following:

- Will be used for active immunization of pregnant individuals at 32 through 36 weeks gestational age
- Will also be used for the prevention of severe LRTD caused by RSV in infants from birth through 6 months of age

**OR**

**3.2.2** Both of the following:

- Patient is 18 to 59 years of age
- Patient is at increased risk for LRTD caused by RSV

**AND**

**4** - For MRESVIA requests ONLY: Trial and failure, intolerance, or contraindication to one of the following: (Applies to MRESVIA only)

- Abrysvo
- Arexvy

Notes	*Prior authorization for ABRYSSO or MRESVIA is NOT required for patients 60 years and older. *Prior authorization for AREXVY is NOT required for patients 50 years and older.
-------	---

## 2 . Revision History

Date	Notes
12/6/2024	Added criteria for Abrysvo due to new indication

Retinal Vascular Disease Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-325193
<b>Guideline Name</b>	Retinal Vascular Disease Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	8/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Preferred: Cimerli	
Diagnosis	Neovascular (wet) age-related macular degeneration (nAMD), Macular edema following retinal vein occlusion (RVO), Diabetic macular edema (DME), Diabetic retinopathy (DR), Myopic choroidal neovascularization (mCNV)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1** - Submission of medical records (e.g., chart notes) documenting one of the following diagnoses:

- Neovascular (wet) age-related macular degeneration (nAMD)
- Macular edema following retinal vein occlusion (RVO)
- Diabetic macular edema (DME)
- Diabetic retinopathy (DR)
- Myopic choroidal neovascularization (mCNV)

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting treatment for a minimum of 90 days with compounded Avastin prepared by a 503(B) Outsourcing Facility has been ineffective in improvement of visual acuity, or not tolerated, or contraindicated (paid pharmacy claims may be used in conjunction with submitted documentation)

**AND**

**3** - Prescribed by or in consultation with an ophthalmologist experienced in the treatment of retinal diseases

Product Name: Preferred: Pavblu	
Diagnosis	Neovascular (wet) age-related macular degeneration (nAMD), Macular edema following retinal vein occlusion (RVO), Diabetic macular edema (DME), Diabetic retinopathy (DR)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting one of the following diagnoses:</p> <ul style="list-style-type: none"> <li>• Neovascular (wet) age-related macular degeneration (nAMD)</li> <li>• Macular edema following retinal vein occlusion (RVO)</li> <li>• Diabetic macular edema (DME)</li> </ul>	

- Diabetic retinopathy (DR)

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting treatment for a minimum of 90 days with compounded Avastin prepared by a 503(B) Outsourcing Facility has been ineffective in improvement of visual acuity, or not tolerated, or contraindicated (paid pharmacy claims may be used in conjunction with submitted documentation)

**AND**

**3** - Prescribed by or in consultation with an ophthalmologist experienced in the treatment of retinal diseases

Product Name:Preferred: Cimerli, Pavblu	
Diagnosis	All indications listed above
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., Improvement in Best Corrected Visual Acuity (BCVA) compared to baseline, stable vision)</p>	

Product Name:Non-Preferred: Eylea Injectable Vial	
Diagnosis	Retinopathy of Prematurity (ROP)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of retinopathy of prematurity (ROP)

**AND**

2 - ONE of the following:

- Patient gestational age at birth less than or equal to 32 weeks
- Patient birth weight less than or equal to 1500 grams

**AND**

3 - Patient weight greater than 800 grams on day of treatment

**AND**

4 - Submission of medical records (e.g., chart notes) documenting retinopathy of prematurity (ROP) is present in at least one eye with one of the following classifications:

- ROP zone 1, stage 1 plus, 2 plus, 3, or 3 plus
- ROP zone 2, stage 2 plus or 3 plus
- AP - ROP (aggressive posterior ROP)

**AND**

5 - Prescribed by or in consultation with an ophthalmologist experienced in the treatment of retinal diseases

Product Name:Non-Preferred: Eylea Injectable Vial

Diagnosis	Retinopathy of Prematurity (RoP)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy as evidenced by the absence of active ROP and unfavorable structural outcomes (e.g., retinal detachment, macular dragging, macular fold, retrolental opacity)

Product Name:Non-Preferred\*: Beovu, Byooviz, Eylea, Eylea HD, Lucentis, Susvimo, Vabysmo, Opuviz, Yesafili, Ahzantive, Enzeevu, and other newly launched retinal vascular disease agents

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred products are Cimerli and Pavblu. \*\*Please note: The drug(s) listed above may require additional review.

Notes	*Patients must use preferred retinal vascular disease agent(s).
-------	---

**2 . Revision History**

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, updated criteria throughout. Updated NP section verbiage.

Revcovi - AZ

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99639
<b>Guideline Name</b>	Revcovi - AZ
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Revcovi	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of severe combined immunodeficiency disease (SCID)  <b>AND</b>	

**2** - Deficiency of adenosine deaminase is confirmed by one of the following:

- Deficiency or absence of adenosine deaminase (ADA) in plasma, lysed erythrocytes, fibroblasts (cultured from amniotic fluid), or chorionic villus
- Increase in deoxyadenosine triphosphate (dATP) levels in erythrocyte lysates compared to laboratory standard
- Decrease in ATP (Adenosine triphosphate) concentration in erythrocytes
- Molecular genetic confirmation of mutations in both alleles of the ADA1 gene
- Positive screening by T cell receptor excision circles (TRECs)

**AND**

**3** - One of the following:

- Patient is not a suitable candidate for hematopoietic cell transplantation (HCT)
- Patient has failed HCT
- Patient is awaiting HCT

**AND**

**4** - Dosing is in accordance with the United States Food and Drug Administration approved labeling

Product Name: Revcovi	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has previously received treatment with Revcovi (elapegedemase) therapy	
<b>AND</b>	
2 - Patient has experienced a positive clinical response to therapy (e.g., normalization of plasma ADA activity, erythrocyte dATP levels, improvement of disease symptoms, etc.)	

**AND**

**3** - Dosing is in accordance with the United States Food and Drug Administration approved labeling

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Reyvow - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99548
<b>Guideline Name</b>	Reyvow - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Reyvow	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderate to severe migraine headaches with or without aura  <b>AND</b>	

**2** - Used for acute treatment of migraine

**AND**

**3** - Patient is 18 years of age or older

**AND**

**4** - Documentation of a one month trial resulting in therapeutic failure, contraindication, or intolerance to **THREE** of the following:

- naratriptan tablets
- rizatriptan tablets/ODT (oral disintegrating tablets)
- sumatriptan tablets/auto injection/cartridge or Imitrex nasal spray (Brand only)
- zolmitriptan tablets/ODT

**AND**

**5** - Prescribed by or in consultation with one of the following specialists with expertise in the acute treatment of migraine:

- Neurologist
- Pain Specialist
- Headache Specialist\*

**AND**

**6** - Prescriber attests to **ALL** of the following:

- Patient has been informed the use of Reyvow may result in significant CNS impairment, and may impact the patient's ability to drive or operate machinery for 8 hours after each dose
- If used concurrently with a benzodiazepine or other drugs that could potentially cause central nervous system (CNS) depression, the prescriber has acknowledged that they have completed an assessment of increased risk for sedation and other cognitive and/or neuropsychiatric adverse events
- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

**AND**

**7** - Both of the following:

**7.1** One of the following

**7.1.1** The patient must have a history of therapeutic failure, contraindication, or intolerance to **THREE** of the following:

- Amitriptyline (Elavil)\*\*
- A beta-blocker (i.e., atenolol, metoprolol, nadolol, propranolol, or timolol)\*\*
- Divalproex sodium [Depakote/Depakote ER (extended-release)]\*\*
- Topiramate (Topamax)\*\*
- VENLAFAXINE [EFFEXOR/EFFEXOR XR (EXTENDED-RELEASE)]\*\*

**OR**

**7.1.2** The patient must be currently treated with one of the following prophylactic therapies unless there is a contraindication or intolerance to **ALL**:

- Amitriptyline (Elavil)\*\*
- A beta-blocker (i.e., atenolol, metoprolol, nadolol, propranolol, or timolol)\*\*
- Divalproex sodium [Depakote/Depakote ER (extended-release)]\*\*
- Topiramate (Topamax)\*\*
- Venlafaxine [Effexor/Effexor XR (extended-release)]\*\*

**AND**

**7.2** Both of the Following

**7.2.1** History of a therapeutic failure after 3 month trial, contraindication, or intolerance to two of the following biologic calcitonin gene-related peptide receptor (CGRP) antagonists for preventive treatment of migraine

- Ajovy (fremanezumab)
- Emgality (galcanezumab)
- Aimovig (erenumab)

**AND**

**7.2.2** History of a therapeutic failure, contraindication, or intolerance to Ubrelvy

Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS) **Drugs may require PA
-------	---

Product Name:Reyvow	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following specialists with expertise in the acute treatment of migraine:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Pain Specialist</li> <li>• Headache Specialist*</li> </ul>	
Notes	*Headache specialists are physicians certified by the United Council f or Neurologic Subspecialties (UCNS)

## 2 . Revision History

Date	Notes
7/13/2021	Updated Guideline

Rezdiffra (resmetirom)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-149969
<b>Guideline Name</b>	Rezdiffra (resmetirom)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Rezdiffra	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of metabolic dysfunction-associated steatohepatitis (MASH), formerly known as nonalcoholic steatohepatitis (NASH)	

**AND**

**2** - Patient does not have cirrhosis (e.g., decompensated cirrhosis)

**AND**

**3** - Submission of medical records (e.g., chart notes) confirming diagnosis has been confirmed by one of the following:

- FibroScan-aspartate aminotransferase (FAST)
- MRI-aspartate aminotransferase (MAST)
- Liver biopsy

**AND**

**4** - Submission of medical records (e.g., chart notes) confirming disease is fibrosis stage F2 or F3 as confirmed by one of the following:

- FibroScan
- Fibrosis-4 index (FIB-4)
- Magnetic Resonance Elastography (MRE)

**AND**

**5** - Presence of greater than or equal to 3 metabolic risk factors (e.g., Type 2 diabetes, hypertension, obesity)

**AND**

**6** - Submission of medical records (e.g., chart notes) confirming drug is used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community based program)

**AND**

**7** - Prescribed by or in consultation with one of the following:

- Gastroenterologist
- Hepatologist

Product Name:Rezdiffra	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient demonstrates positive response to therapy (e.g., NASH resolution, fibrosis stage improvement, etc.)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Submission of medical records (e.g., chart notes) confirming drug will continue to be used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community based program)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Submission of medical records (e.g., chart notes) confirming patient has not progressed to cirrhosis</p>	

**2 . Revision History**

Date	Notes
8/29/2024	New program

Rezurock (belumosudil)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-103329
<b>Guideline Name</b>	Rezurock (belumosudil)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Rezurock	
Diagnosis	Chronic graft-versus-host disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic graft-versus-host disease	

**AND**

**2** - Trial and failure of two or more lines of systemic therapy (e.g., corticosteroids, mycophenolate, etc.)

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Hematologist
- Oncologist
- Physician experienced in the management of transplant patients

Product Name:Rezurock	
Diagnosis	Chronic graft-versus-host disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on therapy	

Product Name:Rezurock	
Diagnosis	Chronic graft-versus-host disease - Twice daily (BID) Therapy
Approval Length	12 month(s)
Guideline Type	Quantity Limit
<b>Approval Criteria</b>	
1 - Patient is using medication concomitantly with one of the following:	

- Strong CYP3A inducer (e.g., carbamazepine, phenobarbital, phenytoin, rifampin)
- Proton pump inhibitor (e.g., omeprazole, pantoprazole, lansoprazole)

## 2 . Revision History

Date	Notes
2/3/2022	New Program

Rezzayo (rezafungin)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-133806
<b>Guideline Name</b>	Rezzayo (rezafungin)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Rezzayo	
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of candidemia or invasive candidiasis with limited or no alternative options  <b>AND</b>	

**2** - Patient is 18 years of age or older

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid pharmacy claims confirming trial and failure, contraindication or intolerance to one of the following:

- generic caspofungin
- generic micafungin

## **2 . Revision History**

Date	Notes
9/26/2023	New program

Rhofade

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99494
<b>Guideline Name</b>	Rhofade
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Rhofade	
Diagnosis	Persistent erythema associated with rosacea
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of persistent erythema associated with rosacea	

**AND**

**2 - ONE of the following:**

**2.1 History of a 30 day or longer trial and failure of one of the following:**

- metronidazole cream, gel, or lotion
- azelaic acid gel

**OR**

**2.2 Contraindication or intolerance to both of the following:**

- metronidazole cream, gel, or lotion
- azelaic acid gel

Product Name: Rhofade	
Diagnosis	Persistent erythema associated with rosacea
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of a positive clinical response to Rhofade therapy	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Rinvoq (upadacitinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-330209
<b>Guideline Name</b>	Rinvoq (upadacitinib)
<b>Formulary</b>	Medicaid - Arizona (AZM, AZMREF, AZMDDD)

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Rinvoq	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:\*\*

**3.1** History of failure to a 3 month trial of one non-biologic disease modifying anti-rheumatic drug (DMARD) (e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine) at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced\*

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following:\*\*\*

A preferred adalimumab biosimilar or Enbrel (etanercept)

infliximab

Orencia (abatacept)

Xeljanz oral tablet (tofacitinib) (IR or XR)

A preferred tocilizumab biosimilar

**AND**

**4** - Not used in combination with other Janus kinase (JAK) inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)\*

Notes

\*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

\*\*PA may be required

\*\*\*Patients requesting initial authorization who were established on th

	erapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.
--	--

Product Name: Rinvoq	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)*</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with a rheumatologist</p>	
Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name: Rinvoq, Rinvoq LQ	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of active polyarticular juvenile idiopathic arthritis (PJIA)

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:\*\*

**3.1** Minimum duration of a 6-week trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:\*

methotrexate

leflunomide

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following:\*\*\*

A preferred adalimumab biosimilar or Enbrel (etanercept)

Orencia (abatacept)

Xeljanz (tofacitinib) oral tablet

A preferred tocilizumab biosimilar

**AND**

**4** - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)\*

Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily). **PA may be required ** *Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.
-------	--

Product Name:Rinvoq, Rinvoq LQ	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)*</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a rheumatologist</p>	
Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name:Rinvoq, Rinvoq LQ	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming a diagnosis of active psoriatic arthritis (PsA)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with one of the following:</p> <p style="padding-left: 40px;">Dermatologist</p> <p style="padding-left: 40px;">Rheumatologist</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:**</p> <p><b>3.1</b> History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced*</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3.2</b> History of failure, contraindication, or intolerance to ALL of the following:***</p> <p style="padding-left: 40px;">A preferred adalimumab biosimilar or Enbrel (etanercept)</p> <p style="padding-left: 40px;">infliximab</p> <p style="padding-left: 40px;">Orencia (abatacept)</p> <p style="padding-left: 40px;">Otezla (apremilast)</p> <p style="padding-left: 40px;">Xeljanz (tofacitinib) oral tablet (IR or XR)</p>	

A preferred ustekinumab biosimilar

**AND**

**4** - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)\*

Notes

\*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily). \*\*PA may be required \*\*  
\*Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.

Product Name: Rinvoq, Rinvoq LQ

Diagnosis Psoriatic Arthritis (PsA)

Approval Length 12 month(s)

Therapy Stage Reauthorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy

**AND**

**2** - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)\*

**AND**

**3** - Prescribed by or in consultation with one of the following:

<p>Dermatologist</p> <p>Rheumatologist</p>	
Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name: Rinvoq	
Diagnosis	Ankylosing Spondylitis (AS), Non-radiographic Axial Spondyloarthritis (nr-AxSpA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - Submission of medical records (e.g., chart notes) confirming one of the following diagnoses:</b></p> <p>Active ankylosing spondylitis (AS)</p> <p>Active non-radiographic axial spondyloarthritis (nr-AxSpA)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2 - Prescribed by or in consultation with a rheumatologist</b></p> <p style="text-align: center;"><b>AND</b></p> <p><b>3 - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:**</b></p> <p><b>3.1 Trial and failure, contraindication, or intolerance to TWO nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen)</b></p>	

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following.\*\*\*

A preferred adalimumab biosimilar or Enbrel (etanercept)

infliximab

Xeljanz (tofacitinib) oral tablet (IR or XR)

**AND**

**4** - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)\*

Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).  **PA may be required  ***Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.
-------	--

Product Name:Rinvoq	
Diagnosis	Ankylosing Spondylitis (AS), Non-radiographic Axial Spondyloarthritis (nr-AxSpA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy	

**AND**

**2** - Not used in combination with other JAK inhibitors, biologic DMARDs, or potent immunosuppressants (e.g., azathioprine or cyclosporine)\*

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Notes

\*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name: Rinvoq

Diagnosis Atopic Dermatitis (AD)

Approval Length 6 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderate to severe atopic dermatitis (AD)

**AND**

**2** - Patient is 12 years of age or older

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting one of the following:

Involvement of at least 10% body surface area (BSA)

SCORing Atopic Dermatitis (SCORAD) index value of at least 25

**AND**

**4** - Prescribed by or in consultation with one of the following:

Dermatologist

Allergist/Immunologist

**AND**

**5** - Submission of medical records (e.g., chart notes) or paid claims history documenting ALL of the following:\*\*

**5.1** History of failure, contraindication, or intolerance to ALL of the following topical therapies:\*\*\*

One topical calcineurin inhibitor [e.g., Elidel (pimecrolimus), Protopic (tacrolimus)]

Eucria (crisaborole)

Opzelura (ruxolitinib) cream

Vtama (tapinarof) cream

Zoryve (roflumilast) 0.15% cream

**AND**

**5.2** Submission of medical records (e.g., chart notes) or paid claims history documenting trial and failure of a minimum 12-week supply of Dupixent (dupilumab)\*\*\*

**AND**

**5.3** Submission of medical records (e.g., chart notes) or paid claims history documenting trial and failure of a minimum 12-week supply of Adbry (tralokinumab-ldrm)\*\*\*

**AND**

**6** - Not used in combination with other JAK inhibitors, biologic immunomodulators (e.g., Dupixent, Adbry), or other immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily). ** PA may be required. ***Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.
-------	--

Product Name: Rinvoq	
Diagnosis	Atopic Dermatitis (AD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by at least ONE of the following:</p> <ul style="list-style-type: none"><li>Reduction in body surface area involvement from baseline</li><li>Reduction in SCORing Atopic Dermatitis (SCORAD) index value from baseline</li></ul> <p><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"><li>Dermatologist</li><li>Allergist/Immunologist</li></ul>	

**AND**

**3** - Not used in combination with other JAK inhibitors, biologic immunomodulators (e.g., Dupixent, Adbry), or other immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes

\*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name: Rinvoq

Diagnosis | Ulcerative Colitis (UC)

Approval Length | 12 month(s)

Therapy Stage | Initial Authorization

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis (UC)

**AND**

**2** - Prescribed by or in consultation with a gastroenterologist

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:\*\*

**3.1** Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

6-mercaptopurine

Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)

Azathioprine

Corticosteroids (e.g., prednisone)

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following:\*\*\*

A preferred adalimumab biosimilar

infliximab

Xeljanz oral tablet (tofacitinib) (IR or XR)

A preferred ustekinumab biosimilar

**AND**

**4** - Not used in combination with other JAK inhibitors, biological therapies for UC, or with potent immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).  **PA may be required  ***Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.
-------	--

Product Name:Rinvoq	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with a gastroenterologist

**AND**

3 - Not used in combination with other JAK inhibitors, biological therapies for UC, or with potent immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes

\*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name: Rinvoq

Diagnosis Crohn's Disease (CD)

Approval Length 12 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's disease (CD)

**AND**

2 - Prescribed by or in consultation with a gastroenterologist

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims documenting BOTH of the following:\*\*

**3.1** Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

6-mercaptopurine

Azathioprine

Methotrexate

Corticosteroids (e.g., prednisone)

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following:\*\*\*

A preferred adalimumab biosimilar

infliximab

A preferred ustekinumab biosimilar

**AND**

**4** - Not used in combination with other JAK inhibitors, biological therapies for CD, or with potent immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes	<p>*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).</p> <p>**PA may be required</p> <p>***Patients requesting initial authorization who were established on therapy via the receipt of a manufacturer supplied sample at no cost in the prescriber's office or any form of assistance from manufacturer sponsored programs shall be required to meet initial authorization criteria as if patient were new to therapy.</p>
-------	---

Product Name:Rinvoq

Diagnosis	Crohn's Disease (CD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by at least one of the following:</p> <p style="padding-left: 40px;">Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</p> <p style="padding-left: 40px;">Reversal of high fecal output state</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a gastroenterologist</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Not used in combination with other JAK inhibitors, biological therapies for CD, or with potent immunosuppressants (e.g., azathioprine, cyclosporine)*</p>	
Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

Product Name: Rinvoq	
Diagnosis	Giant Cell Arteritis (GCA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of giant cell arteritis (GCA)

**AND**

2 - Prescribed by or in consultation with a rheumatologist

**AND**

3 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to BOTH of the following:

A glucocorticoid (e.g., prednisone)

A preferred tocilizumab biosimilar

**AND**

4 - Not used in combination with other JAK inhibitors, biological therapies, or with potent immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes	*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).
-------	---

Product Name: Rinvoq	
Diagnosis	Giant Cell Arteritis (GCA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy</p>	

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

**AND**

**3** - Not used in combination with other JAK inhibitors, biological therapies, or with potent immunosuppressants (e.g., azathioprine, cyclosporine)\*

Notes

\*Rinvoq may be used with concomitant methotrexate, topical or inhaled corticosteroids, and/or low stable dosages of oral corticosteroids (equivalent to 10 mg or less of prednisone daily).

## 2 . Background

Clinical Practice Guidelines			
Table 1. Relative potencies of topical corticosteroids [5]			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment, gel	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream, lotion	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05

	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05
	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
	Triamcinolone acetonide	Cream, ointment	0.5
Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment, lotion	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream, lotion	0.1
	Triamcinolone acetonide	Cream, ointment, lotion	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1

Hydrocortisone acetate	Cream, ointment	0.5-1
------------------------	-----------------	-------

### 3 . Revision History

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, added criteria for new G CA indication, updated criteria throughout

Rituximab

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325194
<b>Guideline Name</b>	Rituximab
<b>Formulary</b>	Medicaid - Arizona (AZM, AZMREF, AZMDDD)

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:PREFERRED: Riabni, Ruxience	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	1 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of moderately- to severely-active rheumatoid arthritis	

**AND**

**2** - Paid claims or submission of medical records (e.g., chart notes) confirming a minimum duration of a 3-month trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:

methotrexate

leflunomide

sulfasalazine

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming that medication is used in combination with methotrexate

Product Name: PREFERRED: Riabni, Ruxience	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	1 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:	
Reduction in the total active (swollen and tender) joint count from baseline	
Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline	
<b>AND</b>	
<b>2</b> - At least 16 weeks have elapsed since last course of therapy	

Product Name: PREFERRED: Riabni, Ruxience	
Diagnosis	Non-Hodgkin's Lymphoma
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1 - One of the following:</b></p> <p><b>1.1 Both of the following:</b></p> <p>Diagnosis of diffuse large B-cell, CD20-positive, non-Hodgkin's lymphoma</p> <p>Used as first-line treatment in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) or other anthracycline-based chemotherapy regimens</p> <p style="text-align: center;"><b>OR</b></p> <p><b>1.2 Both of the following:</b></p> <p>Diagnosis of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma</p> <p>Used as first-line treatment in combination with chemotherapy</p> <p style="text-align: center;"><b>OR</b></p> <p><b>1.3 All of the following:</b></p> <p>Diagnosis of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma</p> <p>Patient achieved a complete or partial response to a rituximab product in combination with chemotherapy</p> <p>Followed by rituximab used as monotherapy for maintenance therapy</p> <p style="text-align: center;"><b>OR</b></p>	

**1.4** Both of the following:

**1.4.1** Diagnosis of low-grade, CD20-positive, B-cell non-Hodgkin's lymphoma

**AND**

**1.4.2** One of the following:

Patient has stable disease following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/ prednisone) chemotherapy

Patient achieved a partial or complete response following first-line treatment with CVP (cyclophosphamide, vincristine, prednisolone/ prednisone) chemotherapy

**OR**

**1.5** Diagnosis of relapsed or refractory, low grade or follicular CD20-positive, B-cell non-Hodgkin's lymphoma.

**OR**

**1.6** All of the following (off-label for Riabni):

**1.6.1** Diagnosis of one of the following previously untreated, advanced stage indications:

CD-20-positive diffuse large B-cell lymphoma (DLBCL)

Burkitt lymphoma (BL)

Burkitt-like lymphoma (BLL)

Mature B-cell acute leukemia (B-AL)

**AND**

**1.6.2** Patient is 6 months of age or older

**AND**

**1.6.3** Used in combination with chemotherapy

Product Name:PREFERRED: Riabni, Ruxience	
Diagnosis	Chronic Lymphocytic Leukemia
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic lymphocytic leukemia	
<b>AND</b>	
2 - Used in combination with fludarabine and cyclophosphamide	

Product Name:PREFERRED: Riabni, Ruxience	
Diagnosis	Wegener's Granulomatosis and Microscopic Polyangiitis
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - One of the following diagnoses:	
Granulomatosis with Polyangiitis (GPA) (Wegener's Granulomatosis)	
Microscopic Polyangiitis	

**AND**

**2** - Used in combination with glucocorticoids (e.g., prednisone)

Product Name:PREFERRED: Riabni, Ruxience

Diagnosis	Pemphigus Vulgaris
-----------	--------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of moderate to severe Pemphigus Vulgaris

Product Name:PREFERRED: Riabni, Ruxience

Diagnosis	Pemphigus Vulgaris
-----------	--------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Patient demonstrates positive clinical response to therapy

Product Name:PREFERRED: Riabni, Ruxience

Diagnosis	Waldenstrom's macroglobulinemia
-----------	---------------------------------

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of relapsed/refractory Waldenstrom's macroglobulinemia (off-label)

Product Name:PREFERRED: Riabni, Ruxience

Diagnosis	Immune or Idiopathic Thrombocytopenic Purpura (Off-Label)
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of immune or idiopathic thrombocytopenic purpura (off-label)

**AND**

2 - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to at least ONE of the following:

Glucocorticoids (e.g., prednisone, methylprednisolone)

Immunoglobulins (e.g., IVIg)

Splenectomy

**AND**

3 - Documented platelet count of less than  $50 \times 10^9 / L$

Product Name:NON-PREFERRED\*: Rituxan, Truxima, and newly launched rituximab products

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
-----------------	--

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred products are Riabni and Ruxience. \*\*Please note: The drug(s) listed above may require additional review.

Notes

\*Patients must use preferred rituximab products.

## 2 . Revision History

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, updated criteria throughout. Updated NP section verbiage.

Ryaltris

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116154
<b>Guideline Name</b>	Ryaltris
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Ryaltris	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Trial and failure to both of the following as separate agents: <ul style="list-style-type: none"><li>generic mometasone nasal spray</li><li>azelastine or olopatadine nasal spray</li></ul>	

## 2 . Revision History

Date	Notes
10/27/2022	New program

Rystiggo (rozanolixizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-150050
<b>Guideline Name</b>	Rystiggo (rozanolixizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Rystiggo	
Diagnosis	Generalized Myasthenia Gravis (gMG)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of generalized myasthenia gravis (gMG)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting ONE of the following:

**2.1** Both of the following:

**2.1.1** Patient is anti-acetylcholine receptor (AChR) antibody positive

**AND**

**2.1.2** Both of the following:

**2.1.2.1** Trial and failure, contraindication, or intolerance to two immunosuppressive therapy (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus) (May be verified via paid pharmacy claims)

**AND**

**2.1.2.2** Trial and failure, contraindication, or intolerance to one of the following: (May be verified via paid pharmacy claims)

- Chronic plasmapheresis or plasma exchange (PE)
- Intravenous immunoglobulin (IVIG) or immunoglobulin (IG) therapy

**OR**

**2.2** Both of the following:

**2.2.1** Patient is anti-muscle-specific tyrosine kinase (MuSK) antibody positive

**AND**

**2.2.2** Both of the following:

**2.2.2.1** Trial and failure, contraindication, or intolerance to two immunosuppressive therapies (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus) (May be verified via paid pharmacy claims)

**AND**

**2.2.2.2** Trial and failure, contraindication, or intolerance to one of the following: (May be verified via paid pharmacy claims)

- Chronic plasmapheresis or plasma exchange (PE)
- Intravenous immunoglobulin (IVIG) or immunoglobulin (IG) therapy
- Rituximab

**AND**

**3** - Prescribed by or in consultation with a neurologist

Product Name:Rystiggo	
Diagnosis	Generalized Myasthenia Gravis (gMG)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy	

## 2 . Revision History

Date	Notes
7/26/2024	Added new GPIs. Updated embedded step requirement, added submission of records/paid claims.

Rytelo (imetelstat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152667
<b>Guideline Name</b>	Rytelo (imetelstat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Rytelo	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or paid claims documenting ALL of the following:  1.1 Diagnosis of myelodysplastic syndrome	

**AND**

**1.2** Disease is low to intermediate-1 risk

**AND**

**1.3** All of the following:

- Hemoglobin less than 10 g/dL
- Baseline absolute neutrophil count of  $1.5 \times 10^9$  /L or greater
- Baseline platelet count of  $75 \times 10^9$  /L or greater

**AND**

**1.4** Both of the following:

- Patient does not have a confirmed mutation with deletion 5q [del(5q)]
- Patient has not received prior treatment with Revlimid (lenalidomide) or hypomethylating agents (e.g., azacitidine, decitabine)

**AND**

**1.5** Patient requires 4 or more red blood cell units over 8 weeks

**AND**

**1.6** One of the following:

- Previous treatment with an erythropoiesis stimulating agent shows no response
- Previous treatment with an erythropoiesis stimulating agent shows loss of response
- Patient is ineligible for treatment with an erythropoiesis stimulating agent

Product Name:Rytelo	
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on therapy</p>	

**2 . Revision History**

Date	Notes
8/27/2024	New program

Samsca

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99642
<b>Guideline Name</b>	Samsca
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Samsca, generic tolvaptan	
Approval Length	30 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following: <ul style="list-style-type: none"><li>Diagnosis of clinically significant euvolemic hyponatremia</li><li>Diagnosis of clinically significant hypervolemic hyponatremia</li></ul>	

**AND**

**2** - Patient has not responded to fluid restriction

**AND**

**3** - Treatment has been initiated or re-initiated in a hospital setting prior to discharge

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Sedative Hypnotics - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-211207
<b>Guideline Name</b>	Sedative Hypnotics - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Rozerem	
Approval Length	12 month(s)
Guideline Type	Step Therapy
<b>Approval Criteria</b>  1 - History of failure, contraindication, or intolerance to a trial of at least two of the following preferred agents: <ul style="list-style-type: none"><li>Eszopiclone (Generic Lunesta)</li><li>Zolpidem/Zolpidem ER (Generic Ambien/Ambien CR)</li><li>Temazepam 15/30mg capsules (Generic Restoril)</li></ul>	

Product Name: Non-Preferred Drugs: Brand Ambien, Brand Ambien CR, Edluar, Brand Intermezzo, generic zolpidem SL tablets, Zolpimist, Belsomra, Dayvigo, estazolam, flurazepam, Brand Halcion, generic triazolam, Brand Lunesta, Quviviq, Brand Restoril, generic temazepam 7.5 mg and 22.5 mg capsules, generic ramelteon, Brand Silenor, generic doxepin, generic zaleplon

Diagnosis	Non-Preferred
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - History of failure, contraindication, or intolerance to a trial of at least two of the following preferred agents:

- Eszopiclone (Generic Lunesta)
- Zolpidem/Zolpidem ER (Generic Ambien/Ambien CR)
- Temazepam 15/30mg capsules (Generic Restoril)

**AND**

2 - For generic ramelteon requests ONLY, patient must have tried and failed Brand Rozerem

Product Name: Brand Ambien, generic zolpidem, Brand Ambien CR, generic zolpidem ER, Edluar, Brand Intermezzo, generic zolpidem SL tablets, Zolpimist, Belsomra, Dayvigo, estazolam, flurazepam, Brand Halcion, generic triazolam, Brand Lunesta, generic eszopiclone, Quviviq, Brand Restoril, generic temazepam, Brand Rozerem, generic ramelteon, Brand Silenor, generic doxepin, generic zaleplon

Diagnosis	Reject 75: Greater than 1 hypnotic in 30 days
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - The requested medication is being used to adjust the dose of the drug

**OR**

**2** - The requested medication will be used in place of the previously prescribed drug, and not in addition to it

**OR**

**3** - The requested medication dosage form will be used in place of the previously prescribed medication dosage form, and not in addition to it

**OR**

**4** - The physician attests they are aware of the multiple sedative hypnotics prescribed to the patient and feels treatment with both medications is medically necessary (Document rationale for use)

Product Name: Brand Ambien, generic zolpidem, Brand Ambien CR, generic zolpidem ER, Edluar, Brand Intermezzo, generic zolpidem SL tablets, Zolpimist, Belsomra, Dayvigo, estazolam, flurazepam, Brand Halcion, generic triazolam, Brand Lunesta, generic eszopiclone, Quviviq, Brand Restoril, generic temazepam, Brand Rozerem, generic ramelteon, Brand Silenor, generic doxepin, generic zaleplon

Diagnosis	Requests for Patients less than 6 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - The patient is unresponsive to other treatment modalities, unless contraindicated (i.e. other medications or behavioral modification attempted)

**AND**

**2** - The physician attests that the requested medication is medically necessary. (Document rationale for use)

## 2 . Revision History

Date	Notes
3/26/2025	Moved Rozerem from NP section to its own ST criteria box and PA req under 6yo sections.

Serevent Diskus - Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99495
<b>Guideline Name</b>	Serevent Diskus - Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Serevent Diskus	
Diagnosis	Asthma
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of asthma  <b>AND</b>	

2 - Patient is 4 years of age or older

**AND**

3 - Patient is also receiving treatment with an inhaled corticosteroid

Product Name:Serevent Diskus

Diagnosis	Exercise-Induced Bronchospasm
-----------	-------------------------------

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of exercise-induced bronchospasm (EIB)

**AND**

2 - Being used for prevention

**AND**

3 - Patient is 4 years of age or older

Product Name:Serevent Diskus

Diagnosis	Bronchospasm associated with chronic obstructive pulmonary disease (COPD)
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of bronchospasm associated with chronic obstructive pulmonary disease (COPD)

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

SGLT-2 Inhibitors - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-229204
<b>Guideline Name</b>	SGLT-2 Inhibitors - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Farxiga, generic dapagliflozin	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 All of the following: <ul style="list-style-type: none"><li>Patient is 10 years of age or older</li><li>Diagnosis of type 2 diabetes mellitus</li></ul>	

- History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin

**OR**

**1.2** One of the following:

- Diagnosis of chronic kidney disease (CKD)
- Diagnosis of heart failure (NYHA class II-IV) with reduced ejection fraction
- Diagnosis of heart failure (NYHA class II-IV) with preserved ejection fraction

**AND**

**2** - For generic dapagliflozin requests ONLY: History of failure, intolerance, or contraindication to Brand Farxiga

Product Name: Jardiance	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - One of the following:</p> <p><b>1.1</b> All of the following:</p> <ul style="list-style-type: none"> <li>• Patient is 10 years of age or older</li> <li>• Diagnosis of type 2 diabetes mellitus</li> <li>• History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin.</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>1.2</b> Both of the following:</p> <ul style="list-style-type: none"> <li>• Requested medication is being used to reduce the risk of cardiovascular death in adults with type 2 diabetes mellitus and established cardiovascular disease</li> </ul>	

- History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin.

**OR**

**1.3** Requested medication is being used for one of the following:

- To reduce the risk of cardiovascular death and hospitalization for heart failure in adults with heart failure
- To reduce the risk of sustained decline in eGFR, end-stage kidney disease, cardiovascular death, and hospitalization in adults with chronic kidney disease at risk of progression.

Product Name: Synjardy, Synjardy XR	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - All of the following:</p> <ul style="list-style-type: none"> <li>• Patient is 10 years of age or older</li> <li>• Diagnosis of type 2 diabetes mellitus</li> <li>• History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - For Synjardy XR requests ONLY; History of failure, intolerance, or contraindication to ALL of the following:</p> <ul style="list-style-type: none"> <li>• Farxiga</li> <li>• Jardiance</li> </ul>	

Product Name: Inovkana, Invokamet, Invokamet XR, Segluromet, Steglatro, Trijardy XR	
Approval Length	12 month(s)

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Both of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of type 2 diabetes mellitus</li> <li>• History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - History of failure, intolerance, or contraindication to ALL of the following:</p> <ul style="list-style-type: none"> <li>• Farxiga</li> <li>• Jardiance</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is 10 years of age or older (applies to Invokana, Invokamet, Invokamet XR ONLY)</p>	

Product Name: Brand Xigduo XR, generic dapagliflozin-metformin	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 All of the following:</p> <ul style="list-style-type: none"> <li>• Patient is 10 years of age or older</li> <li>• Diagnosis of type 2 diabetes mellitus</li> <li>• History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin.</li> </ul>	

**OR**

**1.2** One of the following:

- Diagnosis of chronic kidney disease (CKD)
- Diagnosis of heart failure (NYHA class II-IV) with reduced ejection fraction

**AND**

**2** - For generic dapagliflozin-metformin requests ONLY: History of failure, intolerance, or contraindication to Brand Xigduo XR

Product Name: Brand Bexagliflozin, Brenzavvy, Glyxambi, Qtern, Steglujan

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Both of the following:

- Diagnosis of type 2 diabetes mellitus
- History of failure to metformin at a minimum dose of 1500mg daily for 90 days, or contraindication or intolerance to metformin

**AND**

**2** - History and failure, intolerance, or contraindication to ALL of the following:

- Janumet or Janumet XR
- Januvia
- Jentadueto or Jentadueto XR
- Kombiglyze XR
- Onglyza
- Tradjenta
- Trijardy XR

**AND**

**3** - History of failure, intolerance, or contraindication to ALL of the following:

- Farxiga
- Jardiance

Product Name: Inpefa

Approval Length | 12 month(s)

Guideline Type | Prior Authorization

**Approval Criteria**

**1** - Requested medication is being used to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with one of the following:

- heart failure
- type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors

**AND**

**2** - History of failure, intolerance, or contraindication to Farxiga

**2 . Revision History**

Date	Notes
4/1/2025	Updated criteria for Synjardy/Synjardy XR

Shingrix (zoster vaccine recombinant, adjuvanted)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116193
<b>Guideline Name</b>	Shingrix (zoster vaccine recombinant, adjuvanted)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

### 1 . Criteria

Product Name:Shingrix*	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Vaccine is being used for prevention of herpes zoster (shingles)  <b>AND</b>  2 - Both of the following:	

**2.1** Patient is between 18 to 49 years of age

**AND**

**2.2** Patient is or will be at increased risk of herpes zoster due to immunodeficiency or immunosuppression caused by known disease or therapy

Notes	* Prior authorization is not required for patients 50 years of age and older.
-------	---

## **2 . Revision History**

Date	Notes
10/28/2022	New program

Short-Acting Opioid Products - AZM



## Prior Authorization Guideline

<b>Guideline ID</b>	GL-165180
<b>Guideline Name</b>	Short-Acting Opioid Products - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/14/2025
-----------------	-----------

### Note:

\*\*PLEASE NOTE: This guideline contains criteria for multiple DUR rejection codes. PA Reviewers: Please confirm the correct criteria is being utilized for reviews based on Diagnosis and Guideline Type. Criteria for MME EXCEEDED rejections can be found towards the end of the guideline.

## 1 . Criteria

Product Name: PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine,

levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4

Diagnosis	PA REQUIRED for use of MAT and other Opioids (Reject 88)
Approval Length	*14 Days for surgical procedure, 5 Days for all other requests
Guideline Type	DUR

**Approval Criteria**

1 - Provider attests to notify the prescriber of the MAT therapy and the prescriber of the MAT therapy approves the concurrent opioid therapy.

**AND**

2 - The days supply does not exceed 14 days for a surgical procedure.

**AND**

3 - The days supply does not exceed 5 days for all other requests.

**AND**

4 - There has not been a previous approval in the last 6 months.

Notes	*Approval Length: 14 Days for surgical procedure, 5 Days for all other requests
-------	---

Product Name:NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-

APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4	
Diagnosis	Non-Preferred Reviews*
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - If the request is for a non-preferred medication the patient must have a history of failure, contraindication or intolerance to a trial of at least FIVE preferred short -acting opioids **. </p> <ul style="list-style-type: none"> <li>• hydromorphone (generic Dilaudid)</li> <li>• meperidine</li> <li>• morphine sulfate</li> <li>• oxycodone IR</li> <li>• tramadol 50mg or 100mg tablets</li> <li>• oxycodone w/ acetaminophen (generic Percocet)</li> <li>• oxycodone-ibuprofen</li> <li>• acetaminophen w/ codeine</li> <li>• butalbital-acetaminophen-caffeine w/ codeine (Generic Fioricet)</li> <li>• butalbital-aspirin-caffeine w/cod (generic Fiorinal)</li> <li>• hydrocodone-acetaminophen</li> <li>• hydrocodone-ibuprofen</li> </ul>	
Notes	*This section does NOT apply to cough and cold products.

<p>Product Name:PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4</p>	
Diagnosis	PA Required for > 2 Short Acting Opioids*

Approval Length	** Authorization will be issued for the requested duration, not to exceed 12 months.
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 The requested medication is being used to adjust the dose of the previously prescribed drug</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 The requested medication will be used in place of the previously prescribed drug, and not in addition to it</p> <p style="text-align: center;"><b>OR</b></p> <p>1.3 The requested medication dosage form will be used in place of the previously prescribed medication dosage form, and not in addition to it</p> <p style="text-align: center;"><b>OR</b></p> <p>1.4 The physician attests they are aware of the multiple short-acting opioids prescribed to the patient and feels treatment with all medications is medically necessary (Document rationale for use)</p>	
Notes	*This section does NOT apply to cough and cold products. ** Authorization will be issued for the requested duration, not to exceed 12 months.

Product Name: PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium,

Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxicodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4

Diagnosis	Quantity Limit*
Approval Length	12 month(s)
Guideline Type	Quantity Limit

**Approval Criteria**

1 - The requested dose cannot be achieved by moving to a higher strength of the product

**AND**

2 - The requested dose is within FDA (Food and Drug Administration) approved maximum dose per day, where an FDA maximum dose per day exists (See table in background section)

Notes	*This section does NOT apply to cough and cold products.
-------	--

Product Name:PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxicodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4

Diagnosis	Greater than 5 day supply requests for patients 18 years of age and older**
-----------	---

Approval Length	**Approvals are for 6 months for all of the above with the exception of post-surgical procedures which can be approved for a 14 day supply. Adults may obtain additional fills without PA if the refill is requested within 60 days from the initial fill.
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following conditions or care instances:</p> <ul style="list-style-type: none"> <li>• Active oncology diagnosis</li> <li>• Hospice care</li> <li>• End-of-life care (other than hospice)</li> <li>• Palliative care</li> <li>• Skilled nursing facility care</li> <li>• Traumatic injury, excluding post-surgical procedures</li> <li>• Chronic conditions for which the provider has received PA approval</li> <li>• Post-surgical procedures</li> </ul>	
Notes	**Approvals are for 6 months for all of the above with the exception of post-surgical procedures which can be approved for a 14 day supply. Adults may obtain additional fills without PA if the refill is requested within 60 days from the initial fill.

<p>Product Name:PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4</p>	
Diagnosis	Greater than 5 day supply requests for patients under 18 years of age**
Guideline Type	Quantity Limit

**Approval Criteria**

1 - ONE of the following conditions or care instances:

- Active oncology diagnosis
- Hospice care
- End-of-life care (other than hospice)
- Palliative care
- Children on opioid wean at time of hospital discharge
- Skilled nursing facility care
- Traumatic injury, excluding post-surgical procedures
- Chronic conditions for which the provider has received PA approval
- Post-surgical procedures

Notes

\*Approvals are for 6 months for all of the above with the exception of post-surgical procedures which can be approved for a 14 day supply. Children and adolescents may obtain additional fills without PA for 5 days supply unless the submitted PA supports a longer duration for use

Product Name: PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4

Diagnosis

Opioid Naïve (Not having filled an opioid in the past 120 days)\*

Guideline Type

Morphine Milligram Equivalents (MME)\*\* MME 50.00 exceeded; PA Required for dosage above 50 MEDD

**Approval Criteria**

**1 - Opioid naïve members may receive greater than 50 morphine milligram equivalent (MME) based on the following:**

**1.1 If the request is for 50 MME to 90 MME, ONE of the following (NOTE: If the request exceeds 90 MME please skip this section and proceed to the Exceeding the 90 MME Cumulative Threshold Reviews section):**

**1.1.1 Diagnosis of ONE of the following:**

- Cancer
- End of life pain (including hospice care)
- Palliative care
- Sickle cell anemia

**OR**

**1.1.2 Patient is currently exceeding 50 MME and prescriber attests patient has been on a short-acting opioid in the past 120 days**

**OR**

**1.1.3 Document ALL of the following:**

- The diagnosis associated with the need for pain management with opioid
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression
- The prescriber has acknowledged that they have completed an addiction risk and risk of overdose assessment
- Prescriber attests the member requires more than 50 MME per day to adequately control pain

Notes

\*This section does NOT apply to cough and cold products. \*\*Approval length for cancer, end of life, palliative care, or sickle cell pain will be issued for 12 months. All other approvals will be issued for one month.

Product Name: PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate

supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4

Diagnosis	Cancer/Hospice/End of Life/ Palliative Care/Skilled Nursing Facility/Traumatic Injury Related Pain Exceeding the 90 MME Cumulative Threshold*
Approval Length	12 Months**
Guideline Type	Morphine Milligram Equivalents (MME) Reviews** (MME 90.00 exceeded; PA REQUIRED; Dosage Above MEDD Limit)
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following conditions:</p> <ul style="list-style-type: none"> <li>• Active oncology diagnosis</li> <li>• Hospice</li> <li>• End-of-life care (other than hospice)</li> <li>• Palliative care</li> <li>• Skilled nursing facility care</li> <li>• Traumatic injury, including burns and excluding post-surgical procedures</li> </ul>	
Notes	*This section does NOT apply to cough and cold products. ** The authorization should be entered for an MME of 9999 so as to prevent future disruptions in therapy if the patient's dose is increased.

Product Name: PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine,

levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral son, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4

Diagnosis	Non-cancer/non-hospice/non-end of life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain Exceeding the 90 MME Cumulative Threshold*
Approval Length	** Authorization will be issued for 6 months for non-cancer/non-hospice/non-end- of-life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain related pain up to the current requested MME plus 90 MME. ***If the member has been established on the requested MME dose for at least 30 days and does not meet the medical necessity authorization criteria requirements, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested MME dose.
Therapy Stage	Initial Authorization
Guideline Type	Morphine Milligram Equivalents (MME) Reviews** (MME 90.00 exceeded; PA REQUIRED; Dosage Above MEDD Limit)

**Approval Criteria**

1 - Prescriber attests to ALL of the following:

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment
- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression

**AND**

2 - BOTH of the following:

- Patient has tried and failed non-opioid pain medication (document drug name and date of trial)

<ul style="list-style-type: none"> <li>• Opioid medication doses of less than 90 morphine milligram equivalent (MME) have been tried and did not adequately control pain (document drug regimen or MME and dates of therapy)***</li> </ul>	
Notes	<p>*This section does NOT apply to cough and cold products. ** Authorization will be issued for 6 months for non-cancer/non-hospice/non-end-of-life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain related pain up to the current requested MME plus 90 MME. ***If the member has been established on the requested MME dose for at least 30 days and does not meet the medical necessity authorization criteria requirements, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested MME dose.</p>

<p>Product Name: PREFERRED: APAP-codeine tabs/soln, Ascomp-codeine, generic butalbital-APAP-ASA-codeine, generic butalbital-ASA-codeine, generic butalbital-generic hydromorphone tablets/liquid, Endocet, Hydrocodone-APAP soln, hydrocodone-APAP tablets, hydrocodone-IBP tablets, hydromorphone tablets/liquid, Brand Hydromorphone supp, Brand Lortab elixir, meperidine tablets, morphine sulfate IR tablets/oral soln, Brand Morphine sulfate supp, Brand Nalocet, Brand Oxaydo, oxycodone IR tablets/capsules/soln, Oxycodone-APAP tablets, Brand Oxycodone-APAP 5mg-325mg soln, Brand Prolate tablets, tramadol 50mg and 100mg tablets; NON-PREFERRED: Brand Apadaz, generic belladonna alkaloids-opium, Brand benzhydrocodone-APAP, butorphanol nasal spray, Brand APAP-caffeine-dihydrocodeine caps, codeine, Brand Dilaudid tablets/liquid, Brand Fioricet-codeine, levorphanol, meperidine oral soln, Nucynta, opium tincture, Brand oxycodone-APAP oral soln, Brand oxycodone abuse-deterrent IR tablets, Brand Oxycodone-APAP 10mg-300mg soln, oxymorphone tablets, generic pentazocine-naloxone, Brand Percocet, Brand Prolate oral soln, Qdolo, Brand Roxycodone, Brand Roxybond, Selgentis, Synapryn, generic tramadol-APAP, Brand tramadol oral soln, Brand Tramadol 25mg and 75mg tablets, Brand Trezix, Brand Tylenol/Codeine #4</p>	
Diagnosis	Non-cancer/non-hospice/non-end of life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain Exceeding the 90 MME Cumulative Threshold*
Approval Length	** Authorization will be issued for 6 months for non-cancer/non-hospice/non-end-of-life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain related pain up to the current requested MME plus 90 MME. *** If the member has been established on the requested MME dose for at least 30 days and does not meet the medical necessity authorization criteria requirements, a denial should be issued and a maximum 30-day authorization may be authorized one time for the requested MME dose.
Therapy Stage	Reauthorization
Guideline Type	Morphine Milligram Equivalents (MME) Reviews** (MME 90.00 exceeded; PA REQUIRED; Dosage Above MEDD Limit)

**Approval Criteria**

1 - Prescriber attests to ALL of the following:

- The information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided
- Treatment goals are defined, including estimated duration of treatment
- Treatment plan includes the use of a non-opioid analgesic and/or non-pharmacologic intervention
- Patient has been screened for substance abuse/opioid dependence
- If used in patients with medical comorbidities or if used concurrently with a benzodiazepine or other drugs that could potentially cause drug-drug interactions, the prescriber has acknowledged that they have completed an assessment of increased risk for respiratory depression

**AND**

2 - Identify rationale for not tapering and discontinuing opioid (Document rationale)

**AND**

3 - Patient demonstrates meaningful improvement in pain and function (Document improvement in function or pain score improvement)\*\*\*

Notes

\*This section does NOT apply to cough and cold products. \*\* Authorization will be issued for 6 months for non-cancer/non-hospice/non-end-of-life/non-palliative care/non-skilled nursing facility/non-traumatic injury related pain related pain up to the current requested MME plus 90 MME. \*\*\* If the member has been established on the requested MME dose for at least 30 days and does not meet the medical necessity authorization criteria requirements, a denial should be issued and a maximum 30 day authorization may be authorized one time for the requested MME dose.

**2 . Background**

**Benefit/Coverage/Program Information**

**CDC Recommended Opioid Maximum Morphine Milligram Equivalents per Day\***

Active Ingredient	FDA Label Max Daily Doses
Morphine	None
Hydromorphone	None
Hydrocodone	None
Tapentadol	600mg IR products
Oxymorphone	None
Oxycodone	None
Codeine	360mg
Pentazocine	None
Tramadol	400mg IR products
Meperidine	600mg
Butorphanol nasal	None
Opium	4 suppositories/day Deodorized tincture: 24mg/day Camphorated tincture: 16mg/day
Acetaminophen	4g/day
Aspirin	2080mg/day
Ibuprofen	3200mg/day
Benzhydrocodone**	None
Levorphanol	None

\*Doses are not considered equianalgesic and table does not represent a dose conversion chart.

\*\*Morphine Milligram Equivalents is derived from the package insert.

**3 . Revision History**

Date	Notes
2/14/2025	NP section: removed Roxicodone as t/f option (oxycodone IR is preferred)

Signifor

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99643
<b>Guideline Name</b>	Signifor
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Signifor	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Both of the following:  1.1 Diagnosis of endogenous Cushing's disease (i.e., hypercortisolism is not a result of chronic administration of high dose glucocorticoids)	

**AND**

**1.2** One of the following:

- Pituitary surgery has not been curative for the patient
- Patient is not a candidate for pituitary surgery

Product Name:Signifor	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Signifor therapy	

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Siliq (brodalumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-329257
<b>Guideline Name</b>	Siliq (brodalumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Siliq	
Diagnosis	Plaque Psoriasis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of chronic moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3 percent body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies:

- corticosteroids (e.g., betamethasone, clobetasol)
- vitamin D analogs (e.g., calcitriol, calcipotriene)
- tazarotene
- calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a dermatologist

Product Name: Siliq	
Diagnosis	Plaque Psoriasis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by ONE of the following:</p> <ul style="list-style-type: none"><li>• Reduction the body surface area (BSA) involvement from baseline</li><li>• Improvement in symptoms (e.g., pruritus, inflammation) from baseline</li></ul> <p><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a dermatologist</p>	

## 2 . Revision History

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, updated criteria. Updated "or" to "and" between 1.1 and 1.2

Simponi, Simponi Aria (golimumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-334189
<b>Guideline Name</b>	Simponi, Simponi Aria (golimumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name: Simponi or Simponi Aria	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting all of the following:  1.1 Diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**1.2** One of the following:

**1.2.1** Patient is receiving concurrent therapy with methotrexate (e.g., Rheumatrex, Trexall)

**OR**

**1.2.2** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab (Janssen manufacturer)
- Orencia (abatacept)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred tocilizumab biosimilar

**AND**

**1.4** Prescribed by or in consultation with a rheumatologist

**AND**

**1.5** For Simponi Aria Requests: Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to self-administered Simponi (APPLIES TO REQUESTS FOR SIMPONI ARIA ONLY)

Product Name: Simponi or Simponi Aria	
Diagnosis	Ankylosing Spondylitis (AS)
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) or verification of paid claims documenting all of the following:</p> <p><b>1.1</b> Diagnosis of active ankylosing spondylitis (AS)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> History of failure to two NSAIDs (non-steroidal anti-inflammatory drugs) (e.g., ibuprofen, naproxen) at maximally indicated doses, each used for at least 4 weeks within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.3</b> History of failure, contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"> <li>• A preferred adalimumab biosimilar or Enbrel (etanercept)</li> <li>• Infliximab (Janssen manufacturer)</li> <li>• Xeljanz (tofacitinib) oral tablet (IR or XR)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.4</b> Prescribed by or in consultation with a rheumatologist</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.5</b> For Simponi Aria Requests: Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to self-administered Simponi (APPLIES TO REQUESTS FOR SIMPONI ARIA ONLY)</p>	

Product Name: Simponi or Simponi Aria	
Diagnosis	Rheumatoid Arthritis (RA), Ankylosing Spondylitis (AS)

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) demonstrating positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p>	

Product Name: Simponi or Simponi Aria	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting all of the following:</p> <p>1.1 Diagnosis of active psoriatic arthritis (PsA)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced</p> <p style="text-align: center;"><b>AND</b></p>	

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab (Janssen manufacturer)
- Orencia (abatacept)
- Otezla (apremilast)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**1.4** Prescribed by or in consultation with one of the following:

- Rheumatologist
- Dermatologist

**AND**

**1.5** For Simponi Aria Requests: Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to self-administered Simponi (APPLIES TO REQUESTS FOR SIMPONI ARIA ONLY)

Product Name: Simponi or Simponi Aria	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) demonstrating positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with one of the following:	

- Rheumatologist
- Dermatologist

Product Name: Simponi	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting all of the following:</p> <p>1.1 Diagnosis of moderately to severely active ulcerative colitis (UC)</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 One of the following:</p> <p>1.2.1 Patient is corticosteroid dependent (i.e., an inability to successfully taper corticosteroids without a return of the symptoms of UC)</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2.2 History of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)</li> <li>• 6-mercaptopurine (Purinethol)</li> <li>• Azathioprine (Imuran)</li> <li>• Aminosalicylates (e.g., mesalamine, sulfasalazine)</li> </ul> <p style="text-align: center;"><b>AND</b></p>	

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- Infliximab (Janssen manufacturer)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**1.4** Prescribed by or in consultation with a gastroenterologist

Product Name: Simponi	
Diagnosis	Ulcerative Colitis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) demonstrating positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with a gastroenterologist	

Product Name: Simponi Aria	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

### Approval Criteria

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of active polyarticular juvenile idiopathic arthritis (PJIA)

**AND**

2 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming a minimum duration of a 6-week trial and failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:

- methotrexate
- leflunomide

**AND**

3 - Submission of medical records (e.g., chart notes) documenting a history of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Orenzia (abatacept)
- Xeljanz (tofacitinib) oral tablet
- A preferred tocilizumab biosimilar

**AND**

4 - Prescribed by or in consultation with a rheumatologist

Product Name: Simponi Aria	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

### **Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) demonstrating positive clinical response to therapy as evidenced by at least one of the following:

- Reduction in the total active (swollen and tender) joint count from baseline
- Improvement in symptoms (e.g., pain, stiffness, inflammation) from baseline

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

## **2 . Revision History**

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, updated criteria throughout. PsA indication: changed tocilizumab to ustekinumab.

Sivextro (tedizolid)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242220
<b>Guideline Name</b>	Sivextro (tedizolid)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Sivextro tablets	
Diagnosis	Skin and Skin Structure Infections
Approval Length	6 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - ONE of the following:  1.1 For continuation of therapy upon hospital discharge	

**OR**

**1.2** As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication.

**OR**

**1.3** ALL of the following:

**1.3.1** Diagnosis of acute bacterial skin and skin structure infection (including diabetic foot infections)

**AND**

**1.3.2** BOTH of the following:

- Patient is at least 26 weeks gestational age
- Patient weighs at least 1 kg

**AND**

**1.3.3** ONE of the following diagnoses:

**1.3.3.1** BOTH of the following:

- Acute bacterial skin and skin structure infections
- Infection caused by methicillin-resistant *Staphylococcus aureus* (MRSA) documented by culture and sensitivity report

**OR**

**1.3.3.2** BOTH of the following:

- Empirical treatment of patients with acute bacterial skin and skin structure infections
- Presence of MRSA infection is likely

**AND**

**1.3.4** History of failure, contraindication, or intolerance to linezolid (generic Zyvox)

**AND**

**1.3.5** History of failure, contraindication, or intolerance to ONE of the following antibiotics:

- Sulfamethoxazole-trimethoprim (SMX-TMP)
- A tetracycline
- Clindamycin

**OR**

**1.4** ALL of the following:

**1.4.1** Diagnosis of acute bacterial skin and skin structure infection(including diabetic foot infections)

**AND**

**1.4.2** BOTH of the following:

- Patient is at least 26 weeks gestational age
- Patient weighs at least 1 kg

**AND**

**1.4.3** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Sivextro

**AND**

**1.4.4** History of failure, contraindication, or intolerance to linezolid (generic Zyvox)

**AND**

**1.4.5** History of failure, contraindication, or intolerance to TWO of the following antibiotics:

- Dicloxacillin
- A cephalosporin
- A tetracycline
- Amoxicillin/clavulanate
- Clindamycin
- Sulfamethoxazole-trimethoprim (SMX-TMP)
- A fluoroquinolone

Product Name: Sivextro tablets	
Diagnosis	Off-Label Uses
Approval Length	60 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 For continuation of therapy upon hospital discharge</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication</p> <p style="text-align: center;"><b>OR</b></p> <p>1.3 BOTH of the following:</p> <p>1.3.1 The medication is being prescribed by or in consultation with an infectious disease specialist</p>	

**AND**

**1.3.2** History of failure, contraindication, or intolerance to linezolid (generic Zyvox), if culture and susceptibility confirm susceptibility.

## **2 . Revision History**

Date	Notes
4/24/2025	Added age/weight criterion where applicable due to expanded indication.

Skyclarys (omaveloxolone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-143858
<b>Guideline Name</b>	Skyclarys (omaveloxolone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Skyclarys	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting BOTH of the following: <ul style="list-style-type: none"><li>Diagnosis of Friedreich's ataxia</li><li>Confirmed presence of a mutation in the frataxin (FXN) gene</li></ul>	

**AND**

2 - Prescribed by or in consultation with one of the following:

- Neurologist
- Neurogeneticist
- Physiatrist (Physical Medicine and Rehabilitation Specialist)

Product Name:Skyclarys	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., slowed disease progression, improvement in or stabilization of speech or swallowing, upper/lower limb coordination, upright stability)</p> <p><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"><li>• Neurologist</li><li>• Neurogeneticist</li><li>• Physiatrist (Physical Medicine and Rehabilitation Specialist)</li></ul>	

**2 . Revision History**

Date	Notes
3/19/2024	Updated criteria to remove mFARS scoring. Added examples of positive response to tx in reauth.

Skyrizi (risankizumab-rzaa)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300298
<b>Guideline Name</b>	Skyrizi (risankizumab-rzaa)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Skyrizi SC 150 mg*	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:  1.1 Diagnosis of moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3 percent body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** Minimum duration of a 4-week trial and failure, contraindication, or intolerance to one of the following topical therapies:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies:

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

**AND**

**2 - Prescribed by or in consultation with a dermatologist**

Notes

\*If patient meets criteria above, please approve at GPI-14\*

Product Name:Skyrizi SC 150 mg\*

Diagnosis | Plaque Psoriasis (PsO)

Approval Length | 12 month(s)

Therapy Stage | Reauthorization

Guideline Type | Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy

**AND**

**2 - Prescribed by or in consultation with a dermatologist**

Notes

\*If patient meets criteria above, please approve at GPI-14\*

Product Name:Skyrizi SC 150 mg\*

Diagnosis | Psoriatic Arthritis (PsA)

Approval Length | 12 month(s)

Therapy Stage | Initial Authorization

Guideline Type | Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

**1.1** Diagnosis of active psoriatic arthritis (PsA)

**AND**

**1.2** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Orencia (abatacept)
- Otezla (apremilast)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Dermatologist
- Rheumatologist

Notes

\*If patient meets criteria above, please approve at GPI-14\*

Product Name:Skyrizi SC 150 mg*	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with one of the following:

- Dermatologist
- Rheumatologist

Notes

\*If patient meets criteria above, please approve at GPI-14\*

Product Name:Skyrizi IV

Diagnosis Crohn's Disease (CD)

Approval Length 3 month(s)

Guideline Type Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

1.1 Diagnosis of moderately to severely active Crohn's disease (CD)

**AND**

1.2 Trial and failure, contraindication, or intolerance to one of the following conventional therapies

- 6-mercaptopurine
- Azathioprine
- Methotrexate
- Corticosteroid (e.g., prednisone)

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- A preferred ustekinumab biosimilar

**AND**

**2** - Will be administered as an intravenous induction dose

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name:Skyrizi SC 180mg, 360 mg*	
Diagnosis	Crohn's Disease (CD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following</p> <p><b>1.1</b> Diagnosis of moderately to severely active Crohn's disease (CD)</p> <p><b>AND</b></p> <p><b>1.2</b> History of failure, contraindication, or intolerance to one of the of the following conventional therapies:</p> <ul style="list-style-type: none"><li>• 6-mercaptopurine</li><li>• Azathioprine</li><li>• Methotrexate</li></ul>	

- Corticosteroid (e.g., prednisone)

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- A preferred ustekinumab biosimilar

**AND**

**2** - Will be used as a maintenance dose following the intravenous induction doses

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Notes

\*If patient meets criteria above, please approve at GPI-14\*

Product Name:Skyrizi IV

Diagnosis	Ulcerative Colitis (UC)
-----------	-------------------------

Approval Length	3 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

**1.1** Diagnosis of moderately to severely active ulcerative colitis (UC)

**AND**

**1.2** One of the following:

- Greater than 6 stools per day
- Frequent blood in the stools
- Frequent urgency
- Presence of ulcers
- Abnormal lab values (e.g., hemoglobin, erythrocyte sedimentation rate, C-reactive protein)
- Dependent on, or refractory to, corticosteroids

**AND**

**1.3** Trial and failure, contraindication, or intolerance to one of the following conventional therapies

- 6-mercaptopurine
- Azathioprine
- Corticosteroid (e.g., prednisone)
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)

**AND**

**1.4** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz oral tablet (tofacitinib) (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Will be administered as an intravenous induction dose

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name:Skyrizi SC 180mg, 360 mg*	
Diagnosis	Ulcerative Colitis (UC)

Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following

**1.1** Diagnosis of moderately to severely active ulcerative colitis (UC)

**AND**

**1.2** History of failure, contraindication, or intolerance to one of the of the following conventional therapies:

- 6-mercaptopurine
- Azathioprine
- Corticosteroid (e.g., prednisone)
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Will be used as a maintenance dose following the intravenous induction doses

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Notes	*If patient meets criteria above, please approve at GPI-14*
-------	---

Product Name:Skyrizi SC 180mg, 360 mg*	
Diagnosis	Crohn's Disease (CD), Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by at least one of the following:</p> <ul style="list-style-type: none"> <li>Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li> <li>Reversal of high fecal output state</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a gastroenterologist</p>	
Notes	*If patient meets criteria above, please approve at GPI-14*

## 2 . Revision History

Date	Notes
7/3/2025	Updated preferred agents/embedded steps, updated criteria throughout.

Skysona (elivaldogene autotemcel suspension)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117546
<b>Guideline Name</b>	Skysona (elivaldogene autotemcel suspension)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Skysona	
Approval Length	1 Time Authorization in Lifetime
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of early, active cerebral adrenoleukodystrophy (CALD)  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) documenting molecular genetic testing confirms mutation in the ABCD1 gene

**AND**

**3** - Submission of medical records (e.g., chart notes) confirming ALL of the following:

- Patient has elevated very long chain fatty acid (VLCFA) levels
- Loes score between 0.5 and 9 (inclusive) based on brain MRI assessment [B, 4]
- Brain magnetic resonance imaging (MRI) utilizes Gadolinium enhancement (GdE +) and demonstrates demyelinating lesions [C, 5]
- Neurologic function score (NFS) less than or equal to 1

**AND**

**4** - BOTH of the following:

- Patient is male sex
- Patient is 4 to 17 years of age

**AND**

**5** - Patient is not eligible for an allogeneic hematopoietic stem cell transplant with an HLA-matched sibling donor

**AND**

**6** - Submission of medical records (e.g., chart notes) confirming patient has obtained a negative test result for all of the following prior to cell collection:

- Hepatitis B virus (HBV)
- Hepatitis C virus (HCV)
- Human T-lymphotropic virus 1 and 2 (HTLV-1/HTLV-2)
- Human immunodeficiency virus (HIV)

**AND**

**7** - Patient does not have CALD secondary to head trauma

**AND**

**8** - Discontinue prophylactic anti-retroviral medications (e.g., Truvada, Descovy) for at least one month prior to initiating medications for stem cell mobilization and until all cycles of apheresis are completed

**AND**

**9** - Prescribed by a stem cell transplant physician from a qualified treatment center

**AND**

**10** - Patient has never received Skysona treatment in their lifetime

## **2 . Revision History**

Date	Notes
11/30/2022	New program

Sodium Oxybate Products (Lumryz, Xyrem, Xywav)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-163938
<b>Guideline Name</b>	Sodium Oxybate Products (Lumryz, Xyrem, Xywav)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Lumryz, Brand Xyrem, Generic sodium oxybate, Xywav	
Diagnosis	Narcolepsy with Cataplexy (i.e., Narcolepsy Type 1)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g. chart notes, laboratory values) documenting a diagnosis of narcolepsy with cataplexy (i.e., Narcolepsy Type 1) with BOTH of the following:	

**1.1** The patient has daily periods of irrepressible need to sleep or daytime lapses into sleep occurring for at least three months

**AND**

**1.2** A mean sleep latency of less than or equal to 8 minutes and two or more sleep onset rapid eye movement (REM) periods (SOREMPs) on a Multiple Sleep Latency Test (MSLT) performed according to standard techniques following a normal overnight polysomnogram. A SOREMP (within 15 minutes of sleep onset) on the preceding nocturnal polysomnogram may replace one of the SOREMPs on the MSLT

**AND**

**2** - Physician attestation to BOTH of the following:

**2.1** Patient has experienced cataplexy defined as more than one episode of sudden loss of muscle tone with retained consciousness

**AND**

**2.2** Other causes of sleepiness have been ruled out or treated (including but not limited to obstructive sleep apnea, insufficient sleep syndrome, shift work, the effects of substances or medications, or other sleep disorders)

**AND**

**3** - Prescribed by ONE of the following:

- Neurologist
- Psychiatrist
- Sleep Medicine Specialist

**AND**

**4** - Patient is 7 years of age or older

Product Name:Lumryz, Brand Xyrem, Generic sodium oxybate, Xywav	
Diagnosis	Narcolepsy with Cataplexy (i.e., Narcolepsy Type 1)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting a reduction in frequency of cataplexy attacks associated with therapy</p> <p style="text-align: center;"><b>OR</b></p> <p>2 - Documentation demonstrating reduction in symptoms of excessive daytime sleepiness associated with therapy</p>	

Product Name:Lumryz, Brand Xyrem, Generic sodium oxybate, Xywav	
Diagnosis	Narcolepsy without Cataplexy (i.e., Narcolepsy Type 2)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g. chart notes, lab values) documenting a diagnosis of narcolepsy without cataplexy (i.e., Narcolepsy Type 2) with BOTH of the following:</p> <p>1.1 The patient has daily periods of irrepressible need to sleep or daytime lapses into sleep occurring for at least three months</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 A mean sleep latency of less than or equal to 8 minutes and two or more sleep onset rapid eye movement (REM) periods (SOREMPs) are found on a Multiple Sleep Latency Test</p>	

(MSLT) performed according to standard techniques following a normal overnight polysomnogram. A SOREMP (within 15 minutes of sleep onset) on the preceding nocturnal polysomnogram may replace one of the SOREMPs on the MSLT

**AND**

**2** - Physician attestation to BOTH of the following:

**2.1** Cataplexy is absent

**AND**

**2.2** Other causes of sleepiness have been ruled out or treated (including but not limited to obstructive sleep apnea, insufficient sleep syndrome, shift work, the effects of substances or medications or their withdrawal, sleep phase disorder, or other sleep disorders)

**AND**

**3** - History of failure, contraindication, or intolerance of ALL of the following (MUST be verified via paid pharmacy claims or submission of medical records):

**3.1** ONE of the following:

- Amphetamine based stimulant (e.g., amphetamine, dextroamphetamine)
- Methylphenidate based stimulant

**AND**

**3.2** Armodafanil (Nuvigil)

**AND**

**3.3** Sunosi (solriamfetol)

**AND**

**4** - Prescribed by ONE of the following:

- Neurologist
- Psychiatrist
- Sleep Medicine Specialist

**AND**

**5** - Patient is 7 years of age or older

Product Name:Lumryz, Brand Xyrem, Generic sodium oxybate, Xywav	
Diagnosis	Narcolepsy without Cataplexy (i.e., Narcolepsy Type 2)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting reduction in symptoms of excessive daytime sleepiness associated with therapy</p>	

Product Name:Xywav	
Diagnosis	Idiopathic Hypersomnia (IH)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of idiopathic hypersomnia (IH) confirmed by ALL of the following:</p> <p><b>1.1</b> Patient has experienced daily periods of irrepresible need for sleep or daytime lapses into sleep (i.e., excessive daytime sleepiness) for at least 3 months</p>	

**AND**

**1.2** A multiple sleep latency test (MSLT) documents fewer than two sleep-onset rapid eye movement periods (SOREMPs), or no SOREMPs if the REM sleep latency on the preceding polysomnogram was  $\leq 15$  minutes

**AND**

**1.3** The presence of at least one of the following:

- MSLT shows a mean sleep latency of  $\leq 8$  minutes
- Total 24-hour sleep time is  $\geq 660$  minutes (typically 12 to 14 hours) on 24-hour polysomnography or by wrist actigraphy in association with a sleep log

**AND**

**2** - Physician attestation to BOTH of the following:

**2.1** Cataplexy is absent

**AND**

**2.2** Other causes of sleepiness have been ruled out or treated (including but not limited to obstructive sleep apnea, insufficient sleep syndrome, shift work, the effects of substances or medications or their withdrawal, sleep phase disorder, or other sleep disorders)

**AND**

**3** - Prescribed by with one of the following:

- Neurologist
- Psychiatrist
- Sleep Medicine Specialist

**AND**

**4** - History of failure, contraindication, or intolerance of ALL of the following (MUST be verified via paid pharmacy claims or submission of medical records):

- An amphetamine or methylphenidate based stimulant
- modafinil
- armodafinil

Product Name: Xywav	
Diagnosis	Idiopathic Hypersomnia (IH)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting reduction in symptoms of excessive daytime sleepiness associated with therapy</p>	

## 2 . Revision History

Date	Notes
1/31/2025	IH: Updated prescriber requirement verbiage

Sofdra (sofpironium)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-228336
<b>Guideline Name</b>	Sofdra (sofpironium)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Sofdra	
Diagnosis	Primary Axillary Hyperhidrosis
Approval Length	6 Week(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of primary axillary hyperhidrosis	

**AND**

**2** - Patient has experienced symptoms of disease for a duration of at least 6 months

**AND**

**3** - Patient is 9 years of age or older

**AND**

**4** - Provider attests other causes of axillary hyperhidrosis have been ruled out (e.g., menopause, medications)

**AND**

**5** - Disease frequently interferes with daily activities (e.g., lower productivity at work, lower psychosocial wellbeing)

**AND**

**6** - Trial and failure (minimum 6 weeks supply), contraindication, or intolerance to a topical prescription strength drying agent [e.g., Drysol, Xerac AC (aluminum chloride hexahydrate)]

**AND**

**7** - Trial and failure, contraindication, or intolerance to both of the following:

- Botox
- Qbrexza

**AND**

**8** - Prescribed by or in consultation with a dermatologist

Product Name:Sofdra	
Diagnosis	Primary Axillary Hyperhidrosis
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy (e.g., improvement in productivity, improvement in psychosocial wellbeing)</p>	

## 2 . Revision History

Date	Notes
4/24/2025	Added step through Botox and Qbreza

Sohonos (palovarotene)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-136960
<b>Guideline Name</b>	Sohonos (palovarotene)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Sohonos	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting ALL of the following: 1.1 Diagnosis of Fibrodysplasia Ossificans Progressiva (FOP)	

**AND**

**1.2** Molecular genetic testing confirms mutation in the ACVR1 gene

**AND**

**1.3** One of the following:

**1.3.1** Both of the following:

- Patient is female
- Patient is 8 years of age or older

**OR**

**1.3.2** Both of the following:

- Patient is male
- Patient is 10 years of age or older

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Geneticist
- Orthopedic physician
- Rheumatologist
- Endocrinologist

Product Name: Sohonos	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) documenting that patient demonstrates positive clinical response to therapy (e.g., reduction of volume in new abnormal bone growth)

**2 . Revision History**

Date	Notes
12/1/2023	New program

Somatuline Depot (lanreotide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157634
<b>Guideline Name</b>	Somatuline Depot (lanreotide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Brand Somatuline Depot, generic lanreotide	
Diagnosis	Acromegaly
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of acromegaly	

**AND**

**2** - One of the following:

**2.1** Inadequate response to one of the following:

- Surgery
- Radiotherapy

**OR**

**2.2** Not a candidate for one of the following:

- Surgery
- Radiotherapy

**AND**

**3** - Prescribed by or in consultation with an endocrinologist

**AND**

**4** - Trial and failure, or intolerance to generic octreotide

Product Name: Brand Somatuline Depot, generic lanreotide	
Diagnosis	Acromegaly
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive	

clinical response to therapy, such as a reduction or normalization of IGF-1/GH level for same age and sex

Product Name:Brand Somatuline Depot 120mg/0.5mL, generic lanreotide 120mg/0.5ml	
Diagnosis	Advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NET)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of gastroenteropancreatic neuroendocrine tumor (GEP-NET)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Disease is one of the following:</p> <ul style="list-style-type: none"><li>• Unresectable, locally advanced</li><li>• Metastatic</li></ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with an oncologist</p>	

Product Name:Brand Somatuline Depot 120mg/0.5mL, generic lanreotide 120mg/0.5ml	
Diagnosis	Advanced or metastatic gastroenteropancreatic neuroendocrine tumors (GEP-NET)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting patient does not show evidence of progressive disease while on therapy

Product Name:Brand Somatuline Depot 120mg/0.5mL, generic lanreotide 120mg/0.5ml

Diagnosis	Carcinoid Syndrome
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of carcinoid syndrome

**AND**

2 - Used to reduce the frequency of short-acting somatostatin analog rescue therapy

**AND**

3 - Prescribed by or in consultation with one of the following:

- Endocrinologist
- Oncologist

**AND**

4 - Trial and failure, or intolerance to generic octreotide

Product Name:Brand Somatuline Depot 120mg/0.5mL, generic lanreotide 120mg/0.5ml

Diagnosis	Carcinoid Syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
10/25/2024	Updated naming conventions of targets, removed off-label designation for lanreotide/carcinoid syndrome

Somavert

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99644
<b>Guideline Name</b>	Somavert
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Somavert	
Diagnosis	Acromegaly
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - All of the following:  1.1 Diagnosis of acromegaly by ONE of the following:	

- Serum GH (growth hormone) level greater than 1 ng/mL (nanograms per milliliter) after a 2 hour oral glucose tolerance test (OGTT) at time of diagnosis
- Elevated serum IGF-1 (Insulin-like growth factor-1) levels (above the age and gender adjusted normal range as provided by the physician's lab) at time of diagnosis

**AND**

**1.2** One of the following:

**1.2.1** Inadequate response to one of the following:

- Surgery
- Radiation therapy
- Dopamine agonist (e.g., bromocriptine, cabergoline) therapy

**OR**

**1.2.2** Not a candidate for all of the following:

- Surgery
- Radiation therapy
- Dopamine agonist (e.g., bromocriptine, cabergoline) therapy

**AND**

**1.3** Inadequate response, intolerance, or contraindication to one of the following somatostatin analogs:

- Sandostatin (octreotide) or Sandostatin LAR
- Somatuline Depot (lanreotide)

**OR**

**2** - Patient is currently on Somavert therapy for acromegaly

Product Name: Somavert	
Diagnosis	Acromegaly

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Somavert therapy</p>	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Soriatane

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99496
<b>Guideline Name</b>	Soriatane
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Brand Soriatane, Generic acitretin	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of severe psoriasis  <b>AND</b>	

**2** - Prescribed or recommended by a dermatologist

**AND**

**3** - One of the following:

**3.1** Patient is unresponsive to other therapies (e.g., topical corticosteroids, topical vitamin D analogs, tazarotene, methotrexate)

**OR**

**3.2** Other therapies are contraindicated based on the patient's clinical condition

**AND**

**4** - One of the following:

- Greater than or equal to 10% body surface area involvement
- Palmoplantar, facial, or genital involvement
- Severe scalp psoriasis

Product Name: Brand Soriatane, Generic acitretin	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Soriatane therapy	
<b>AND</b>	
2 - Prescribed or recommended by a dermatologist	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Sotyktu (deucravacitinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300299
<b>Guideline Name</b>	Sotyktu (deucravacitinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Sotyktu	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims confirming ALL of the following:  1.1 Diagnosis of moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3% body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** History of failure to one of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- corticosteroids (e.g., betamethasone, clobetasol)
- vitamin D analogs (e.g., calcitriol, calcipotriene)
- tazarotene
- calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure of a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies:

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a dermatologist

**AND**

**3** - Not used in combination with other potent immunosuppressants (e.g., azathioprine, cyclosporine)

Product Name:Sotyktu	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming positive clinical response to therapy as evidenced by ONE of the following:</p> <ul style="list-style-type: none"><li>• Reduction the body surface area (BSA) involvement from baseline</li><li>• Improvement in symptoms (e.g., pruritus, inflammation) from baseline</li></ul> <p><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation with a dermatologist</p> <p><b>AND</b></p> <p><b>3</b> - Not used in combination with other potent immunosuppressants (e.g., azathioprine, cyclosporine)</p>	

## 2 . Revision History

Date	Notes
7/3/2025	Updated preferred agents/embedded steps, updated criteria.

Spevigo (spesolimab-sbzo)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147674
<b>Guideline Name</b>	Spevigo (spesolimab-sbzo)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Spevigo IV	
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of generalized pustular psoriasis (GPP)  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) confirming patient has a moderate to severe GPP flare based on one of the following:

- Presence of fresh pustules (new appearance or worsening of pustules)
- At least 5% of body surface area (BSA) covered with erythema and the presence of pustules
- A Generalized Pustular Psoriasis Physician Global Assessment (GPPPGA) total score of at least 3 (moderate)
- GPPPGA pustulation sub score of at least 2 (mild)

**AND**

**3** - Both of the following:

- Patient is 12 years of age or older
- Patient weighs at least 40 kg

**AND**

**4** - Prescribed by or in consultation with a dermatologist

**AND**

**5** - Patient has not already received two infusions of Spevigo for a single flare

Product Name:Spevigo SC	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming diagnosis of generalized pustular psoriasis (GPP) as defined by both of the following:	

- Primary, sterile, macroscopically visible pustules on non-acral skin (excluding cases where pustulation is restricted to psoriatic plaques)
- Disease is relapsing (> 1 episode) or persistent (> 3 months)

**AND**

**2** - Subcutaneous formulation will not be used to treat GPP flare

**AND**

**3** - Both of the following:

- Patient is 12 years of age or older
- Patient weighs at least 40 kg

**AND**

**4** - Prescribed by or in consultation with a dermatologist

Product Name:Spevigo SC	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., reduction in number of flares)</p>	

## 2 . Revision History

Date	Notes
------	-------

5/23/2024	Updated IV formulation criteria due to expanded indication and added criteria for new SC formulation.
-----------	---

Spinraza- Arizona

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99729
<b>Guideline Name</b>	Spinraza- Arizona
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Spinraza	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of spinal muscular atrophy (SMA) type I, II, or III made by, or in consultation with, a neurologist with expertise in the diagnosis of SMA	

**AND**

**2** - Submission of medical records (e.g., chart notes, laboratory values) confirming both of the following:

**2.1** The mutation or deletion of genes in chromosome 5q resulting in one of the following:

- Homozygous gene deletion or mutation (e.g., homozygous deletion of exon 7 at locus 5q13)
- Compound heterozygous mutation (e.g., deletion of SMN1 exon 7 [allele 1] and mutation of SMN1 [allele 2])

**AND**

**2.2** Patient has at least 2 copies of SMN2

**AND**

**3** - Patient is not dependent on invasive ventilation or tracheostomy

**AND**

**4** - Patient is not dependent on use of non-invasive ventilation beyond use for naps and nighttime sleep

**AND**

**5** - Submission of medical records (e.g., chart notes, laboratory values) or claims history of the baseline exam of one of the following exams (based on patient age and motor ability) to establish baseline motor ability:

- Hammersmith Infant Neurological Exam Part 2 (HINE-2) (infant to early childhood)
- Hammersmith Functional Motor Scale Expanded (HFMSE)
- Upper Limb Module (ULM) Test (Non ambulatory)
- Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND)

**AND**

**6** - Prescribed by, or in consultation with, a neurologist with expertise in the treatment of SMA

**AND**

**7** - One of the following:

**7.1** Patient has not previously received gene replacement therapy for the treatment of SMA

**OR**

**7.2** One of the following:

**7.2.1** Both of the following:

**7.2.1.1** Patient recently received gene replacement therapy within the previous 6 months

**AND**

**7.2.1.2** Patient has experienced a declination in clinical status since receipt of gene replacement therapy

**OR**

**7.2.2** Both of the following:

**7.2.2.1** Patient has previously received gene replacement therapy

**AND**

**7.2.2.2** Patient has experienced a declination in clinical status that represents a potential abatement of gene therapy efficacy

**AND**

**8** - Spinraza is to be administered intrathecally by, or under the direction of, healthcare professionals experienced in performing lumbar punctures

**AND**

**9** - Spinraza dosing for SMA is within accordance with the United States Food and Drug Administration approved labeling: maximum dosing of 12 milligrams for each loading dose

Product Name: Spinraza	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of spinal muscular atrophy (SMA) type I, II, or III made by, or in consultation with, a neurologist with expertise in the diagnosis of SMA</p> <p><b>AND</b></p> <p><b>2</b> - Submission of medical records (e.g., chart notes, laboratory values) or claims history confirming both of the following:</p> <p><b>2.1</b> The mutation or deletion of genes in chromosome 5q resulting in one of the following:</p> <ul style="list-style-type: none"><li>• Homozygous gene deletion or mutation (e.g., homozygous deletion of exon 7 at locus 5q13)</li><li>• Compound heterozygous mutation (e.g., deletion of SMN1 exon 7 [allele 1] and mutation of SMN1 [allele 2])</li></ul> <p><b>AND</b></p> <p><b>2.2</b> Patient has at least 2 copies of SMN2</p>	

**AND**

**3** - Patient is not dependent on invasive ventilation or tracheostomy

**AND**

**4** - Patient is not dependent on use of non-invasive ventilation beyond use for naps and nighttime sleep

**AND**

**5** - One of the following:

**5.1** Patient has not previously received gene replacement therapy for the treatment of SMA

**OR**

**5.2** Both of the following:

**5.2.1** Patient has previously received gene replacement therapy

**AND**

**5.2.2** Patient has experienced a declination in clinical status that represented a potential failure or abatement of gene therapy efficacy

**AND**

**6** - Submission of medical records (e.g., chart notes, laboratory values) or claims history with the most recent results (less than 1 month prior to request) documenting a positive clinical response from pretreatment baseline status to Spinraza therapy as demonstrated by one of the following exams:

**6.1** Both of the following for Hammersmith Infant Neurological Exam Part 2 (HINE-2) milestones:

**6.1.1** One of the following:

- Improvement or maintenance of previous improvement of at least 2 point (or maximal score) increase in ability to kick
- Improvement or maintenance of previous improvement of at least 1 point increase in any other HINE-2 milestone (e.g., head control, rolling, sitting, crawling, etc.), excluding voluntary grasp

**AND**

**6.1.2** One of the following:

- The patient exhibited improvement or maintenance of previous improvement in more HINE motor milestones than worsening, from pretreatment baseline (net positive improvement)
- Achieved and maintained any new motor milestones when they would otherwise be unexpected to do so (e.g., sit unassisted, stand, walk)

**OR**

**6.2** One of the following for Hammersmith Functional Motor Scale Expanded (HFMSSE):

- Improvement or maintenance of previous improvement of at least a 3 point increase in score from pretreatment baseline
- Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**OR**

**6.3** One of the following for Upper Limb Module (ULM):

- Improvement or maintenance of previous improvement of at least a 2 point increase in score from pretreatment baseline
- Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**OR**

**6.4** One of the following for Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders (CHOP INTEND):

**6.4.1** Improvement or maintenance of previous improvement of at least a 4 point increase in score from pretreatment baseline

**OR**

**6.4.2** Patient has achieved and maintained any new motor milestone from pretreatment baseline when they would otherwise be unexpected to do so

**OR**

**6.4.3** Both of the following:

- Patient was prescribed Spinraza due to clinical declination after receipt of gene therapy
- Patients clinical status has stabilized after receipt of Spinraza therapy

**AND**

**7** - Prescribed by, or in consultation with, a neurologist with expertise in the treatment of SMA

**AND**

**8** - Spinraza is to be administered intrathecally by, or under the direction of, healthcare professionals experienced in performing lumbar punctures

**AND**

**9** - Spinraza dosing for SMA is within accordance with the United States Food and Drug Administration approved labeling: maximum dosing of 12 milligrams every 4 months, starting 4 months after the last loading dose

## 2 . Revision History

Date	Notes
------	-------

5/25/2021	7/1 Implementation
-----------	--------------------

Spiriva (generic tiotropium) products

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144742
<b>Guideline Name</b>	Spiriva (generic tiotropium) products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/21/2024
-----------------	-----------

## 1 . Criteria

Product Name:generic tiotropium bromide	
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Requests for generic tiotropium bromide (generic Spiriva Handihaler) should be denied. The plan's preferred products are Brand Spiriva Handihaler and Spiriva Respimat	
Notes	Note: Clinical Program: Brand Over Generic-Not Covered

## 2 . Revision History

Date	Notes
3/21/2024	Update guideline to add note that calls out brand is preferred

Spravato, ketamine - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164878
<b>Guideline Name</b>	Spravato, ketamine - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Spravato, ketamine	
Diagnosis	Major Depressive Disorder (Treatment-Resistant)
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has a confirmed diagnosis of major depressive disorder as defined by the DSM-V (Diagnostic and Statistical Manual of Mental Disorders) criteria and is treatment resistant	

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Requested medication is prescribed by, or in consultation with, a psychiatric provider

**AND**

**4** - ONE of the following:

**4.1** Patient does not have an active substance use disorder (SUD)

**OR**

**4.2** BOTH of the following:

- Patient has an active substance use disorder
- Patient is currently receiving treatment

**AND**

**5** - ONE of the following:

**5.1** Patient has experienced an inadequate response during the current depressive episode with BOTH of the following therapies:

**5.1.1** Two antidepressants from at least two different classes [must include one of each AHCCCS (Arizona Health Care Cost Containment System) preferred agents: SSRI (selective serotonin reuptake inhibitor), SNRI (serotonin-norepinephrine reuptake inhibitor), or bupropion] having different mechanisms of action at the maximally tolerated labeled dose, each used for at least 4-6 weeks

**AND**

**5.1.2** At least TWO augmentation therapies below for at least 4 weeks:

- SSRI or SNRI, and a second-generation antipsychotic used concomitantly (aripiprazole, quetiapine, risperidone, olanzapine)
- SSRI or SNRI, and lithium used concomitantly
- SSRI or SNRI, and liothyronine (T3) used concomitantly
- SSRI or SNRI, and mirtazapine
- SSRI and bupropion and buspirone

**OR**

**5.2** Patient has active suicidal ideation and urgent symptom control is necessary

**AND**

**6** - Requested medication is used in combination with an oral antidepressant (e.g., duloxetine, escitalopram, sertraline, venlafaxine)

**AND**

**7** - Requested medication is administered under the direct supervision of a healthcare provider

**AND**

**8** - Provider is certified in the Spravato REMS (risk evaluation and mitigation strategy) program (Applies to Spravato requests ONLY)

**AND**

**9** - Patient must be monitored by a health care provider for at least 2 hours after administration

Product Name: Spravato, ketamine

Diagnosis

Major Depressive Disorder (Treatment-Resistant)

Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Provider attests that the patient has documented improvement or sustained improvement in depressive symptoms from baseline</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient use of requested medication is in combination with an oral antidepressant</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient administers requested medication under the direct supervision of a healthcare provider</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Provider is certified in the Spravato REMS (risk evaluation and mitigation strategy) program (applies to Spravato requests ONLY)</p> <p style="text-align: center;"><b>AND</b></p> <p>5 - Patient must continue to be monitored by a health care provider for at least 2 hours after administration</p>	

Product Name: Spravato, ketamine	
Diagnosis	Requests for Patients less than 6 years of age
Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - The patient is unresponsive to other treatment modalities, unless contraindicated (i.e. other medications or behavioral modification attempted)

**AND**

2 - The physician attests that the requested medication is medically necessary. (Document rationale for use)

Product Name: Spravato, ketamine	
Diagnosis	Depressive symptoms in an adult with major depressive disorder (MDD) with acute suicidal ideation or behavior
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of major depressive disorder according to the current Diagnostic and Statistical Manual of Mental Disorders (DSM) (i.e., DSM-5) criteria</p> <p><b>AND</b></p> <p>2 - Patient is experiencing an acute suicidal ideation or behavior</p> <p><b>AND</b></p> <p>3 - Patient is receiving newly initiated or optimized oral antidepressant</p> <p><b>AND</b></p>	

4 - Provider and/or the provider's healthcare setting is certified in the Spravato REMS (Risk Evaluation and Mitigation Strategy) program (applies to Spravato requests ONLY)

## 2 . Background

Benefit/Coverage/Program Information		
HCPCS Codes		
CODE	DESCRIPTION	LAY DESCRIPTION
96360	Intravenous infusion, hydration; initial, 31 minutes to 1 hour	A physician or an assistant under direct physician supervision infuses a hydration solution (prepackaged fluid and electrolytes) for 31 minutes to one hour through an intravenous catheter inserted by needle into a patient's vein or by infusion through an existing indwelling intravascular access catheter or port. Report 96361 for each additional hour beyond the first hour. Intravenous infusion for hydration lasting 30 minutes or less is not reported.
96361	Intravenous infusion, hydration; each additional hour (List separately in addition to code for primary procedure)	See 96360
96365	Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); initial, up to 1 hour	A physician or an assistant under direct physician supervision injects or infuses a therapeutic, prophylactic (preventive), or diagnostic medication other than chemotherapy or other highly complex drugs or biologic agents via intravenous route. Infusions are administered through

		<p>an intravenous catheter inserted by needle into a patient's vein or by injection or infusion through an existing indwelling intravascular access catheter or port.</p> <p>Report 96365 for the initial hour and 96366 for each additional hour. Report 96367 for each additional sequential infusion of a different substance or drug, up to one hour, and 96368 for each concurrent infusion of substances other than chemotherapy or other highly complex drugs or biologic agents.</p>	
96366	<p>Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); each additional hour (List separately in addition to code for primary procedure)</p>	See 96365	
96367	<p>Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); additional sequential infusion of a new drug/substance, up to 1 hour (List separately in addition to code for primary procedure)</p>	See 96365	
96368	<p>Intravenous infusion, for therapy, prophylaxis, or diagnosis (specify substance or drug); concurrent infusion (List separately in addition to code for primary procedure)</p>	<p>A physician or an assistant under direct physician supervision injects or infuses a therapeutic, prophylactic (preventive), or diagnostic medication other than chemotherapy or other highly complex drugs or biologic agents via intravenous route. Infusions are administered through an intravenous catheter inserted by</p>	

		<p>needle into a patient's vein or by injection or infusion through an existing indwelling intravascular access catheter or port.</p> <p>Report 96365 for the initial hour and 96366 for each additional hour. Report 96367 for each additional sequential infusion of a different substance or drug, up to one hour, and 96368 for each concurrent infusion of substances other than chemotherapy or other highly complex drugs or biologic agents.</p>
96374	<p>Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); intravenous push, single or initial substance/drug</p>	<p>The physician or an assistant under direct physician supervision administers a therapeutic, prophylactic, or diagnostic substance by subcutaneous or intramuscular injection (96372), intra-arterial injection (96373), or by push into an intravenous catheter or intravascular access device (96374 for a single or initial substance, 96375 for each additional sequential IV push of a new substance, and 96376 for each additional sequential IV push of the same substance after 30 minutes have elapsed). The push technique involves an infusion of less than 15 minutes. Code 96376 may be reported only by facilities.</p>
96375	<p>Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); each additional sequential intravenous push of a new substance/drug (List separately in addition to</p>	<p>See 96374</p>

	code for primary procedure)		
96376	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); each additional sequential intravenous push of the same substance/drug provided in a facility (List separately in addition to code for primary procedure)	See 96374 Code 96376 may be reported only by facilities.	
96379	Unlisted therapeutic, prophylactic, or diagnostic intravenous or intra-arterial injection or infusion		

### 3 . Revision History

Date	Notes
2/25/2025	Added new GPI for ketamine injection

Strensiq

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99646
<b>Guideline Name</b>	Strensiq
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Strensiq	
Diagnosis	perinatal/infantile or juvenile-onset hypophosphatasia (HPP)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - All of the following:  1.1 Diagnosis of perinatal/infantile or juvenile-onset hypophosphatasia based on all of the following:	

**1.1.1** One of the following:

- Onset of clinical signs and symptoms of hypophosphatasia prior to age 18 years (e.g., respiratory insufficiency, vitamin B6 responsive seizures, hypotonia, failure to thrive, delayed walking, waddling gait, dental abnormalities, low trauma fractures)
- Radiographic evidence supporting the diagnosis of hypophosphatasia at the age of onset prior to age 18 years (e.g., craniosynostosis, infantile rickets, non-traumatic fractures)

**AND**

**1.1.2** One of the following:

**1.1.2.1** Both of the following:

- Patient has low level activity of serum alkaline phosphatase (ALP) evidenced by an ALP level below the age-adjusted normal range
- Patient has an elevated level of tissue non-specific alkaline phosphatase (TNSALP) substrate (e.g. serum pyridoxal 5'-phosphate [PLP] level, serum or urine phosphoethanolamine [PEA] level, urinary inorganic pyrophosphate [PPi level])

**OR**

**1.1.2.2** Confirmation of tissue-nonspecific alkaline phosphatase (TNSALP) gene mutation by ALPL genomic DNA testing\*

**AND**

**1.2** Prescribed by one of the following:

- Endocrinologist
- A specialist experienced in the treatment of metabolic bone disorders

**AND**

**1.3** One of the following:

**1.3.1** Both of the following:

- Diagnosis of perinatal/infantile-onset hypophosphatasia

- Coverage will be provided up to a maximum supply limit of 9 mg/kg/week

**OR**

**1.3.2** Both of the following:

- Diagnosis of juvenile-onset hypophosphatasia
- Coverage will be provided up to a maximum supply limit of 6 mg/kg/week

**AND**

**1.4** One of the following:

**1.4.1** Patient is prescribed Strensiq 18 mg/0.45 mL, Strensiq 28 mg/0.7 mL, or Strensiq 40 mg/mL vials

**OR**

**1.4.2** Both of the following:

- Patient is prescribed Strensiq 80 mg/0.8 mL vial
- Patient's weight is greater than or equal to 40 kg

**AND**

**1.5** Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

Notes	*Results of prior genetic testing can be submitted as confirmation of diagnosis of HPP, however please note that the provider should confirm coverage status of any new genetic testing under the patient's United Healthcare plan prior to ordering
-------	--

Product Name:Strensiq	
Diagnosis	perinatal/infantile or juvenile-onset hypophosphatasia (HPP)
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - All of the following:</p> <p><b>1.1</b> Clinically relevant decrease from baseline in tissue non-specific alkaline phosphatase (TNSALP) substrate (e.g. serum pyridoxal 5'-phosphate [PLP] level, serum or urine phosphoethanolamine [PEA] level, urinary inorganic pyrophosphate [PPi level])</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> Prescribed by one of the following:</p> <ul style="list-style-type: none"> <li>• Endocrinologist</li> <li>• A specialist experienced in the treatment of metabolic bone diseases</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.3</b> One of the following:</p> <p><b>1.3.1</b> Both of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of perinatal/infantile-onset hypophosphatasia</li> <li>• Coverage will be provided up to a maximum supply limit of 9 mg/kg/week</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p><b>1.3.2</b> Both of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of juvenile-onset hypophosphatasia</li> <li>• Coverage will be provided up to a maximum supply limit of 6 mg/kg/week</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.4</b> One of the following:</p>	

**1.4.1** Patient is prescribed Strensiq 18 mg/0.45 mL, Strensiq 28 mg/0.7 mL, or Strensiq 40 mg/mL vials

**OR**

**1.4.2** Both of the following

- Patient is prescribed Strensiq 80 mg/0.8 mL vials
- Patient's weight is greater than or equal to 40 kg

**AND**

**1.5** Prescriber attests to the following: the information provided is true and accurate to the best of their knowledge and they understand that UnitedHealthcare may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Sublingual Immunotherapy (SLIT)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-221194
<b>Guideline Name</b>	Sublingual Immunotherapy (SLIT)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2025
-----------------	----------

## 1 . Criteria

Product Name:All products	
Diagnosis	Patients 21 years of age and older
Approval Length	N/A - All requests for patients 21 years of age and older should be DENIED as benefit exclusion
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Requests for patients 21 years of age and older are not covered	
Notes	Approval Length: N/A - All requests for patients 21 years of age and older should be denied as a benefit exclusion.

Product Name:Grastek	
Diagnosis	Grass pollen-induced allergic rhinitis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of moderate to severe grass pollen-induced allergic rhinitis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Diagnosis confirmed by one of the following:</p> <ul style="list-style-type: none"> <li>• Positive skin test to Timothy grass or cross-reactive grass pollens (eg, Sweet Vernal, Orchard/Cocksfoot, Perennial Rye, Kentucky blue/June grass, Meadow Fescue, or Redtop)</li> <li>• in vitro testing for pollen-specific IgE antibodies for Timothy grass or cross-reactive grass pollens (e.g., Sweet Vernal, Orchard/Cocksfoot, Perennial Rye, Kentucky blue/June grass, Meadow Fescue, or Redtop)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Treatment is started or will be started at least 12 weeks before the beginning of the grass pollen season</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - History of failure, contraindication, or intolerance to two of the following:</p> <ul style="list-style-type: none"> <li>• oral antihistamine [e.g. cetirizine (Zyrtec)]</li> <li>• intranasal antihistamine [e.g. azelastine (Astelin)]</li> <li>• intranasal corticosteroid [e.g. fluticasone (Flonase)]</li> <li>• leukotriene inhibitor [e.g. montelukast (Singulair)]</li> </ul> <p style="text-align: center;"><b>AND</b></p>	

**5** - Not received in combination with similar cross-reactive grass pollen immunotherapy (e.g., Oralair)

**AND**

**6** - Patient does not have unstable and/or uncontrolled asthma

**AND**

**7** - Prescribed by or in consultation with a specialist in allergy and immunology

**Product Name:Grastek**

Diagnosis	Grass pollen-induced allergic rhinitis
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Documentation of positive clinical response to Grastek therapy

**Product Name:Oralair**

Diagnosis	Grass pollen-induced allergic rhinitis
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of moderate to severe grass pollen-induced allergic rhinitis

**AND**

**2** - Diagnosis confirmed by one of the following:

- Positive skin test to any of the five grass species contained in Oralair [(i.e., Sweet Vernal, Orchard, Perennial Rye, Timothy, and Kentucky Blue grass mixed pollens) or cross-reactive grass pollens (e.g., Cocksfoot, Meadow Fescue, or Redtop)]
- in vitro testing for pollen-specific IgE antibodies for any of the five grass species contained in Oralair [(i.e., Sweet Vernal, Orchard, Perennial Rye, Timothy, and Kentucky Blue grass mixed pollens) or cross-reactive grass pollens (e.g., Cocksfoot, Meadow Fescue, or Redtop)]

**AND**

**3** - Treatment is started or will be started at least 4 months before the beginning of the grass pollen season

**AND**

**4** - History of failure, contraindication, or intolerance to two of the following:

- oral antihistamine [e.g. cetirizine (Zyrtec)]
- intranasal antihistamine [e.g. azelastine (Astelin)]
- intranasal corticosteroid [e.g. fluticasone (Flonase)]
- leukotriene inhibitor [e.g. montelukast (Singulair)]

**AND**

**5** - Not received in combination with similar cross-reactive grass pollen immunotherapy (e.g., Grastek)

**AND**

**6** - Patient does not have unstable and/or uncontrolled asthma

**AND**

7 - Prescribed by or in consultation with a specialist in allergy and immunology

Product Name:Oralair

Diagnosis	Grass pollen-induced allergic rhinitis
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to Oralair therapy

Product Name:Ragwitek

Diagnosis	Short ragweed pollen-induced allergic rhinitis
-----------	--

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of moderate to severe short ragweed pollen-induced allergic rhinitis

**AND**

2 - Diagnosis confirmed by one of the following:

- Positive skin test to short ragweed pollen
- in vitro testing for pollen-specific IgE antibodies for short ragweed pollen

**AND**

**3** - Treatment is started or will be started at least 12 weeks before the beginning of the short ragweed pollen season

**AND**

**4** - History of failure, contraindication, or intolerance to two of the following:

- oral antihistamine [e.g. cetirizine (Zyrtec)]
- intranasal antihistamine [e.g. azelastine (Astelin)]
- intranasal corticosteroid [e.g. fluticasone (Flonase)]
- leukotriene inhibitor [e.g. montelukast (Singulair)]

**AND**

**5** - Patient does not have unstable and/or uncontrolled asthma

**AND**

**6** - Prescribed by or in consultation with a specialist in allergy and immunology

Product Name:Ragwitek	
Diagnosis	Short ragweed pollen-induced allergic rhinitis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Ragwitek therapy	

Product Name:Odactra	
Diagnosis	House dust mite (HDM)-induced allergic rhinitis
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of house dust mite (HDM)-induced allergic rhinitis.</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Diagnosis confirmed by one of the following:</p> <ul style="list-style-type: none"> <li>• Positive skin test to licensed house dust mite allergen extracts</li> <li>• in vitro testing for IgE antibodies to Dermatophagoides farinae or Dermatophagoides pteronyssinus house dust mites</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - History of failure, contraindication, or intolerance to two of the following:</p> <ul style="list-style-type: none"> <li>• oral antihistamine [e.g. cetirizine (Zyrtec)]</li> <li>• intranasal antihistamine [e.g. azelastine (Astelin)]</li> <li>• intranasal corticosteroid [e.g. fluticasone (Flonase)]</li> <li>• leukotriene inhibitor [e.g. montelukast (Singulair)]</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Patient does not have unstable and/or uncontrolled asthma</p> <p style="text-align: center;"><b>AND</b></p> <p>5 - Prescribed by or in consultation with a specialist in allergy and immunology</p> <p style="text-align: center;"><b>AND</b></p> <p>6 - Patient is between 5 and 20 years of age*</p>	

Notes	*Odactra is not covered in patients 21 years of age or older
-------	--

Product Name:Odactra	
Diagnosis	House dust mite (HDM)-induced allergic rhinitis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Odactra therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is between 5 and 20 years of age*</p>	
Notes	*Odactra is not covered in patients 21 years of age or older

## 2 . Revision History

Date	Notes
3/26/2025	Updated age criterion for Odactra due to expanded approval.

Sublocade, Brixadi - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157633
<b>Guideline Name</b>	Sublocade, Brixadi - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Brixadi, Sublocade	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following: <b>1.1</b> All of the following: <b>1.1.1</b> Patient has severe Opioid Use Disorder (OUD) as defined by the DSM-5 (Diagnostic	

and Statistical Manual of Mental Disorders, Fifth Edition) OUD Diagnostic Tool and has a demonstrated history of non-adherence to oral medications

**AND**

**1.1.2** Patient is currently maintained on 8mg to 24mg per day dose of oral, sublingual, or transmucosal buprenorphine product equivalent prior to initiation of Brixadi or Sublocade

**AND**

**1.1.3** Patient will not receive supplemental oral, sublingual, or transmucosal buprenorphine for greater than 6 weeks after initiation of Brixadi or Sublocade

**AND**

**1.1.4** Patient is receiving psychosocial interventions as part of a comprehensive medication assisted treatment (MAT) program

**AND**

**1.1.5** Prescriber checks the Arizona State Board of Pharmacy Controlled Substance Prescription Monitoring Program (CSPMP) database prior to each monthly injection of Brixadi or Sublocade

**AND**

**1.1.6** Dosing of Brixadi or Sublocade is in accordance with the U. S. Food and Drug Administration approved labeling

**OR**

**1.2** Brixadi or Sublocade is being requested due to circumstances other than non-adherence to oral medications. Document circumstance(s).

Product Name: Brixadi, Sublocade

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Physician documentation that the patient has experienced a positive clinical response to Brixadi or Sublocade therapy, as defined by the provider

**AND**

**2** - One of the following:

**2.1** All of the following:

**2.1.1** Patient will not receive supplemental oral, sublingual, or transmucosal buprenorphine for greater than 6 weeks after Brixadi or Sublocade therapy initiation

**AND**

**2.1.2** Patient is receiving psychosocial interventions as part of a comprehensive medication assisted treatment (MAT) program

**AND**

**2.1.3** Prescriber checks the Arizona State Board of Pharmacy Controlled Substance Prescription Monitoring Program (CSPMP) database prior to each monthly injection of Brixadi or Sublocade

**AND**

**2.1.4** Dosing of Brixadi or Sublocade is in accordance with the U. S. Food and Drug Administration approved labeling

**OR**

**2.2** Brixadi or Sublocade is being requested due to circumstances other than non-adherence to oral medications. Document circumstance(s).

## 2 . Revision History

Date	Notes
10/25/2024	Remove min day requirement for buprenorphine prior to initiation from 1.1.2

Suboxone - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-115713
<b>Guideline Name</b>	Suboxone - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/20/2022
-----------------	------------

## 1 . Criteria

Product Name:Generic buprenorphine-naloxone film	
Approval Length	N/A - Requests for generic buprenorphine hcl-naloxone film should not be approved
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Requests for generic buprenorphine-naloxone film are not authorized and will not be approved	
Notes	Approval Length: N/A - Requests for generic buprenorphine-naloxone film should not be approved. Patient need to use Brand Suboxone film or other preferred alternatives.

Product Name:Zubsolv, Bunavail	
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - The patient has a Diagnostic and Statistical Manual, Fifth Edition, Text Revision, (DSM-V-TR) diagnosis of opioid use disorder</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - The patient must have a reason or special circumstance that they cannot use the preferred products</p> <ul style="list-style-type: none"> <li>• brand Suboxone Film</li> <li>• buprenorphine (generic Subutex)</li> <li>• buprenorphine HCl/naloxone Tab (Generic Suboxone Tab)</li> <li>• naloxone</li> <li>• naltrexone</li> <li>• Narcan (naloxone)</li> <li>• Sublocade (buprenorphine)</li> <li>• Vivitrol (naltrexone microspheres)</li> </ul>	
Notes	*Up to 24 mg per day of Suboxone, or equivalent dosing of an alternative medication, will be authorized for the initial period.

Product Name:Zubsolv, Bunavail	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - The patient has been prescribed a buprenorphine product for the purpose of opioid use disorder maintenance therapy

**AND**

2 - The patient must have a reason or special circumstance that they cannot use the preferred products

**AND**

3 - Patient must have tried Suboxone film or buprenorphine-naloxone ODT tablets

Notes

\* Up to 16 mg per day of Suboxone, or equivalent dosing of an alternative medication, will be authorized for the reauthorization period.

Product Name: Brand suboxone, generic buprenorphine hcl-naloxone, buprenorphine/naloxone sublingual tablet, Zubsolv, Bunavail \*

Approval Length

3 month(s)

Therapy Stage

Initial Authorization

Guideline Type

Quantity Limit

**Approval Criteria**

1 - Physician has provided rationale for needing to exceed the buprenorphine daily limit

**AND**

2 - The requested dosage cannot be achieved using the plan accepted quantity limit of a different dose or formulation

Notes

\* This criteria applies to requests exceeding 24 mg of buprenorphine or equivalent

Product Name: Brand suboxone, generic buprenorphine hcl-naloxone, buprenorphine/naloxone sublingual tablet, Zubsolv, Bunavail \*

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - Physician has provided rationale for needing to exceed the buprenorphine daily limit</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - The requested dosage cannot be achieved using the plan accepted quantity limit of a different dose or formulation</p>	
Notes	*This criteria applies to requests exceeding 16 mg of buprenorphine or equivalent

## 2 . Revision History

Date	Notes
10/20/2022	Removed reference to PDL

Sucraid

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164414
<b>Guideline Name</b>	Sucraid
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Sucraid	
Approval Length	N/A- Requests for drugs not covered by Medicaid must be obtained through manufacturer for Compassionate Use
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of congenital sucrase-isomaltase deficiency (CSID) as confirmed by one of the following:	

**1.1** Duodenal biopsy showing low sucrose activity and normal amounts of other disaccharides

**OR**

**1.2** All of the following:

- Stool pH less than 6
- Negative lactose breath test
- Increase in breath hydrogen greater than 10 ppm (parts per million) when challenged with sucrose after fasting

**AND**

**2** - Prescribed by or in consultation with a gastroenterologist or rare disease specialist

**AND**

**3** - Will be used with a sucrose-free, low starch diet

**AND**

**4** - Provider attests that the requested medication will be obtained under compassionate use\*

Notes

\*Approval Length: N/A- Requests for drugs not covered by Medicaid must be obtained through manufacturer for Compassionate Use. Providers must contact QOL Medical 1-866-469-3773

Product Name: Sucraid

Approval Length

N/A- Requests for drugs not covered by Medicaid must be obtained through manufacturer for Compassionate Use

Therapy Stage

Reauthorization

Guideline Type

Prior Authorization

**Approval Criteria**

1 - Prescribed by or in consultation with a gastroenterologist or rare disease specialist

**AND**

2 - Will be used with a sucrose-free, low starch diet

**AND**

3 - Provider attests that the patient has achieved a clinically meaningful response while on Sucraid therapy, defined as at least a 50 percent reduction in all of the following:

- Symptoms of abdominal pain, cramps, bloating, gas, vomiting
- Number of stools per day
- Watery, loose stool consistency
- Number of symptomatic days

**AND**

4 - Provider attests that the requested medication will be obtained under compassionate use\*

Notes	*Approval Length: N/A- Requests for drugs not covered by Medicaid must be obtained through manufacturer for Compassionate Use. Providers must contact QOL Medical 1-866-469-3773
-------	--

## 2 . Revision History

Date	Notes
1/30/2025	Updated PA to remove approval duration, added note directing provider to contact mfg for compassionate use.

Sunlenca (lenacapavir sodium)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269197
<b>Guideline Name</b>	Sunlenca (lenacapavir sodium)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Sunlenca	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 Submission of medical records (e.g., chart notes) documenting all of the following:  1.1.1 Diagnosis of HIV-1 infection	

**AND**

**1.1.2** Both of the following:

**1.1.2.1** Patient is heavily treatment-experienced with multidrug resistance as confirmed by a resistance assay

**AND**

**1.1.2.2** Patient is failing their current antiretroviral regimen due to one of the following:

- Resistance
- Intolerance
- Safety considerations

**AND**

**1.1.3** Patient is currently taking, or will be prescribed, an active and optimized background antiretroviral therapy regimen

**AND**

**1.1.4** Prescribed by or in consultation with a clinician with HIV expertise

**OR**

**1.2** For continuation of prior therapy

## **2 . Revision History**

Date	Notes
5/29/2025	Added new GPI for Sunlenca tablet

Sunosi (solriamfetol)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-206195
<b>Guideline Name</b>	Sunosi (solriamfetol)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/27/2025
-----------------	-----------

## 1 . Criteria

Product Name:Sunosi	
Diagnosis	Narcolepsy
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g. chart notes, lab values) documenting a diagnosis of narcolepsy with BOTH of the following:	

**1.1** The patient has daily periods of irrepressible need to sleep or daytime lapses into sleep occurring for at least three months.

**AND**

**1.2** A mean sleep latency of less than or equal to 8 minutes and two or more sleep onset rapid eye movement (REM) periods (SOREMPs) are found on a multiple sleep latency test (MSLT) performed according to standard techniques following a normal overnight polysomnogram. A SOREMP (within 15 minutes of sleep onset) on the preceding nocturnal polysomnogram may replace one of the SOREMPs on the MSLT.

**AND**

**2** - Physician attestation to the following:

- Other causes of sleepiness have been ruled out or treated (including but not limited to obstructive sleep apnea, insufficient sleep syndrome, shift work, the effects of substances or medications or their withdrawal, sleep phase disorder, or other sleep disorders)

**AND**

**3** - History of failure, contraindication, or intolerance to BOTH of the following:

**3.1** ONE of the following:

- Amphetamine based stimulant (e.g., amphetamine, dextroamphetamine)
- Methylphenidate based stimulant

**AND**

**3.2** Armodafinil

**AND**

**4** - Prescribed by one of the following:

- Neurologist

- Psychiatrist
- Sleep Medicine Specialist

Product Name:Sunosi	
Diagnosis	Narcolepsy
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Reduction in symptoms of excessive daytime sleepiness associated with Sunosi therapy</p>	

Product Name:Sunosi	
Diagnosis	Obstructive Sleep Apnea
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g. chart notes, lab values) documenting a diagnosis of obstructive sleep apnea with ONE of the following:</p> <p>1.1 Fifteen or more obstructive respiratory events per hour of sleep confirmed by a sleep study</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 BOTH of the following:</p> <p>1.2.1 Five or more obstructive respiratory events per hour of sleep confirmed by a sleep study</p>	

**AND**

**1.2.2** ONE or more of the following sign/symptoms are present:

- Daytime sleepiness
- Nonrestorative sleep
- Fatigue
- Insomnia
- Waking up with breath holding, gasping, or choking
- Habitual snoring noted by bed partner or other observer
- Observed apnea

**AND**

**2** - BOTH of the following:

**2.1** Standard treatments for the underlying airway obstruction (e.g., continuous positive airway pressure [CPAP], bi-level positive airway pressure [BiPAP]) have been used for one month or longer

**AND**

**2.2** Patient is fully compliant with ongoing treatment(s) for the underlying airway obstruction

**AND**

**3** - History of failure, contraindication, or intolerance to armodafinil

**AND**

**4** - Prescribed by one of the following:

- Neurologist
- Psychiatrist
- Sleep Medicine Specialist

Product Name:Sunosi	
Diagnosis	Obstructive Sleep Apnea
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Reduction in symptoms of excessive daytime sleepiness associated with Sunosi therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient continues to be fully compliant with ongoing treatment(s) for the underlying airway obstruction (e.g. continuous positive airway pressure [CPAP], bi-level positive airway pressure [BiPAP])</p>	

## 2 . Revision History

Date	Notes
2/27/2025	Narcolepsy initial auth: Corrected connector for 1.1 /1.2 to AND.

Sutent

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99767
<b>Guideline Name</b>	Sutent
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Sutent	
Diagnosis	Gastrointestinal Stromal Tumor (GIST)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of gastrointestinal stromal tumor (GIST)	

**AND**

**2** - History of failure, contraindication, or intolerance to Gleevec (imatinib)

Product Name:Sutent	
Diagnosis	Renal Cell Carcinoma (RCC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of renal cell carcinoma (RCC)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <p>2.1 Disease has relapsed</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Diagnosis of Stage IV disease</p> <p style="text-align: center;"><b>OR</b></p> <p>2.3 BOTH of the following:</p> <p>2.3.1 Used in adjuvant setting</p> <p style="text-align: center;"><b>AND</b></p>	

**2.3.2** Patient has a high risk of recurrence following nephrectomy

Product Name:Sutent	
Diagnosis	Islet Cell Tumor / Progressive Pancreatic Neuroendocrine Tumors (pNET)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of islet cell tumor / progressive pancreatic neuroendocrine tumors (pNET)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Disease is ONE of the following:</p> <ul style="list-style-type: none"><li>• Unresectable, locally advanced</li><li>• Metastatic</li></ul>	

Product Name:Sutent	
Diagnosis	Soft Tissue Sarcoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of ONE of the following:</p> <ul style="list-style-type: none"><li>• Alveolar soft part sarcoma (ASPS)</li><li>• Angiosarcoma</li></ul>	

- Solitary fibrous tumor / hemangiopericytoma

Product Name: Sutent	
Diagnosis	Thyroid Carcinoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 ALL of the following:</p> <p>1.1.1 Diagnosis of ONE of the following:</p> <ul style="list-style-type: none"> <li>• Follicular carcinoma</li> <li>• Hürthle cell carcinoma</li> <li>• Papillary carcinoma</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>1.1.2 ONE of the following:</p> <ul style="list-style-type: none"> <li>• Unresectable locoregional recurrent disease</li> <li>• Persistent disease</li> <li>• Metastatic disease</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>1.1.3 ONE of the following:</p> <ul style="list-style-type: none"> <li>• Patient has symptomatic disease</li> <li>• Patient has progressive disease</li> </ul>	

**AND**

**1.1.4** Disease is refractory to radioactive iodine treatment

**OR**

**1.2** ALL of the following:

**1.2.1** Diagnosis of medullary thyroid carcinoma

**AND**

**1.2.2** ONE of the following:

- Patient has progressive disease
- Patient has symptomatic metastatic disease

**AND**

**1.2.3** History of failure, contraindication, or intolerance to ONE of the following:

- Caprelsa (vandetanib)
- Cometriq (cabozantinib)

<b>Product Name: Sutent</b>	
Diagnosis	Chordoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of recurrent chordoma	

Product Name:Sutent	
Diagnosis	Central Nervous System Cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of surgically inaccessible meningiomas</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Disease is recurrent</li> <li>• Disease is progressive</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Further radiation is not possible</p>	

Product Name:Sutent	
Diagnosis	Thymic Carcinoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of thymic carcinoma</p>	

**AND**

**2** - Used as second-line following a failure, contraindication, or intolerance to a first-line chemotherapy regimen (e.g., carboplatin/paclitaxel)

Product Name:Sutent	
Diagnosis	Gastrointestinal Stromal Tumor (GIST), Renal Cell Carcinoma (RCC), Islet Cell Tumor / Progressive Pancreatic Neuroendocrine Tumors (pNET), Soft Tissue Sarcoma, Thyroid Carcinoma, Chordoma, Central Nervous System Cancer, Thymic Carcinoma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on Sutent therapy	

Product Name:Sutent	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Sutent will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.	

Product Name:Sutent	
Diagnosis	NCCN Recommended Regimens

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Sutent therapy</p>	

## 2 . Revision History

Date	Notes
6/2/2021	Arizona Medicaid 7.1 Implementation

Syfovre (pegcetacoplan)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-124881
<b>Guideline Name</b>	Syfovre (pegcetacoplan)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Syfovre	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of geographic atrophy (GA) secondary to age-related macular degeneration (AMD) as confirmed by one of the following: <ul style="list-style-type: none"><li>Fundus photography (e.g. fundus autofluorescence [FAF])</li></ul>	

- Optical coherence tomography (OCT)
- Fluorescein angiography

**AND**

**2** - GA is not secondary to any other conditions (e.g., Stargardt disease, cone rod dystrophy, toxic maculopathies)

**AND**

**3** - Prescribed by or in consultation with an ophthalmologist experienced in the treatment of retinal diseases

Product Name: Syfovre	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., reduction in growth rate of GA lesion)</p>	

## 2 . Revision History

Date	Notes
4/20/2023	New Program

Symdeko

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99649
<b>Guideline Name</b>	Symdeko
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Symdeko	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystic fibrosis (CF)  <b>AND</b>	

**2** - Submission of laboratory result documenting ONE of the following:

**2.1** The patient is homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene

**OR**

**2.2** The patient has at least ONE mutation in the CFTR gene that is responsive to Symdeko (See Table in Background Section)

**AND**

**3** - The patient is greater than or equal to 6 years of age

**AND**

**4** - Prescribed by or in consultation with a specialist affiliated with a CF care center

Product Name: Symdeko

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Provider attests that the patient has achieved a clinically meaningful response while on Symdeko therapy to ONE of the following:

- Lung function as demonstrated by percent predicted expiratory volume in 1 second (ppFEV1)
- Body mass index (BMI)
- Pulmonary exacerbations
- Quality of life as demonstrated by Cystic Fibrosis Questionnaire-Revised (CFQ-R) respiratory domain score

**AND**

2 - Prescribed by, or in consultation with, a specialist affiliated with a cystic fibrosis (CF) care center

## 2 . Background

### Benefit/Coverage/Program Information

**Table 1 CFTR Gene Mutations**

A1067T	D1270N	F1052V	R1070W	S945L	3272-26A→G
A455E	D579G	F1074L	R117C	S977F	3849+10kbC→T
D110E	E193K	K1060T	R347H		711+3A→G
D110H	E56K	L206W	R352Q		2789+5G→A
D1152H	E831X	P67L	R74W		

## 3 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Symlin

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99499
<b>Guideline Name</b>	Symlin
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Symlin	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Patient must have ONE of the following diagnoses: <ul style="list-style-type: none"><li>Type 1 diabetes</li><li>Type 2 diabetes</li></ul>	

**AND**

2 - Concurrent use of insulin therapy

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Synagis

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-117156
<b>Guideline Name</b>	Synagis
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/21/2022
-----------------	------------

### Note:

**\*\*PLEASE NOTE: PA IS NOT REQUIRED FOR CHILDREN UNDER 2 YEARS OF AGE\*\***

## 1 . Criteria

Product Name:Synagis*	
Diagnosis	Prematurity
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - BOTH of the following:  1.1 Patient is an infant born before 29 weeks, 0 days gestation	

**AND**

**1.2** Patient is less than 12 months of age at the start of RSV “season”

**AND**

**2** - Administered during RSV season\*\*

**AND**

**3** - Monthly dose of Synagis does not exceed 15 milligram per kilogram per dose

**AND**

**4** - Monthly dose of Synagis does not exceed 5 doses per single RSV “season”\*\*\*

**AND**

**5** - The patient does not meet ONE of the following situations

- Infants and children with hemodynamically insignificant heart disease (e.g., secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus)
- Infants with congenital heart disease and cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure
- Infants with cardiomyopathy sufficiently mild that they do not require pharmacotherapy
- Routine use of prophylaxis in children with Down syndrome [unless qualifying heart disease, CLD, airway clearance issues (the inability to clear secretions from the upper airway because of ineffective cough), or prematurity (less than 29 weeks, 0 days gestation) is present]
- Routine use of prophylaxis in children with cystic fibrosis (unless indications noted in proven indications above are present)
- Administration of monthly Synagis prophylaxis after an infant or child has experienced a breakthrough RSV hospitalization during the current season if child had met criteria for palivizumab
- Prophylaxis for primary asthma prevention or to reduce subsequent episodes of wheezing in infants and children
- Synagis prophylaxis for prevention of nosocomial disease

<ul style="list-style-type: none"> <li>Treatment of symptomatic RSV disease</li> </ul>	
Notes	<p>*NOTE: Approval for up to 5 doses per single RSV “season”</p> <p>** Information regarding RSV season may be found at:</p> <ul style="list-style-type: none"> <li>Centers for Disease and Prevention (CDC) surveillance reports (<a href="http://www.cdc.gov/surveillance/nrevss/rsv/index.html">http://www.cdc.gov/surveillance/nrevss/rsv/index.html</a>)</li> <li><a href="http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx">http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx</a></li> </ul> <p>***NOTE: Infants in a neonatal intensive care unit who qualify for prophylaxis may receive the first dose 48 to 72 hours before discharge to home or promptly after discharge. If the first dose is administered in the hospital, this dose will be considered the first dose of the maximum 5 dose series for the season. And any subsequent doses received in the hospital setting, are also considered as part of the maximum 5 dose series. For infants born during the RSV “season,” fewer than 5 monthly doses may be needed.</p>

Product Name:Synagis*	
Diagnosis	Chronic Lung Disease (CLD)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 ALL of the following for patients age 0 to less than 12 months:</p> <p>1.1.1 The patient is a preterm infant defined as gestational age less than 32 weeks, 0 days</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.2 Patient has developed chronic lung disease (CLD) of prematurity</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.3 There was a requirement for greater than 21% oxygen for at least the first 28 days after birth</p>	

**OR**

**1.2** ALL of the following for patients age greater than or equal to 12 months to less than 24 months:

**1.2.1** The patient was born at less than 32 weeks, 0 days gestation

**AND**

**1.2.2** The patient required at least 28 days of oxygen after birth

**AND**

**1.2.3** The patient continues to require supplemental oxygen, diuretics, or chronic systemic corticosteroid therapy within 6 months of the start of the second RSV “season”

**AND**

**2** - Administered during RSV season\*\*

**AND**

**3** - Monthly dose of Synagis does not exceed 15 milligram per kilogram per dose

**AND**

**4** - Monthly dose of Synagis does not exceed 5 doses per single RSV “season”\*\*\*

**AND**

**5** - The patient does not meet ONE of the following situations

- Infants and children with hemodynamically insignificant heart disease (e.g., secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus)
- Infants with congenital heart disease and cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure
- Infants with cardiomyopathy sufficiently mild that they do not require pharmacotherapy
- Routine use of prophylaxis in children with Down syndrome [unless qualifying heart disease, CLD, airway clearance issues (the inability to clear secretions from the upper airway because of ineffective cough), or prematurity (less than 29 weeks, 0 days gestation) is present]
- Routine use of prophylaxis in children with cystic fibrosis (unless indications noted in proven indications above are present)
- Administration of monthly Synagis prophylaxis after an infant or child has experienced a breakthrough RSV hospitalization during the current season if child had met criteria for palivizumab
- Prophylaxis for primary asthma prevention or to reduce subsequent episodes of wheezing in infants and children
- Synagis prophylaxis for prevention of nosocomial disease
- Treatment of symptomatic RSV disease

Notes	<p>*NOTE: Approval for up to 5 doses per single RSV “season”</p> <p>** Information regarding RSV season may be found at:</p> <ul style="list-style-type: none"> <li>• Centers for Disease and Prevention (CDC) surveillance reports (<a href="http://www.cdc.gov/surveillance/nrevss/rsv/index.html">http://www.cdc.gov/surveillance/nrevss/rsv/index.html</a>)</li> <li>• <a href="http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx">http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx</a></li> </ul> <p>***NOTE: Infants in a neonatal intensive care unit who qualify for prophylaxis may receive the first dose 48 to 72 hours before discharge to home or promptly after discharge. If the first dose is administered in the hospital, this dose will be considered the first dose of the maximum 5 dose series for the season. And any subsequent doses received in the hospital setting, are also considered as part of the maximum 5 dose series. For infants born during the RSV “season,” fewer than 5 monthly doses may be needed.</p>
-------	--

Product Name: Synagis*	
Diagnosis	Congenital Heart Disease (CHD)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 ONE of the following for patients age 0 to less than 12 months:</p>	

**1.1.1** Patient has hemodynamically significant congenital heart disease (CHD) including ONE of the following:

- Acyanotic heart disease and receiving medication to control congestive heart failure and will require cardiac surgical procedures
- Moderate to severe pulmonary hypertension
- Documentation that decisions regarding prophylaxis for infants with cyanotic heart defects were made in consultation with a pediatric cardiologist

**OR**

**1.1.2** The patient is undergoing cardiac transplantation during the RSV “season”

**OR**

**1.2** BOTH of the following:

**1.2.1** The patient is greater than or equal to 12 months to less than 24 months of age:

**AND**

**1.2.2** ONE of the following:

- After cardiac bypass
- At the conclusion of extracorporeal membrane oxygenation
- The patient is undergoing cardiac transplantation during the RSV “season”

**AND**

**2** - Administered during RSV season\*\*

**AND**

**3** - Monthly dose of Synagis does not exceed 15 milligram per kilogram per dose

**AND**

4 - Monthly dose of Synagis does not exceed 5 doses per single RSV "season"\*\*\*

**AND**

5 - The patient does not meet ONE of the following situations

- Infants and children with hemodynamically insignificant heart disease (e.g., secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus)
- Infants with congenital heart disease and cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure
- Infants with cardiomyopathy sufficiently mild that they do not require pharmacotherapy
- Routine use of prophylaxis in children with Down syndrome [unless qualifying heart disease, CLD, airway clearance issues (the inability to clear secretions from the upper airway because of ineffective cough), or prematurity (less than 29 weeks, 0 days gestation) is present]
- Routine use of prophylaxis in children with cystic fibrosis (unless indications noted in proven indications above are present)
- Administration of monthly Synagis prophylaxis after an infant or child has experienced a breakthrough RSV hospitalization during the current season if child had met criteria for palivizumab
- Prophylaxis for primary asthma prevention or to reduce subsequent episodes of wheezing in infants and children
- Synagis prophylaxis for prevention of nosocomial disease
- Treatment of symptomatic RSV disease

Notes

\*NOTE: Approval for up to 5 doses per single RSV "season"  
\*\* Information regarding RSV season may be found at:  
• Centers for Disease and Prevention (CDC) surveillance reports (<http://www.cdc.gov/surveillance/nrevss/rsv/index.html>)  
• <http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx>  
\*\*\*NOTE: Infants in a neonatal intensive care unit who qualify for prophylaxis may receive the first dose 48 to 72 hours before discharge to home or promptly after discharge. If the first dose is administered in the hospital, this dose will be considered the first dose of the maximum 5 dose series for the season. And any subsequent doses received in the hospital setting, are also considered as part of the maximum 5 dose series. For infants born during the RSV "season," fewer than 5 monthly doses may be needed.

Product Name: Synagis\*

Diagnosis Congenital abnormalities of the airway or neuromuscular disease

Guideline Type Prior Authorization

## **Approval Criteria**

**1** - ALL of the following:

**1.1** Patient is age 0 to less than 12 months

**AND**

**1.2** Patient has ONE of the following:

- Neuromuscular disease
- A congenital anomaly that impairs the ability to clear secretions from the lower airway because of ineffective cough

**AND**

**2** - Administered during RSV season\*\*

**AND**

**3** - Monthly dose of Synagis does not exceed 15 milligram per kilogram per dose

**AND**

**4** - Monthly dose of Synagis does not exceed 5 doses per single RSV "season"\*\*\*

**AND**

**5** - The patient does not meet ONE of the following situations

- Infants and children with hemodynamically insignificant heart disease (e.g., secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus)
- Infants with congenital heart disease and cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure
- Infants with cardiomyopathy sufficiently mild that they do not require pharmacotherapy

<ul style="list-style-type: none"> <li>• Routine use of prophylaxis in children with Down syndrome [unless qualifying heart disease, CLD, airway clearance issues (the inability to clear secretions from the upper airway because of ineffective cough), or prematurity (less than 29 weeks, 0 days gestation) is present]</li> <li>• Routine use of prophylaxis in children with cystic fibrosis (unless indications noted in proven indications above are present)</li> <li>• Administration of monthly Synagis prophylaxis after an infant or child has experienced a breakthrough RSV hospitalization during the current season if child had met criteria for palivizumab</li> <li>• Prophylaxis for primary asthma prevention or to reduce subsequent episodes of wheezing in infants and children</li> <li>• Synagis prophylaxis for prevention of nosocomial disease</li> <li>• Treatment of symptomatic RSV disease</li> </ul>	
Notes	<p>*NOTE: Approval for up to 5 doses per single RSV “season”</p> <p>** Information regarding RSV season may be found at:</p> <ul style="list-style-type: none"> <li>• Centers for Disease and Prevention (CDC) surveillance reports (<a href="http://www.cdc.gov/surveillance/nrevss/rsv/index.html">http://www.cdc.gov/surveillance/nrevss/rsv/index.html</a>)</li> <li>• <a href="http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx">http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx</a></li> </ul> <p>***NOTE: Infants in a neonatal intensive care unit who qualify for prophylaxis may receive the first dose 48 to 72 hours before discharge to home or promptly after discharge. If the first dose is administered in the hospital, this dose will be considered the first dose of the maximum 5 dose series for the season. And any subsequent doses received in the hospital setting, are also considered as part of the maximum 5 dose series. For infants born during the RSV “season,” fewer than 5 monthly doses may be needed.</p>

Product Name: Synagis*	
Diagnosis	Immunocompromised children less than 24 months of age
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - BOTH of the following:</p> <p>1.1 Patient is less than 24 months of age</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 The patient is immunocompromised (e.g. receiving cancer chemotherapy, undergoing hematopoietic stem cell transplantation, or solid organ transplantation)</p>	

**AND**

**2** - Administered during RSV season\*\*

**AND**

**3** - Monthly dose of Synagis does not exceed 15 milligram per kilogram per dose

**AND**

**4** - Monthly dose of Synagis does not exceed 5 doses per single RSV “season”\*\*\*

**AND**

**5** - The patient does not meet ONE of the following situations

- Infants and children with hemodynamically insignificant heart disease (e.g., secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus)
- Infants with congenital heart disease and cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure
- Infants with cardiomyopathy sufficiently mild that they do not require pharmacotherapy
- Routine use of prophylaxis in children with Down syndrome [unless qualifying heart disease, CLD, airway clearance issues (the inability to clear secretions from the upper airway because of ineffective cough), or prematurity (less than 29 weeks, 0 days gestation) is present]
- Routine use of prophylaxis in children with cystic fibrosis (unless indications noted in proven indications above are present)
- Administration of monthly Synagis prophylaxis after an infant or child has experienced a breakthrough RSV hospitalization during the current season if child had met criteria for palivizumab
- Prophylaxis for primary asthma prevention or to reduce subsequent episodes of wheezing in infants and children
- Synagis prophylaxis for prevention of nosocomial disease
- Treatment of symptomatic RSV disease

Notes

\*NOTE: Approval for up to 5 doses per single RSV “season”

\*\* Information regarding RSV season may be found at:

• Centers for Disease and Prevention (CDC) surveillance reports (<http>

	<p>://www.cdc.gov/surveillance/nrevss/rsv/index.html)</p> <ul style="list-style-type: none"> <li>• <a href="http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx">http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx</a></li> </ul> <p>***NOTE: Infants in a neonatal intensive care unit who qualify for prophylaxis may receive the first dose 48 to 72 hours before discharge to home or promptly after discharge. If the first dose is administered in the hospital, this dose will be considered the first dose of the maximum 5 dose series for the season. And any subsequent doses received in the hospital setting, are also considered as part of the maximum 5 dose series. For infants born during the RSV "season," fewer than 5 monthly doses may be needed.</p>
--	--

Product Name:Synagis*	
Diagnosis	Cystic fibrosis (CF)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 BOTH of the following for patients age 0 to less than 12 months:</p> <p>1.1.1 Patient has cystic fibrosis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.1.2 Patient has clinical evidence of at least ONE of the following:</p> <ul style="list-style-type: none"> <li>• Chronic lung disease (CLD)</li> <li>• Nutritional compromise</li> <li>• Failure to thrive defined as weight for length less than the 10th percentile on a pediatric growth chart</li> </ul> <p style="text-align: center;"><b>OR</b></p> <p>1.2 BOTH of the following:</p> <p>1.2.1 Patient is greater than or equal to 12 months to less than 24 months of age</p>	

**AND**

**1.2.2** Patient has manifestations of severe lung disease including ONE of the following:

- Previous hospitalization for pulmonary exacerbation in the first year of life
- Abnormalities on chest radiography or chest computed tomography that persists when stable
- Weight for length less than the 10th percentile on a pediatric growth chart

**AND**

**2** - Administered during RSV season\*\*

**AND**

**3** - Monthly dose of Synagis does not exceed 15 milligram per kilogram per dose

**AND**

**4** - Monthly dose of Synagis does not exceed 5 doses per single RSV "season"\*\*\*

**AND**

**5** - The patient does not meet ONE of the following situations

- Infants and children with hemodynamically insignificant heart disease (e.g., secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus)
- Infants with congenital heart disease and cardiac lesions adequately corrected by surgery, unless they continue to require medication for congestive heart failure
- Infants with cardiomyopathy sufficiently mild that they do not require pharmacotherapy
- Routine use of prophylaxis in children with Down syndrome [unless qualifying heart disease, CLD, airway clearance issues (the inability to clear secretions from the upper airway because of ineffective cough), or prematurity (less than 29 weeks, 0 days gestation) is present]
- Routine use of prophylaxis in children with cystic fibrosis (unless indications noted in proven indications above are present)

	<ul style="list-style-type: none"> <li>• Administration of monthly Synagis prophylaxis after an infant or child has experienced a breakthrough RSV hospitalization during the current season if child had met criteria for palivizumab</li> <li>• Prophylaxis for primary asthma prevention or to reduce subsequent episodes of wheezing in infants and children</li> <li>• Synagis prophylaxis for prevention of nosocomial disease</li> <li>• Treatment of symptomatic RSV disease</li> </ul>
Notes	<p>*NOTE: Approval for up to 5 doses per single RSV “season”</p> <p>** Information regarding RSV season may be found at:</p> <ul style="list-style-type: none"> <li>• Centers for Disease and Prevention (CDC) surveillance reports (<a href="http://www.cdc.gov/surveillance/nrevss/rsv/index.html">http://www.cdc.gov/surveillance/nrevss/rsv/index.html</a>)</li> <li>• <a href="http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx">http://uhc-cs-10.uhc.com/sites/cspm/CSSP/Pages/Synagis.aspx</a></li> </ul> <p>***NOTE: Infants in a neonatal intensive care unit who qualify for prophylaxis may receive the first dose 48 to 72 hours before discharge to home or promptly after discharge. If the first dose is administered in the hospital, this dose will be considered the first dose of the maximum 5 dose series for the season. And any subsequent doses received in the hospital setting, are also considered as part of the maximum 5 dose series. For infants born during the RSV “season,” fewer than 5 monthly doses may be needed.</p>

## 2 . Background

Benefit/Coverage/Program Information
<p><b>Additional Information</b></p> <p>In most of North America, peak RSV activity typically occurs between November and March, usually beginning in November or December, peaking in January or February, and ending by the end of March or sometime in April. Communities in the southern United States, particularly some communities in the state of Florida, tend to experience the earliest onset of RSV. Data from the Centers for Disease Control and Prevention (CDC) have identified variations in the onset and offset of the RSV “season” in the state of Florida that could affect the timing of Synagis administration. <sup>10</sup></p> <ul style="list-style-type: none"> <li>• Despite varied onsets, the RSV “season” is of the same duration (5 months) in the different regions of Florida.</li> <li>• On the basis of the epidemiology of RSV in Alaska, particularly in remote regions where the burden of RSV disease is significantly greater than the general US population, the selection of Alaska Native infants eligible for prophylaxis may differ from the remainder of the United States. Clinicians may wish to use RSV surveillance data generated by the state of Alaska to assist in determining onset and end of the RSV season for qualifying infants.</li> </ul>

- Limited information is available concerning the burden of RSV disease among Native American populations. However, special consideration may be prudent for Navajo and White Mountain Apache infants in the first year of life.

For analysis of National Respiratory and Enteric Virus Surveillance System (NREVSS) reports in the CDC Morbidity and Mortality Weekly Report, season onset is defined as the first of 2 consecutive weeks during which the mean percentage of specimens testing positive for RSV antigen is  $\geq 10\%$  and RSV “season” offset is defined as the last of 2 consecutive weeks during which the mean percentage of positive specimens is  $\geq 10\%$ . Use of specimens to determine the start of the RSV “season” requires that the number of specimens tested be statistically significant.

### 3 . Revision History

Date	Notes
11/21/2022	Added note to guideline, PA not required for children under 2 yo

Systane, Refresh, Gonak, Genteal, Tears Naturale

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99534
<b>Guideline Name</b>	Systane, Refresh, Gonak, Genteal, Tears Naturale
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:brand Systane, brand Refresh, brand Gonak, brand Genteal, Tears Naturale	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, contraindication, or intolerance to ALL of the following: <ul style="list-style-type: none"><li>Generic equivalents for drops, ointments and gel formulations for Systane, Refresh, Gonak, Genteal, Tears Naturale, and Generic equivalent to the requested brand product</li><li>sodium chloride ophthalmic ointment</li></ul>	

## 2 . Revision History

Date	Notes
5/20/2021	Arizona Medicaid 7.1 Implementation

Talicia and rifabutin

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-164155
<b>Guideline Name</b>	Talicia and rifabutin
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2025
-----------------	----------

## 1 . Criteria

Product Name:generic rifabutin	
Diagnosis	Mycobacterium Avium Complex Prophylaxis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Mycobacterium Avium Complex Prophylaxis	

**AND**

**2** - Prescribed by or in consultation with an HIV or infectious disease specialist

**AND**

**3** - Member has failed azithromycin or clarithromycin or is intolerant to the medication due to significant adverse effects or both are contraindicated

**AND**

**4** - The requested dosage does not exceed 450 mg per day

Product Name:generic rifabutin	
Diagnosis	Mycobacterium Avium Complex Prophylaxis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Member is responding positively to therapy	

Product Name:generic rifabutin	
Diagnosis	Mycobacterium Avium Complex Prophylaxis
Approval Length	12 month(s)
Guideline Type	Quantity Limit
<b>Approval Criteria</b>	

1 - For doses that exceed 450mg, the use of this drug is supported by information from ONE of the following appropriate compendia of current literature:

- Food and Drug Administration (FDA) approved indications and limits
- Published practice guidelines and treatment protocols
- Comparative data evaluating the efficacy, type and frequency of side effects and potential drug interactions among alternative products as well as the risks, benefits and potential member outcomes
- Drug Facts and Comparisons
- American Hospital Formulary Service Drug Information
- United States Pharmacopeia – Drug Information
- DRUGDEX Information System
- UpToDate
- MicroMedex
- Peer-reviewed medical literature, including randomized clinical trials, outcomes, research data and pharmaco-economic studies
- Other drug reference resources

Product Name:generic rifabutin	
Diagnosis	Helicobacter pylori Infection (off-label)
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of H. pylori infection</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed in combination with amoxicillin and a proton pump inhibitor</p>	

Product Name:Talicia	
Diagnosis	Helicobacter pylori Infection
Approval Length	14 Day(s)
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Diagnosis of H. pylori infection

**AND**

**2** - The medication is prescribed by or in consultation with a gastroenterologist or infectious disease specialist

**AND**

**3** - One of the following:

**3.1** Both of the following:

**3.1.1** Trial and failure, contraindication, or intolerance to Bismuth quadruple therapy (e.g., bismuth and metronidazole and tetracycline and proton pump inhibitor [PPI])

**AND**

**3.1.2** Trial and failure, contraindication, or intolerance to BOTH of the following suggested first line treatment regimens:

- Clarithromycin based therapy (e.g., clarithromycin based triple therapy, clarithromycin based concomitant therapy) or PCAB-clarithromycin triple therapy
- Rifabutin triple therapy

**OR**

**3.2** Both of the following:

**3.2.1** Culture and sensitivity report indicate resistance or lack of susceptibility of H. pylori to all first-line treatment regimens except Rifabutin triple therapy

**AND**

**3.2.2** Member must have tried and failed rifabutin triple therapy

Product Name:generic rifabutin	
Diagnosis	Tuberculosis (off-label)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of tuberculosis infection</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with an HIV or infectious disease specialist</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Current treatment with protease inhibitors or non-nucleoside reverse transcriptase inhibitors (NNRTIs) for the treatment of HIV infection</p>	

Product Name:generic rifabutin	
Diagnosis	Tuberculosis (off-label)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Member is responding positively to therapy</p>	

## 2 . Revision History

Date	Notes
1/30/2025	Updated guideline targets, removed dosing chart

Taltz (ixekizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-329256
<b>Guideline Name</b>	Taltz (ixekizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Taltz	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:  1.1 Diagnosis of chronic moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3 percent body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** History of failure to ONE of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies:

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with a dermatologist

Product Name:Taltz

Diagnosis	Plaque Psoriasis (PsO)
-----------	------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Documentation of positive clinical response to therapy

**AND**

**2** - Prescribed by or in consultation with a dermatologist

Product Name:Taltz

Diagnosis	Psoriatic Arthritis (PsA)
-----------	---------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

**1.1** Diagnosis of active psoriatic arthritis (PsA)

**AND**

**1.2** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following preferred biologic products

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Orenzia (abatacept)
- Otezla (apremilast)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Product Name:Taltz	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

**AND**

**2** - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Product Name:Taltz	
Diagnosis	Ankylosing Spondylitis (AS), Non-radiographic axial spondyloarthritis (nr-axSpA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

**1.1** One of the following diagnoses:

- Active ankylosing spondylitis (AS)
- Active non-radiographic axial spondyloarthritis (nr-axSpA)

**AND**

**1.2** History of failure to TWO nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen) at maximally indicated doses, each used for at least 4 weeks within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name:Taltz	
Diagnosis	Ankylosing Spondylitis (AS), Non-radiographic axial spondyloarthritis (nr-axSpA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p>	

## 2 . Revision History

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, updated criteria throughout. PsA, added 'or verification of paid claims'

Tarceva

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99779
<b>Guideline Name</b>	Tarceva
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	Pancreatic Cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of pancreatic cancer	

**AND**

**2** - Disease is ONE of the following:

- Locally advanced
- Unresectable
- Metastatic

**AND**

**3** - Used in combination with Gemzar (gemcitabine)

Product Name: Brand Tarceva, generic erlotinib	
Diagnosis	Pancreatic Cancer
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on Tarceva therapy	

Product Name: Brand Tarceva, generic erlotinib	
Diagnosis	Non-Small Cell Lung Cancer (NSCLC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of non-small cell lung cancer (NSCLC)	

**AND**

**2** - Disease is ONE of the following:

- Metastatic
- Recurrent

**AND**

**3** - ONE of the following:

- Tumors are positive for epidermal growth factor receptor (EGFR)exon 19 deletions
- Tumors are positive for exon 21 (L858R) substitution mutations
- Tumors are positive for a known sensitizing EGFR mutation (e.g. in-frame exon 20 insertions, exon 18 G719 mutation, exon 21 L861Q mutation)

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	Non-Small Cell Lung Cancer (NSCLC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on Tarceva therapy	

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	Chordoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of chordoma

Product Name: Brand Tarceva, generic erlotinib

Diagnosis	Chordoma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient does not show evidence of progressive disease while on Tarceva therapy

Product Name: Brand Tarceva, generic erlotinib

Diagnosis	Kidney Cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Both of the following:

- Diagnosis of kidney cancer
- Disease is stage IV or relapsed

**AND**

2 - Disease is of non-clear cell histology

Product Name: Brand Tarceva, generic erlotinib	
Diagnosis	Kidney Cancer
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Tarceva therapy</p>	

Product Name: Brand Tarceva, generic erlotinib	
Diagnosis	Central Nervous System (CNS) Cancers
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of metastatic brain cancer from Non-Small Cell Lung Cancer (NSCLC)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Tumors are positive for epidermal growth factor receptor (EGFR) exon 19 deletions</li> <li>• Tumors are positive for exon 21 (L858R) substitution mutations</li> <li>• Tumors are positive for a known sensitizing EGFR mutation (e.g., in-frame exon 20 insertions, exon 18 G719 mutation, exon 21 L861Q mutation)</li> </ul>	

Product Name: Brand Tarceva, generic erlotinib	
Diagnosis	Central Nervous System (CNS) Cancers
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Tarceva therapy</p>	

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	Vulvar cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of vulvar cancer</p>	

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	Vulvar cancer
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Tarceva therapy</p>	

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Tarceva will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.</p>	

Product Name:Brand Tarceva, generic erlotinib	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Tarceva therapy</p>	

## 2 . Revision History

Date	Notes
6/2/2021	Arizona Medicaid 7.1 Implementation

Targretin

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99771
<b>Guideline Name</b>	Targretin
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Targretin caps, generic bexarotene caps, Targretin gel	
Diagnosis	Cutaneous T-Cell Lymphoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of cutaneous T-cell lymphoma (CTCL)	

**AND**

**2** - History of failure, contraindication, or intolerance to at least one prior therapy (including skin-directed therapies [e.g., corticosteroids (clobetasol, diflorasone, halobetasol, augmented betamethasone dipropionate), phototherapy, or systemic therapies [e.g. Interferons])

Product Name: Brand Targretin caps, generic bexarotene caps, Targretin gel	
Diagnosis	Cutaneous T-Cell Lymphoma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has not had disease progression while on therapy	

Product Name: Brand Targretin caps, generic bexarotene caps, Targretin gel	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Targretin will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.	

Product Name: Brand Targretin caps, generic bexarotene caps, Targretin gel	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Targretin therapy</p>	

## 2 . Revision History

Date	Notes
6/2/2021	Arizona Medicaid 7.1 Implementation

Tarpeyo (budesonide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-113527
<b>Guideline Name</b>	Tarpeyo (budesonide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/8/2022
-----------------	----------

### 1 . Criteria

Product Name: Tarpeyo	
Approval Length	9 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of primary immunoglobulin A nephropathy (IgAN) as confirmed by a kidney biopsy  <b>AND</b>	

**2** - Patient is at risk of rapid disease progression [e.g., generally a urine protein-to-creatinine ratio (UPCR) greater than or equal to 1.5 g/g, or by other criteria such as clinical risk scoring using the International IgAN Prediction Tool]

**AND**

**3** - Used to reduce proteinuria

**AND**

**4** - Estimated glomerular filtration rate (eGFR) greater than or equal to 35 mL/min/1.73 m<sup>2</sup>

**AND**

**5** - One of the following:

**5.1** Patient has been on a minimum 90-day trial of a maximally tolerated dose and will continue to receive therapy with one of the following:

- An angiotensin-converting enzyme (ACE) inhibitor (e.g., benazepril, lisinopril)
- An angiotensin II receptor blocker (ARB) (e.g., losartan, valsartan)

**OR**

**5.2** Patient has a contraindication or intolerance to both ACE inhibitors and ARBs

**AND**

**6** - Trial and failure, contraindication, or intolerance to another glucocorticoid (e.g., methylprednisolone, prednisone)

**AND**

**7** - Prescribed by or in consultation with a nephrologist

## 2 . Revision History

Date	Notes
9/8/2022	Removed references, no clinical criteria changes.

Tasigna

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99772
<b>Guideline Name</b>	Tasigna
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Tasigna	
Diagnosis	Chronic Myeloid Leukemia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of chronic myeloid leukemia	

**AND**

**2** - ONE of the following:

**2.1** Patient is not a candidate for imatinib (Gleevec) as attested by physician

**OR**

**2.2** Patient is currently on Tasigna therapy

Product Name:Tasigna	
Diagnosis	Chronic Myeloid Leukemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on Tasigna therapy	

Product Name:Tasigna	
Diagnosis	Gastrointestinal Stromal Tumor (GIST)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of progressive gastrointestinal stromal tumor (GIST)	

**AND**

**2** - History of failure, contraindication, or intolerance to ALL of the following:

- Gleevec (imatinib)
- Sutent (sunitinib)
- Stivarga (regorafenib)

Product Name:Tasigna	
Diagnosis	Gastrointestinal Stromal Tumor (GIST)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient does not show evidence of progressive disease while on Tasigna therapy	

Product Name:Tasigna	
Diagnosis	Acute Lymphoblastic Leukemia (ALL)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ALL)	

Product Name:Tasigna	
Diagnosis	Acute Lymphoblastic Leukemia (ALL)

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Tasigna therapy</p>	

Product Name:Tasigna	
Diagnosis	Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of myeloid/lymphoid neoplasms with eosinophilia and ABL1 (gene) rearrangement</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Neoplasm is in blast or chronic phase</p>	

Product Name:Tasigna	
Diagnosis	Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient does not show evidence of progressive disease while on Tassigna therapy

**Product Name:Tassigna**

Diagnosis	NCCN Recommended Regimens
-----------	---------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Tassigna will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.

**Product Name:Tassigna**

Diagnosis	NCCN Recommended Regimens
-----------	---------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to Tassigna therapy

**2 . Revision History**

Date	Notes
------	-------

6/2/2021	Arizona Medicaid 7.1 Implementation
----------	-------------------------------------

Tecelra (afamitresgene autoleucel)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-157796
<b>Guideline Name</b>	Tecelra (afamitresgene autoleucel)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Tecelra*	
Diagnosis	Synovial Sarcoma
Approval Length	1 Time Authorization in Lifetime
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming ALL of the following: <b>1.1</b> Diagnosis of synovial sarcoma	

**AND**

**1.2** Disease is one of the following:

- Unresectable
- Metastatic

**AND**

**1.3** Both of the following:

**1.3.1** Patient is HLA-A\*02:01P, HLA-A\*02:02P, HLA-A\*02:03P, or HLA-A\*02:06P positive

**AND**

**1.3.2** Patient does not have HLA-A\*02:05P in either allele

**AND**

**1.4** Presence of MAGE-A4 antigen as detected by a U.S. Food and Drug Administration (FDA)-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims confirming patient has received prior therapy with one of the following:

- An anthracycline (e.g. doxorubicin)
- Ifosfamide

**AND**

**3** - Patient is 16 years of age or older

**AND**

4 - Submission of medical records (e.g., chart notes) or paid claims confirming patient has never received Tecelra treatment in their lifetime

Notes

\*For medical PA, Final approval must be approved by the member's plan. If approved, the transaction will go through POS.

## 2 . Revision History

Date	Notes
10/25/2024	New program

Tegsedi

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99652
<b>Guideline Name</b>	Tegsedi
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Tegsedi	
Diagnosis	Hereditary transthyretin-mediated (hATTR) amyloidosis with polyneuropathy
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - BOTH of the following:	

- Diagnosis of Hereditary transthyretin-mediated (hATTR) amyloidosis with polyneuropathy
- Documentation that the patient has a pathogenic transthyretin (TTR) mutation (e.g., V30M)

**AND**

**2** - Prescribed by or in consultation with a neurologist

**AND**

**3** - Documentation of ONE of the following:

- Patient has a baseline polyneuropathy disability (PND) score less than or equal to IIIb
- Patient has a baseline familial amyloidotic polyneuropathy (FAP) Stage 1 or 2
- Patient has a baseline neuropathy impairment (NIS) score greater than or equal to 10 and less than or equal to 130

**AND**

**4** - Patient has not had a liver transplant

**AND**

**5** - Presence of clinical signs and symptoms of the disease (e.g., peripheral sensorimotor polyneuropathy, autonomic neuropathy, motor disability, etc.)

**AND**

**6** - Patient is not receiving Tegsedi in combination with ONE of the following:

- Oligonucleotide agents [e.g., Onpattro (patisiran)]
- Tafamidis (e.g., Vyndaqel, Vyndamax)

Product Name: Tegsedi

Diagnosis	Hereditary transthyretin-mediated (hATTR) amyloidosis with polyneuropathy
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient has previously received treatment with Tegsedi

**AND**

2 - Prescribed by or in consultation with a neurologist

**AND**

3 - Documentation of ONE of the following:

- Patient continues to have a polyneuropathy disability (PND) score less than or equal to IIIb
- Patient continues to have a familial amyloidotic polyneuropathy (FAP) Stage 1 or 2
- Patient continues to have a neuropathy impairment (NIS) score greater than or equal to 10 and less than or equal to 130

**AND**

4 - Documentation that the patient has experienced a positive clinical response to Tegsedi therapy (e.g., improved neurologic impairment, motor function, quality of life, slowing of disease progression, etc.)

**AND**

5 - Patient is not receiving Tegsedi in combination with ONE of the following:

- Oligonucleotide agents [e.g., Onpattro (patisiran)]

- Tafamidis (e.g., Vyndaqel, Vyndamax)

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Tepezza (teprotumumab-trbw)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-127089
<b>Guideline Name</b>	Tepezza (teprotumumab-trbw)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Tepezza	
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting diagnosis of thyroid eye disease (TED)  <b>AND</b>	

**2** - Prescribed by or in consultation with one of the following:

- Endocrinologist
- Specialist with expertise in the treatment of TED
- Ophthalmologist

**AND**

**3** - Treatment with Tepezza has not exceeded a total of 8 infusions

## **2 . Revision History**

Date	Notes
6/26/2023	Removed criteria for TED severity due to expanded indication, added ophthalmologist as prescriber option.

Test Strips

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-120594
<b>Guideline Name</b>	Test Strips
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Non-preferred Test Strip Products	
Approval Length	12 month(s)
Guideline Type	Step Therapy
<b>Approval Criteria</b> 1 - History of failure, contraindication, or intolerance to BOTH of the following*: <ul style="list-style-type: none"><li>True Metrix</li><li>Accu-Chek</li></ul>	

<b>OR</b>	
2 - Patient is on an insulin pump	
<b>OR</b>	
3 - Patient is visually impaired	
Notes	*See background section for plan specific preferred agents

Product Name: Preferred or non-preferred test strip products	
Approval Length	12 month(s)
Guideline Type	Quantity Limit
<p><b>Approval Criteria</b></p> <p>1 - ONE of the following:</p> <p>1.1 For Insulin Dependent or Pregnant patients, the physician must confirm the patient requires a greater quantity because of more frequent blood glucose testing (e.g., patients on intravenous insulin infusions)</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2 For Non-Insulin Dependent Patients, ONE the following:</p> <p>1.2.1 The patient is experiencing or is prone to hypoglycemia or hyperglycemia and requires additional testing to achieve glycemic control</p> <p style="text-align: center;"><b>OR</b></p> <p>1.2.2 The patient's physician is adjusting medications and the patient requires additional blood glucose testing during this time</p>	

**OR**

**1.2.3** The patient's physician is adjusting MNT (medical nutrition therapy) and the patient requires additional blood glucose testing during this time

**OR**

**1.2.4** The patient requires additional testing due to fluctuations in blood glucose due to physical activity or exercise

**OR**

**1.2.5** Other circumstances where prescribing physician confirms that the patient requires a greater quantity because of more frequent blood glucose testing (clinical review required by OptumRx reviewing pharmacist and/or medical director)

## 2 . Background

Benefit/Coverage/Program Information	
<b>Preferred Test Strips According to Plan</b>	
PLAN	PREFERRED TEST STRIPS
Arizona Complete Health	OneTouch Ultra test strips OneTouch Verio test strips
Care1st	OneTouch Ultra test strips OneTouch Verio test strips

MercyCare	OneTouch meters and strips (all OneTouch products)	
Banner University Family Care	Freestyle OneTouch Ultra test strips OneTouch Verio test strips	
Health Choice	Accu-Check products	
Molina	True Metrix	
UHC/C&S AZ	OneTouch Ultra test strips OneTouch Verio test strips	

### 3 . Revision History

Date	Notes
1/27/2023	Updated background chart, no changes to criteria

Testosterone - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242225
<b>Guideline Name</b>	Testosterone - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name: PREFERRED: generic testosterone 1.62% gel (generic for Androgel), Depo-testosterone, testosterone cypionate, testosterone enanthate; NON-PREFERRED: generic testosterone 1% gel/pump, Brand Androgel, Brand Testim, generic testosterone TD gel (generic Testim), Brand Vogelxo gel/pump, generic testosterone 1.62% gel, generic testosterone TD gel/pump (generic Vogelxo), generic testosterone 2% TD gel (generic Fortesta), testosterone topical 30mg/act solution, Azmiro, Jatenzo, Kyzatrex, Natesto, Tlando, Undecatrex	
Diagnosis	Hypogonadism
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:

**1.1** TWO pre-treatment serum total testosterone levels less than 300 ng/dL (less than 10.4 nmol/L) or less than the reference range for the lab, taken at separate times (Document lab value and date for both levels)

**OR**

**1.2** BOTH of the following:

**1.2.1** Patient has a condition that may cause altered sex-hormone binding globulin (SHBG) (e.g., thyroid disorder, HIV disease, liver disorder, diabetes, obesity)

**AND**

**1.2.2** ONE pre-treatment calculated free or bioavailable testosterone level less than 50 pg/mL (less than 5 ng/dL or less than 0.17 nmol/L) or less than the reference range for the lab (This may require treatment to be temporarily held. Document lab value and date)

**OR**

**1.3** Patient has a history of ONE of the following:

- Bilateral orchiectomy
- Panhypopituitarism
- A genetic disorder known to cause hypogonadism (e.g., congenital anorchia, Klinefelter's syndrome)

**AND**

**2** - Patient is NOT taking ONE of the following growth hormones, unless diagnosed with panhypopituitarism:

- Genotropin
- Humatrope
- Norditropin FlexPro
- Nutropin AQ
- Omnitrope

- Saizen

**AND**

**3** - Patient is NOT taking with an Aromatase inhibitor (eg, Arimidex [anastrozole], Femara [letrozole], or Aromasin [exemestane])

**AND**

**4** - Patient was male at birth

**AND**

**5** - Diagnosis of hypogonadism

**AND**

**6** - ONE of the following:

- Significant reduction in weight (less than 90 percent ideal body weight) (e.g., AIDS wasting syndrome)
- Osteopenia
- Osteoporosis
- Decreased bone density
- Decreased libido
- Organic cause of testosterone deficiency (eg, injury, tumor, infection, or genetic defects)

**AND**

**7** - If the request is for a NON-PREFERRED drug: Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to generic testosterone 1.62% gel (generic for Androgel by preferred manufacturers)

Product Name: PREFERRED: generic testosterone 1.62% gel (generic for Androgel), Depo-testosterone, testosterone cypionate, testosterone enanthate; NON-PREFERRED: generic testosterone 1% gel/pump, Brand Androgel, Brand Testim, generic testosterone TD gel

(generic Testim), Brand Vogelxo gel/pump, generic testosterone 1.62% gel, generic testosterone TD gel/pump (generic Vogelxo), generic testosterone 2% TD gel (generic Fortesta), testosterone topical 30mg/act solution, Azmiro, Jatenzo, Kyzatrex, Natesto, Tlando, Undecatrex

Diagnosis	Gender Dysphoria
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Patient is using hormones to change physical characteristics

**AND**

2 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of gender dysphoria, as defined by the current version of the Diagnostic and Statistical Manual of Mental Disorders (DSM)

**AND**

3 - Patient is NOT taking ONE of the following growth hormones, unless diagnosed with panhypopituitarism:

- Genotropin
- Humatrope
- Norditropin FlexPro
- Nutropin AQ
- Omnitrope
- Saizen

**AND**

4 - Patient is NOT taking with an Aromatase inhibitor (eg, Arimidex [anastrozole], Femara [letrozole], Aromasin [exemestane])

**AND**

**5** - If the request is for a NON-PREFERRED drug: Submission of medical records (e.g., chart notes) or paid claims confirming a history of failure or intolerance to generic testosterone 1.62% gel (generic for Androgel by preferred manufacturers)

Product Name: PREFERRED: generic testosterone 1.62% gel (generic for Androgel), Depo-testosterone, testosterone cypionate, testosterone enanthate; NON-PREFERRED: generic testosterone 1% gel/pump, Brand Androgel, Brand Testim, generic testosterone TD gel (generic Testim), Brand Vogelxo gel/pump, generic testosterone 1.62% gel, generic testosterone TD gel/pump (generic Vogelxo), generic testosterone 2% TD gel (generic Fortesta), testosterone topical 30mg/act solution, Azmiro, Jatenzo, Kyzatrex, Natesto, Tlando, Undecatrex

Diagnosis	Hypogonadism, Gender Dysphoria
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting ONE of the following:

**1.1** Follow-up total serum testosterone level drawn within the past 12 months is within or below the normal male limits of the reporting lab (document value and date)

**OR**

**1.2** Follow-up total serum testosterone level drawn within the past 12 months is outside of upper male limits of normal for the reporting lab and the dose is adjusted (document value and date)

**OR**

**1.3** BOTH of the following:

**1.3.1** Patient has a condition that may cause altered sex-hormone binding globulin (SHBG) (e.g., thyroid disorder, HIV disease, liver disorder, diabetes, obesity)

**AND**

**1.3.2** ONE of the following:

**1.3.2.1** Follow-up calculated free or bioavailable testosterone level drawn within the past 12 months is within or below the normal male limits of the reporting lab (document lab value and date)

**OR**

**1.3.2.2** Follow-up calculated free or bioavailable testosterone level drawn within the past 12 months is outside of upper male limits of normal for the reporting lab and the dose is adjusted (document value and date)

**AND**

**2** - Patient is NOT taking ONE of the following growth hormones, unless diagnosed with panhypopituitarism:

- Genotropin
- Humatrope
- Norditropin FlexPro
- Nutropin AQ
- Omnitrope
- Saizen

**AND**

**3** - Patient is NOT taking with an Aromatase inhibitor (eg, Arimidex [anastrozole], Femara [letrozole], Aromasin [exemestane])

**AND**

**4** - If the request is for a NON-PREFERRED drug: Submission of medical records (e.g., chart

notes) or paid claims confirming a history of failure or intolerance to generic testosterone 1.62% gel (generic for Androgel by preferred manufacturers)

## 2 . Revision History

Date	Notes
4/24/2025	Removed inactive products, updated product name sections, updated verbiage in embedded step

Tezruly (terazosin) oral solution

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-241269
<b>Guideline Name</b>	Tezruly (terazosin) oral solution
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name: Tezruly	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Trial and failure, or intolerance to terazosin capsules/tablets  <b>OR</b>  2 - Patient is unable to swallow oral capsules/tablets	

**2 . Revision History**

Date	Notes
4/24/2025	New program

Tezspire (tezepelumab-ekko)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-121766
<b>Guideline Name</b>	Tezspire (tezepelumab-ekko)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2023
-----------------	----------

## 1 . Criteria

Product Name: Tezspire	
Approval Length	6 Month(s) [A]
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of severe asthma	

**AND**

**2** - Patient is 12 years of age or older

**AND**

**3** - One of the following: [2,3]

- Patient has had two or more asthma exacerbations requiring systemic corticosteroids (e.g., prednisone) within the past 12 months
- Prior asthma-related hospitalization within the past 12 months

**AND**

**4** - Patient is currently being treated with one of the following unless there is a contraindication or intolerance to these medications:

**4.1** Both of the following: [2,3]

- High-dose inhaled corticosteroid (ICS) (i.e., greater than 500 mcg fluticasone propionate equivalent/day)
- Additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium)

**OR**

**4.2** One maximally-dosed combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Symbicort [budesonide/formoterol], Breo Ellipta [fluticasone/vilanterol]) [B]

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Allergist/Immunologist

Product Name: Tezspire	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy as evidenced by one of the following:</p> <ul style="list-style-type: none"> <li>• A reduction in asthma exacerbations</li> <li>• Improvement in forced expiratory volume in 1 second (FEV1) from baseline</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications [4]</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Pulmonologist</li> <li>• Allergist/Immunologist</li> </ul>	

## 2 . Endnotes

- A. The Global Initiative for Asthma (GINA) Global Strategy for Asthma Management and Prevention update recommends that patients with asthma should be reviewed regularly to monitor their symptom control, risk factors and occurrence of exacerbations, as well as to document the response to any treatment changes. Ideally, after initiation of treatment, patients should be re-evaluated in 3 to 6 months. [4]
- B. The Global Initiative for Asthma (GINA) Global Strategy for Asthma Management and Prevention guideline recommend patients with severe asthma should be treated with maximal optimized high dose ICS-LABA therapy. [4]

### 3 . Revision History

Date	Notes
2/27/2023	Added new GPI

Thalomid

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99780
<b>Guideline Name</b>	Thalomid
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Thalomid	
Diagnosis	Multiple Myeloma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of multiple myeloma	

Product Name:Thalomid	
Diagnosis	Multiple Myeloma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Thalomid therapy</p>	

Product Name:Thalomid	
Diagnosis	Erythema Nodosum Leprosum (ENL)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of moderate to severe erythema nodosum leprosum (ENL)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <p>2.1 Used for acute treatment</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Used as maintenance therapy for prevention &amp; suppression of cutaneous manifestations of ENL recurrence</p>	

Product Name:Thalomid
-----------------------

Diagnosis	Erythema Nodosum Leprosum (ENL)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Thalomid therapy	

Product Name:Thalomid	
Diagnosis	Aphthous Stomatitis or Ulcer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of severe, recurrent aphthous stomatitis or ulcer	

Product Name:Thalomid	
Diagnosis	Aphthous Stomatitis or Ulcer
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Thalomid therapy	

Product Name:Thalomid	
-----------------------	--

Diagnosis	Pyoderma Gangrenosum
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of pyoderma gangrenosum</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Used as third line treatment</p>	

Product Name:Thalomid	
Diagnosis	Pyoderma Gangrenosum
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Thalomid therapy</p>	

Product Name:Thalomid	
Diagnosis	Cutaneous Manifestations Systemic Lupus Erythematosus (SLE)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Diagnosis of cutaneous manifestations of systemic lupus erythematosus (SLE)

Product Name:Thalomid

Diagnosis	Cutaneous Manifestations Systemic Lupus Erythematosus (SLE)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to Thalomid therapy

Product Name:Thalomid

Diagnosis	B-Cell Lymphomas
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of Castleman's Disease (CD)

**AND**

2 - NOT used as first line therapy

Product Name:Thalomid

Diagnosis	B-Cell Lymphomas
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Thalomid therapy</p>	

Product Name:Thalomid	
Diagnosis	Myelofibrosis-Associated Anemia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of primary myelofibrosis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p>2.1 Both of the following:</p> <p>2.1.1 Serum erythropoietin levels less than 500 mU/mL</p> <p style="text-align: center;"><b>AND</b></p> <p>2.1.2 History of failure, contraindication, or intolerance to erythropoietins [e.g., Procrit (epoetin alfa)]</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Serum erythropoietin levels greater than or equal to 500 mU/mL</p>	

Product Name:Thalomid	
Diagnosis	Myelofibrosis-Associated Anemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation that member has evidence of symptom improvement or reduction in spleen-liver volume while on Thalomid</p>	

Product Name:Thalomid	
Diagnosis	Acquired Immunodeficiency Syndrome (AIDS)- Related Kaposi Sarcoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of Acquired Immunodeficiency Syndrome (AIDS)- Related Kaposi Sarcoma</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is currently being treated with antiretroviral therapy (ART)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Not used as first line therapy</p>	

Product Name:Thalomid	
Diagnosis	AIDS- Related Kaposi Sarcoma

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Thalomid therapy</p>	

Product Name:Thalomid	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Thalomid will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.</p>	

Product Name:Thalomid	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Thalomid therapy</p>	

## 2 . Revision History

Date	Notes
6/3/2021	Arizona Medicaid 7.1 Implementation

Thrombopoiesis Stimulating Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-146016
<b>Guideline Name</b>	Thrombopoiesis Stimulating Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li><li>• Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	5/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Preferred Drugs: Nplate, Promacta tablet	
Diagnosis	Chronic Immune Thrombocytopenia (ITP)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of chronic immune thrombocytopenia (ITP)	

**AND**

**2** - History of failure, contraindication, or intolerance to ONE of the following:

- Corticosteroids
- Immunoglobulins
- Splenectomy

Notes

\*Note: Drugs may require PA

Product Name: Non-Preferred Drugs: Alvaiz, Doptelet, Promacta powder pack/oral suspension, Tavalisse

Diagnosis Chronic Immune Thrombocytopenia (ITP)

Approval Length 12 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

**1** - Diagnosis of chronic immune thrombocytopenia (ITP)

**AND**

**2** - One of the following:

**2.1** Both of the following:

**2.1.1** History of failure, contraindication, or intolerance to ONE of the following:

- Corticosteroids
- Immunoglobulins
- Splenectomy

**AND**

**2.1.2** History of failure, contraindication, or intolerance to BOTH of the following preferred alternatives\*:

- Promacta Tablet (eltrombopag)\*
- Nplate (romiplostim)\*

**OR**

**2.2** Patient is currently stable on requested non-preferred medication

Notes	*Note: Drugs may require PA
-------	-----------------------------

Product Name:Alvaiz, Doptelet, Nplate, Promacta tablets, Promacta powder pack/oral suspension, Tavalisse	
Diagnosis	Chronic Immune (idiopathic) thrombocytopenia (ITP)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

Product Name:Alvaiz, Promacta tablets, Promacta powder pack/oral suspension	
Diagnosis	Severe Aplastic Anemia
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of severe aplastic anemia</p>	

**AND**

**2** - One of the following:

**2.1** Used in combination with standard immunosuppressive therapy [e.g., Atgam (antithymocyte globulin equine), Thymoglobulin (antithymocyte globulin rabbit), cyclosporine]

**OR**

**2.2** History of failure, contraindication, or intolerance to at least one course of immunosuppressive therapy [e.g., Atgam (antithymocyte globulin equine), Thymoglobulin (antithymocyte globulin rabbit), cyclosporine]

**AND**

**3** - For Alvaiz and Promacta powder pack/oral suspension requests ONLY: clinical rationale for use instead of preferred Promacta tablet

Product Name:Alvaiz, Promacta tablets, Promacta powder pack/oral suspension	
Diagnosis	Severe Aplastic Anemia
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	

Product Name:Alvaiz, Promacta tablet	
Diagnosis	Chronic Hepatitis C-associated Thrombocytopenia
Approval Length	6 month(s)
Therapy Stage	Initial Authorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of chronic Hepatitis C-associated thrombocytopenia</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <ul style="list-style-type: none"> <li>• Planning to initiate and maintain interferon-based treatment</li> <li>• Currently receiving interferon-based treatment</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - For Alvaiz requests ONLY: History of failure, contraindication, or intolerance to Promacta tablet</p>	

Product Name:Alvaiz, Promacta tablet	
Diagnosis	Chronic Hepatitis C-associated Thrombocytopenia
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is currently on antiviral interferon therapy for treatment of chronic Hepatitis C</p>	

Product Name:Doptelet, Mulpleta	
Diagnosis	Thrombocytopenia
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of thrombocytopenia</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has chronic liver disease</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is scheduled to undergo a procedure</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - History of failure, contraindication, or intolerance to BOTH of the following preferred alternatives*:</p> <ul style="list-style-type: none"> <li>• Promacta Tablets (eltrombopag)*</li> <li>• Nplate (romiplostim)*</li> </ul>	
Notes	*Note: Drugs may require PA

Product Name:Nplate	
Diagnosis	Hematopoietic Syndrome of Acute Radiation Syndrome [HS-ARS]
Approval Length	6 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of Hematopoietic Syndrome of Acute Radiation Syndrome [HS-ARS]

**AND**

2 - Patient is receiving myelosuppressive doses of radiation

**2 . Revision History**

Date	Notes
4/23/2024	Added Alvaiz as NP target

Tobramycin Inhalation - ARIZONA

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99653
<b>Guideline Name</b>	Tobramycin Inhalation - ARIZONA
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Bethkis, Kitabis	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of cystic fibrosis (CF)	

Product Name:Brand TOBI Nebulizer Solution, generic tobramycin solution for inhalation, TOBI Podhaler	
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of cystic fibrosis (CF)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Lung infection with positive culture demonstrating Pseudomonas aeruginosa infection</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - History of failure, intolerance, or contraindication to BOTH of the following</p> <ul style="list-style-type: none"> <li>• Brand Bethkis</li> <li>• Kitabis</li> </ul>	

Product Name: Brand TOBI Nebulizer Solution, generic tobramycin solution for inhalation, TOBI Podhaler	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

## 2 . Revision History

Date	Notes
------	-------

3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1
-----------	--

Tocilizumab (Actemra, Tofidence, Tyenne)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325195
<b>Guideline Name</b>	Tocilizumab (Actemra, Tofidence, Tyenne)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Preferred: Tyenne SC/IV	
Diagnosis	Rheumatoid Arthritis
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:  1.1 Diagnosis of moderately to severely active Rheumatoid Arthritis (RA)	

**AND**

**1.2** History of failure to a 3 month trial of ONE non-biologic disease modifying anti-rheumatic drug (DMARD) [e.g., methotrexate, leflunomide, sulfasalazine, hydroxychloroquine] at maximally indicated doses within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**2** - Prescribed by or in consultation with a rheumatologist

Product Name: Preferred: Tyenne SC/IV	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:</p> <p><b>1.1</b> Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis</p> <p><b>AND</b></p> <p><b>1.2</b> History of failure to a minimum duration of a 6-week trial, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:</p> <ul style="list-style-type: none"><li>• leflunomide</li><li>• methotrexate</li></ul> <p><b>AND</b></p>	

2 - Prescribed by, or in consultation with, a rheumatologist

Product Name:Preferred: Tyenne SC/IV	
Diagnosis	Systemic Juvenile Idiopathic Arthritis (SJIA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:</p> <p>1.1 Diagnosis of active systemic juvenile idiopathic arthritis</p> <p style="text-align: center;"><b>AND</b></p> <p>1.2 History of failure, contraindication, or intolerance to one of the following conventional therapies at maximally tolerated doses:</p> <ul style="list-style-type: none"><li>• Minimum duration of a 3-month trial and failure of methotrexate</li><li>• Minimum duration of a 1-month trial of nonsteroidal anti-inflammatory drug (NSAID) (e.g., ibuprofen, naproxen)</li><li>• Minimum duration of a 2-week trial of systemic glucocorticoid (e.g., prednisone)</li></ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by, or in consultation with, a rheumatologist</p>	

Product Name:Preferred: Tyenne SC/IV	
Diagnosis	Giant Cell Arteritis (GCA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:

1.1 Diagnosis of giant cell arteritis (GCA)

**AND**

1.2 History of failure, contraindication, or intolerance to a glucocorticoid (e.g., prednisone)

**AND**

2 - Prescribed by or in consultation with a rheumatologist

Product Name:Preferred: Tyenne SC/IV	
Diagnosis	Rheumatoid Arthritis, Polyarticular Juvenile Idiopathic Arthritis (PJIA), Systemic Juvenile Idiopathic Arthritis (SJIA), Giant Cell Arteritis (GCA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by, or in consultation with, a rheumatologist	

Product Name:Preferred: Tyenne SC
-----------------------------------

Diagnosis	Systemic Sclerosis-Associated Interstitial Lung Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting a diagnosis of active systemic sclerosis-associated interstitial lung disease (SSc-ILD) as documented by ALL of the following:

**1.1** ONE of the following:

**1.1.1** Skin thickening of the fingers of both hands extending proximal to the metacarpophalangeal joints

**OR**

**1.1.2** TWO of the following:

- Skin thickening of the fingers (e.g., puffy fingers, sclerodactyly of the fingers)
- Fingertip lesions (e.g., digital tip ulcers, fingertip pitting scars)
- Telangiectasia
- Abnormal nailfold capillaries
- Pulmonary arterial hypertension
- Raynaud's phenomenon
- SSc-related autoantibodies (e.g., anticentromere, anti-topoisomerase I, anti-RNA polymerase III)

**AND**

**1.2** Presence of interstitial lung disease as determined by finding evidence of pulmonary fibrosis on HRCT (high-resolution computed tomography), involving at least 10% of the lungs

**AND**

**2** - Prescribed by, or in consultation with, a pulmonologist

Product Name:Preferred: Tyenne SC	
Diagnosis	Systemic Sclerosis-Associated Interstitial Lung Disease
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by, or in consultation with, a pulmonologist</p>	

Product Name:Preferred: Tyenne IV	
Diagnosis	Coronavirus disease 2019 (COVID-19)
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of COVID-19</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is hospitalized (tocilizumab is only FDA approved when used for COVID 19 patients in an inpatient setting)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Currently receiving systemic corticosteroids</p>	

**AND**

4 - Patient requires one of the following:

- Supplemental oxygen
- Non-invasive mechanical ventilation
- Invasive mechanical ventilation
- Extracorporeal membrane oxygenation (ECMO)

Notes

NOTE: Tocilizumab is only FDA approved when used for COVID-19 patients in an inpatient setting

Product Name:Non-Preferred\*: Actemra SC/IV, Tofidence IV, and newly launched tocilizumab products

Approval Length

Requests for Non-Preferred biosimilars are not approved at this time

Guideline Type

Prior Authorization

### Approval Criteria

1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred product is Tyenne (tocilizumab-aazg). \*\*Please note: The drug(s) listed above may require additional review.

Notes

\*Patients must use preferred tocilizumab biosimilar(s).

## 2 . Revision History

Date	Notes
7/16/2025	Updated preferred agents and clinical criteria, updated NP section verbiage.

Topical Capsaicin Products

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-136966
<b>Guideline Name</b>	Topical Capsaicin Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	12/1/2023
-----------------	-----------

### 1 . Criteria

Product Name:Diclareal	
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming diagnosis of osteoarthritis of the knees</p> <p style="text-align: center;"><b>AND</b></p>	

**2** - Submission of medical records (e.g., chart notes, paid claims history) documenting history of failure to ALL of the following:

- diclofenac 1% topical gel
- diclofenac 2% topical solution
- topical capsaicin cream/patch

Product Name: Trubrex	
Approval Length	3 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) confirming requested medication is being used for the treatment of acute and chronic pain in muscles and joints associated with muscle soreness, strains, sprains, arthritis, simple backache, muscle stiffness, etc</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Submission of medical records (e.g., chart notes, paid claims history) documenting trial and failure, contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"> <li>• diclofenac 1% topical gel</li> <li>• topical capsaicin cream/patch</li> <li>• topical lidocaine patch</li> </ul>	

## 2 . Revision History

Date	Notes
12/1/2023	New program

Topical NSAIDs

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99574
<b>Guideline Name</b>	Topical NSAIDs
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Brand Flector Patch, generic diclofenac epolamine 1.3% patch	
Approval Length	2 Week(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of acute pain due to minor strains, sprains, or contusions  <b>AND</b>  2 - ONE of the following:	

**2.1** The patient did not receive adequate pain relief when treated with at least three preferred non-steroidal anti-inflammatory drugs (NSAIDs) (An inadequate response to treatment is defined as pain and/or inflammatory symptoms not resolved after 14 days of therapy)

- Diclofenac DR (Generic Voltaren)
- Diclofenac ER (Generic Voltaren ER)
- Etodolac (Generic Lodine)
- Etodolac ER (Generic Lodine ER)
- Fenoprofen (Generic Nalfon)
- Flurbiprofen (Generic Ansaid)
- Ibuprofen
- Indomethacin (Generic Indocin)
- Ketorolac (Generic Toradol)
- Mefenamic (Generic Ponstel)
- Meloxicam (Generic Mobic)
- Nabumetone (Generic Relafen)
- Nabumetone DS (Generic Relafen DS)
- Naproxen (Generic Anaprox)
- Naproxen DR (Generic Anaprox DR)
- Naproxen EC (Generic Anaprox EC)
- Oxaprozin (Generic Daypro)
- Piroxicam (Generic Feldene)
- Sulindac (Generic Clinoril)

**OR**

**2.2** The patient has one of the following risk factors for NSAID-induced adverse GI (gastrointestinal) events:

- Patient is greater than or equal to 65 years of age
- Prior history of peptic, gastric, or duodenal ulcer
- History of NSAID-related ulcer
- History of clinically significant GI (gastrointestinal) bleeding
- Untreated or active H. Pylori gastritis
- Concurrent use of oral corticosteroids (e.g. prednisone, prednisolone, dexamethasone)
- Concurrent use of anticoagulants (e.g. warfarin, heparin)
- Concurrent use of antiplatelets (e.g. aspirin including low-dose, clopidogrel)

Product Name:Pennsaid 2%, diclofenac sodium soln 1.5%	
Approval Length	12 month(s)
Guideline Type	Prior Authorization

## **Approval Criteria**

1 - Patient has a diagnosis of pain due to osteoarthritis of the knee(s)

**AND**

2 - ONE of the following:

**2.1** The patient did not receive adequate pain relief when treated with at least three preferred non-steroidal anti-inflammatory drugs (NSAIDs) (An inadequate response to treatment is defined as pain and/or inflammatory symptoms not resolved after 14 days of therapy)

- Diclofenac DR (Generic Voltaren)
- Diclofenac ER (Generic Voltaren ER)
- Etodolac (Generic Lodine)
- Etodolac ER (Generic Lodine ER)
- Fenoprofen (Generic Nalfon)
- Flurbiprofen (Generic Ansaid)
- Ibuprofen
- Indomethacin (Generic Indocin)
- Ketorolac (Generic Toradol)
- Mefenamic (Generic Ponstel)
- Meloxicam (Generic Mobic)
- Nabumetone (Generic Relafen)
- Nabumetone DS (Generic Relafen DS)
- Naproxen (Generic Anaprox)
- Naproxen DR (Generic Anaprox DR)
- Naproxen EC (Generic Anaprox EC)
- Oxaprozin (Generic Daypro)
- Piroxicam (Generic Feldene)
- Sulindac (Generic Clinoril)

**OR**

**2.2** The patient has one of the following risk factors for NSAID-induced adverse GI (gastrointestinal) events:

- Patient is greater than or equal to 65 years of age
- Prior history of peptic, gastric, or duodenal ulcer
- History of NSAID-related ulcer
- History of clinically significant GI bleeding
- Untreated or active H. Pylori gastritis

- Concurrent use of oral corticosteroids (e.g. prednisone, prednisolone, dexamethasone)
- Concurrent use of anticoagulants (e.g. warfarin, heparin)
- Concurrent use of antiplatelets (e.g. aspirin including low-dose, clopidogrel)

**AND**

**3** - Patient has a history of failure, intolerance, or contraindication to diclofenac topical gel 1% (Rx formulation), or Voltaren OTC (over the counter)

**Product Name:** generic diclofenac topical gel 1% (Rx formulation), Voltaren OTC

Approval Length	12 month(s)
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - The patient has a diagnosis of pain due to osteoarthritis of joints amenable to topical treatment, including but not limited to the hands, knees, ankles, elbows, feet, and wrists

**AND**

**2** - ONE of the following:

**2.1** The patient did not receive adequate pain relief when treated with at least three preferred non-steroidal anti-inflammatory drugs (NSAIDs) (An inadequate response to treatment is defined as pain and/or inflammatory symptoms not resolved after 14 days of therapy)

- Diclofenac DR (Generic Voltaren)
- Diclofenac ER (Generic Voltaren ER)
- Etodolac (Generic Lodine)
- Etodolac ER (Generic Lodine ER)
- Fenoprofen (Generic Nalfon)
- Flurbiprofen (Generic Ansaid)
- Ibuprofen
- Indomethacin (Generic Indocin)
- Ketorolac (Generic Toradol)
- Mefenamic (Generic Ponstel)
- Meloxicam (Generic Mobic)
- Nabumetone (Generic Relafen)
- Nabumetone DS (Generic Relafen DS)

- Naproxen (Generic Anaprox)
- Naproxen DR (Generic Anaprox DR)
- Naproxen EC (Generic Anaprox EC)
- Oxaprozin (Generic Daypro)
- Piroxicam (Generic Feldene)
- Sulindac (Generic Clinoril)

**OR**

**2.2** The patient has one of the following risk factors for NSAID-induced adverse GI (gastrointestinal) events:

- Patient is greater than or equal to 65 years of age
- Prior history of peptic, gastric, or duodenal ulcer
- History of NSAID-related ulcer
- History of clinically significant GI bleeding
- Untreated or active H. Pylori gastritis
- Concurrent use of oral corticosteroids (e.g. prednisone, prednisolone, dexamethasone)
- Concurrent use of anticoagulants (e.g. warfarin, heparin)
- Concurrent use of antiplatelets (e.g. aspirin including low-dose, clopidogrel)

Trastuzumab Products

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325196
<b>Guideline Name</b>	Trastuzumab Products
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Preferred: Ogivri	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The drug is being used as indicated by National Comprehensive Cancer Network (NCCN) guidelines with a Category of Evidence and Consensus of 1, 2A, or 2B	

Product Name:Non-Preferred*: Herceptin, Hercessi, Herzuma, Kanjinti, Ontruzant, Trazimera, and newly launched trastuzumab products
--

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred product is Ogivri (trastuzumab-dkst). **Please note: The drug(s) listed above may require additional review.</p>	
Notes	*Patients must use preferred trastuzumab biosimilar(s).

## 2 . Revision History

Date	Notes
7/16/2025	New program, updated NP criteria verbiage.

Tremfya (guselkumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-329258
<b>Guideline Name</b>	Tremfya (guselkumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Tremfya SC 100mg*	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:  1.1 Diagnosis of chronic moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3% body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** History of failure to one of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)

**AND**

**1.4** History of failure of a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.5** History of failure, contraindication, or intolerance to one of the following topical therapies

- Vtama
- Zoryve 0.3% cream

**AND**

**1.6** History of failure, contraindication, or intolerance to ALL of the following preferred biologic products:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- Infliximab
- Otezla (apremilast)
- A preferred ustekinumab biosimilar

<b>AND</b>	
<b>2 - Prescribed by or in consultation with a dermatologist</b>	
Notes	*If patient meets criteria above, please approve at GPI-14

Product Name:Tremfya SC 100mg*	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2 - Prescribed by or in consultation with a dermatologist</b></p>	
Notes	*If patient meets criteria above, please approve at GPI-14

Product Name:Tremfya SC 100mg*	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting ALL of the following:</p>	

**1.1** Diagnosis of active psoriatic arthritis (PsA)

**AND**

**1.2** History of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**1.3** History of failure, contraindication, or intolerance to ALL of the following preferred biologic products:

- A preferred adalimumab biosimilar or Enbrel (etanercept)
- infliximab
- Orencia (abatacept)
- Otezla (apremilast)
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**2** - Prescribed by, or in consultation with, ONE of the following:

- Rheumatologist
- Dermatologist

Notes	*If patient meets criteria above, please approve at GPI-14
-------	--

Product Name:Tremfya SC 100mg*	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with ONE of the following:

- Rheumatologist
- Dermatologist

Notes

\*If patient meets criteria above, please approve at GPI-14

Product Name: Tremfya IV

Diagnosis Crohn's Disease (CD)

Approval Length 3 month(s)

Guideline Type Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's Disease (CD)

**AND**

2 - One of the following:

- Frequent diarrhea and abdominal pain
- At least 10% weight loss
- Complications such as obstruction, fever, abdominal mass
- Abnormal lab values (e.g., C-reactive protein [CRP])
- CD Activity Index (CDAI) greater than 220

**AND**

3 - Submission of medical records (e.g., chart notes) or verification of paid claims history documenting BOTH of the following:

**3.1** Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

- Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)
- 6-mercaptopurine (Purinethol)
- Azathioprine (Imuran)
- Methotrexate (Rheumatrex, Trexall)

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following (document drug, date, and duration of trial):

- A preferred adalimumab biosimilar
- infliximab
- A preferred ustekinumab biosimilar

**AND**

**4** - Prescribed by or in consultation with a gastroenterologist

**AND**

**5** - Will be administered as an intravenous induction dose

Product Name:Tremfya SC	
Diagnosis	Crohn's Disease (CD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active Crohn's disease	

**AND**

**2** - Will be used as a maintenance dose following the intravenous induction doses

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name:Tremfya IV

Diagnosis	Ulcerative Colitis (UC)
-----------	-------------------------

Approval Length	3 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis (UC)

**AND**

**2** - One of the following:

- Greater than 6 stools per day
- Frequent blood in the stools
- Frequent urgency
- Presence of ulcers
- Abnormal lab values (e.g., hemoglobin, erythrocyte sedimentation rate, C-reactive protein)
- Dependent on, or refractory to, corticosteroids

**AND**

**3** - Submission of medical records (e.g., chart notes) or verification of paid claims history documenting BOTH of the following:

**3.1** Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

- 6-mercaptopurine
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)
- Azathioprine
- Corticosteroids (e.g., prednisone)

**AND**

**3.2** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz oral tablet (tofacitinib) (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**4** - Prescribed by or in consultation with a gastroenterologist

**AND**

**5** - Will be administered as an intravenous induction dose

Product Name:Tremfya SC	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis (UC)	

**AND**

**2** - Will be used as a maintenance dose following the intravenous induction doses

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name:Tremfya SC	
Diagnosis	Crohn's Disease (CD), Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:	
<ul style="list-style-type: none"><li>• Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li><li>• Reversal of high fecal output state</li></ul>	
<b>AND</b>	
<b>2</b> - Prescribed by or in consultation with a gastroenterologist	

## 2 . Revision History

Date	Notes
------	-------

7/16/2025	Updated preferred agents/embedded steps, updated criteria throughout. PsA: updated "or" to "and" between 1 and 2
-----------	--

Tretinoin Capsules - ARIZONA

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99513
<b>Guideline Name</b>	Tretinoin Capsules - ARIZONA
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name:Tretinoin capsules	
Diagnosis	Acute Promyelocytic Leukemia (APL)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of acute promyelocytic leukemia	

Product Name:Tretinoin capsules	
Diagnosis	Acute Promyelocytic Leukemia (APL)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to tretinoin capsules</p>	

Product Name:Tretinoin capsules	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Tretinoin capsules will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.</p>	

Product Name:Tretinoin capsules	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Documentation of positive clinical response to tretinoin capsules

## 2 . Revision History

Date	Notes
4/8/2021	7/1 Implementation

Tretinoin Topical

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99591
<b>Guideline Name</b>	Tretinoin Topical
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Retin-A cream and gel*	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Patient is 26 years of age or less</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** Both of the following:

- Patient is greater than 26 years of age
- Diagnosis of acne vulgaris

**AND**

**2** - The patient must have a history of therapeutic failure, contraindication, or intolerance to ALL of the following:

- benzoyl peroxide
- topical clindamycin
- topical erythromycin

Notes	*Only Brand Covered
-------	---------------------

## 2 . Revision History

Date	Notes
10/29/2021	Changed effective date to 12/1/21

Trikafta (elexacaftor/tezacaftor/ivacaftor)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-125904
<b>Guideline Name</b>	Trikafta (elexacaftor/tezacaftor/ivacaftor)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Trikafta (80-40-60 mg) granules packet, Trikafta (100-50-75 mg) granules packet	
Diagnosis	Cystic Fibrosis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of cystic fibrosis (CF)	

**AND**

**2** - Submission of laboratory results documenting that the patient has at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or a mutation in the CFTR gene that is responsive to Trikafta based on in vitro data

**AND**

**3** - Patient is between 2 and 6 years of age

**AND**

**4** - Prescribed by, or in consultation with, a specialist affiliated with a CF care center

Product Name: Trikafta (50-25-37.5 mg) tablet pack, Trikafta (100-50-75 mg) tablet pack

Diagnosis	Cystic Fibrosis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Diagnosis of cystic fibrosis (CF)

**AND**

**2** - Submission of laboratory results documenting that the patient has at least one F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or a mutation in the CFTR gene that is responsive to Trikafta based on in vitro data

**AND**

3 - The patient is 6 years of age or older

**AND**

4 - Prescribed by, or in consultation with, a specialist affiliated with a CF care center

Product Name: Trikafta granules packets, Trikafta tablet packs	
Diagnosis	Cystic Fibrosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Provider attests that the patient has achieved a clinically meaningful response while on Trikafta therapy to ONE of the following:	
<ul style="list-style-type: none"><li>• Lung function as demonstrated by percent predicted expiratory volume in 1 second (ppFEV1)</li><li>• Body mass index (BMI)</li><li>• Pulmonary exacerbations</li><li>• Quality of life as demonstrated by Cystic Fibrosis Questionnaire-Revised (CFQ-R) respiratory domain score</li></ul>	
<b>AND</b>	
2 - Prescribed by, or in consultation with, a specialist affiliated with a cystic fibrosis (CF) care center	

## 2 . Revision History

Date	Notes
5/19/2023	Added criteria for ages 2-6, with new corresponding granules packets.



Triptans - AZM

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269192
<b>Guideline Name</b>	Triptans - AZM
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Brand Amerge, Brand Imitrex tablets, Brand Imitrex injection, generic sumatriptan 6mg PFS, generic almotriptan, brand Maxalt, brand Maxalt MLT, Onzetra Xsail, brand Relpax, Symbravo, Brand Treximet, generic sumatriptan-naproxen, Zembrace, brand Zomig, brand Zomig ZMT, brand Frova, generic frovatriptan, Tosymra	
Diagnosis	Non-preferred products
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of migraine headaches with or without aura	

**AND**

**2** - Patient has a history of failure, contraindication, or intolerance to a trial of at least three preferred products (document drugs, duration, and date of trials)\*

- naratriptan (generic Amerge)
- rizatriptan (generic Maxalt)
- sumatriptan (Generic Imitrex)
- zolmitriptan (Generic Zomig)
- eletriptan (generic Relpax)

**AND**

**3** - For Brand Treximet and generic sumatriptan-naproxen requests ONLY, BOTH of the following (applies to Brand/generic Treximet ONLY):

**3.1** Trial, failure, contraindication or intolerance to BOTH of the following:

- naproxen
- sumatriptan

**AND**

**3.2** Physician has provided rationale for needing to use fixed-dose combination therapy with requested medication instead of taking individual products in combination (i.e., naproxen and sumatriptan)

**AND**

**4** - For Symbravo requests ONLY, BOTH of the following (applies to Symbravo ONLY):

**4.1** Trial, failure, contraindication or intolerance to BOTH of the following:

- meloxicam
- rizatriptan

**AND**

**4.2** Physician has provided rationale for needing to use fixed-dose combination therapy with requested medication instead of taking individual products in combination (i.e., meloxicam and rizatriptan)

Product Name: Brand Imitrex (inj, cartridge, auto-injector and PFS), generic sumatriptan (inj, cartridge, auto-injector and PFS)\*

Diagnosis	Migraine Headaches with or without Aura
Approval Length	12 month(s)
Guideline Type	Quantity Limits

**Approval Criteria**

1 - Diagnosis of migraine headaches with or without aura

**AND**

2 - Prescribed by or in consultation with one of the following:

- Neurologist
- Pain management specialist

**AND**

3 - Patient is currently receiving prophylactic therapy with at least ONE of the following:

3.1 Amitriptyline (Elavil)

**OR**

3.2 One of the following beta-blockers:

- atenolol
- metoprolol
- nadolol\*\*
- propranolol

- timolol\*\*

**OR**

**3.3** Divalproex sodium (Depakote/Depakote ER)

**OR**

**3.4** OnabotulinumtoxinA (Botox) \*\*\*

**OR**

**3.5** Topiramate (Topamax)

**OR**

**3.6** Venlafaxine (Effexor/Effexor XR)

**OR**

**3.7** Calcitonin gene-related peptide (CGRP) receptor antagonists [e.g., Aimovig (erenumab), Emgality (galcanezumab)]

**AND**

**4** - One of the following:

**4.1** Higher dose or quantity is supported in the dosage and administration section of the manufacturer's prescribing information

**OR**

**4.2** Higher dose or quantity is supported by one of the following compendia:

- American Hospital Formulary Service Drug Information
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**OR**

**4.3** Physician provides evidence from published biomedical literature to support safety and additional efficacy at doses/quantities greater than those approved by the Food and Drug Administration (FDA) for the diagnosis indicated

**AND**

**5** - Physician acknowledges that the potential benefit outweighs the risk associated with the higher dose or quantity

Notes

\* See "Quantity Limits" table in background section for quantity limits  
 \*\* Nadolol and timolol are non-preferred and should not be included in denial to provider  
 \*\*\* OnabotulinumtoxinA (Botox) is a medical benefit, should not be included in denial to provider

Product Name: Brand Imitrex (inj, cartridge, auto-injector and PFS), generic sumatriptan (inj, cartridge, auto-injector and PFS)\*

Diagnosis	Cluster Headaches
Approval Length	12 month(s)
Guideline Type	Quantity Limit

**Approval Criteria**

**1** - Diagnosis of cluster headaches

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Neurologist

- Pain management specialist

**AND**

**3** - Patient has experienced at least 2 cluster periods lasting from 7 days to 365 days, separated by pain-free periods lasting at least three months.

**AND**

**4** - One of the following:

**4.1** Higher dose or quantity is supported in the dosage and administration section of the manufacturer's prescribing information

**OR**

**4.2** Higher dose or quantity is supported by one of the following compendia:

- American Hospital Formulary Service Drug Information
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**OR**

**4.3** Physician provides evidence from published biomedical literature to support safety and additional efficacy at doses/quantities greater than those approved by the Food and Drug Administration (FDA) for the diagnosis indicated

**AND**

**5** - Physician acknowledges that the potential benefit outweighs the risk associated with the higher dose or quantity

Notes

\* See "Quantity Limits" table in background section for quantity limits

Product Name: Brand Amerge, generic naratriptan, Brand Frova, generic frovatriptan, Brand Imitrex tablets and nasal spray, generic sumatriptan tablets and nasal spray, generic

almotriptan, Brand Maxalt and Maxalt MLT, generic rizatriptan and rizatriptan MLT, Onzetra Xsail, Brand Relpax, generic eletriptan, Symbravo, Brand Treximet, generic sumatriptan-naproxen, Zembrace Sym Touch, Brand Zomig and Zomig ZMT, generic zolmitriptan and zolmitriptan ZMT, brand Zomig nasal, generic zolmitriptan nasal spray, Tosymra \*

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Quantity Limit
----------------	----------------

### Approval Criteria

1 - Diagnosis of migraine headaches with or without aura

**AND**

2 - Prescribed by or in consultation with one of the following:

- Neurologist
- Pain management specialist

**AND**

3 - Patient is currently receiving prophylactic therapy with at least ONE of the following:

3.1 Amitriptyline (Elavil)

**OR**

3.2 One of the following beta-blockers:

- atenolol
- metoprolol
- nadolol\*\*
- propranolol
- timolol\*\*

**OR**

3.3 Divalproex sodium (Depakote/Depakote ER)

**OR**

**3.4** OnabotulinumtoxinA (Botox) \*\*\*

**OR**

**3.5** Topiramate (Topamax)

**OR**

**3.6** Venlafaxine (Effexor/Effexor XR)

**OR**

**3.7** Calcitonin gene-related peptide (CGRP) receptor antagonists [e.g., Aimovig (erenumab), Emgality (galcanezumab)]

**AND**

**4** - One of the following:

**4.1** Higher dose or quantity is supported in the dosage and administration section of the manufacturer's prescribing information

**OR**

**4.2** Higher dose or quantity is supported by one of the following compendia:

- American Hospital Formulary Service Drug Information
- Thomson Micromedex DrugDex
- Clinical pharmacology
- United States Pharmacopoeia-National Formulary (USP-NF)

**OR**

**4.3** Physician provides evidence from published biomedical literature to support safety and additional efficacy at doses/quantities greater than those approved by the FDA (Food and Drug Administration) for the diagnosis indicated

**AND**

**5** - Physician acknowledges that the potential benefit outweighs the risk associated with the higher dose or quantity

Notes

\* See "Quantity Limits" table in background section for quantity limits  
\*\* Nadolol and timolol are non-preferred and should not be included in denial to provider  
\*\*\* OnabotulinumtoxinA (Botox) is a medical benefit, should not be included in denial to provider

Product Name:generic zolmitriptan nasal spray

Approval Length | 12 month(s)

Guideline Type | Step Therapy

**Approval Criteria**

1 - Patient has a history of failure, contraindication, or intolerance to a trial of Brand Zomig Nasal Spray

**2 . Background**

**Benefit/Coverage/Program Information**

**Quantity Limits**

Drug Name	Strength	Quantity Limit
Brand Amerge	1mg, 2.5mg	9 tabs/month

generic naratriptan		
Brand Frova	2.5mg	9 tabs/month
Generic frovatriptan		
Brand Imitrex tablets	25mg, 50mg, 100mg	9 tabs/month
generic sumatriptan tablets		
Brand Maxalt	5mg, 10mg	9 tabs/month
Generic rizatriptan		
Brand Maxalt MLT	5mg, 10mg	9 tabs/month
Generic rizatriptan ODT		
Generic almotriptan	6.25mg, 12.5mg	6 tabs/month
Relpax	20mg, 40mg	6 tabs/month
Generic eletriptan		
Brand Zomig	2.5mg, 5mg	6 tabs/month
Generic zolmitriptan		
Brand Zomig ZMT	2.5mg, 5mg	6 tabs/month
Generic zolmitriptan ODT		
Brand Imitrex Nasal Spray	5mg, 20mg	6 spray devices/month
Generic sumatriptan nasal spray		
Zomig Nasal Spray	2.5mg, 5mg	6 spray devices/month
Treximet	85mg/500 mg	9 tabs/month
Generic sumatriptan/naproxen		
Symbravo (rizatriptan/meloxicam)	10mg/20mg	9 tabs/month
Onzetra Xsail	11mg	1 box (8 units)/month
Zembrace SymTouch	3mg	1 box (4 units)/month

Brand Imitrex Generic Sumatriptan Autoinjector/Cartridge Refills	4mg/0.5mL 6mg/0.5mL	8 autoinjectors or cartridge refills/month (4 boxes/month)
Brand Imitrex Generic Sumatriptan Vials	6mg/0.5mL	10 vials/month (2 boxes/month)
Generic Sumatriptan Pre-filled Syringe	6mg/0.5mL	8 prefilled syringes (4 boxes/month)
Tosymra nasal spray	10mg	6 units per month

### 3 . Revision History

Date	Notes
5/29/2025	Added Symbravo as NP target, updated criteria.

Tryngolza (olezarsen sodium)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-193190
<b>Guideline Name</b>	Tryngolza (olezarsen sodium)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Tryngolza	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of familial chylomicronemia syndrome (FCS) (type 1 hyperlipoproteinemia)	

**AND**

**2** - Submission of medical records (e.g., chart notes) documenting one of the following:

**2.1** Genetic confirmation of biallelic pathogenic variants in FCS-causing genes (i.e., LPL, GPIHBP1, APOA5, APOC2, or LMF1)

**OR**

**2.2** A North American FCS (NAFCS) Score of greater than or equal to 45

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting both of the following:

**3.1** One of the following:

**3.1.1** Patient has tried or will receive treatment with standard of care triglyceride lowering therapy (i.e., prescription omega-3 fatty acid and a fibrate)

**OR**

**3.1.2** Patient has an intolerance to standard of care triglyceride lowering therapy (i.e., prescription omega-3 fatty acid and a fibrate)

**AND**

**3.2** Baseline fasting triglyceride levels are greater than or equal to 880 mg/dL prior to treatment with requested drug

**AND**

**4** - Requested drug will be used as adjunct to a low-fat diet

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Cardiologist
- Endocrinologist
- Gastroenterologist
- Lipid specialist (lipidologist)

Product Name: Tryngolza

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., reduction in triglyceride levels from baseline)

**2 . Revision History**

Date	Notes
2/25/2025	New program

Tryvio (aprocitentan)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-155841
<b>Guideline Name</b>	Tryvio (aprocitentan)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

### 1 . Criteria

Product Name:Tryvio	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of hypertension  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes) or verification of paid claims confirming patient has not achieved target blood pressure (e.g., systolic blood pressure [SBP] less than 130 mmHg) after treatment with ALL of the following antihypertensive medications from different classes for an adequate duration (minimum 4 weeks each) at a maximally tolerated dose:

**2.1** One of the following:

- Angiotensin converting enzyme (ACE) inhibitor (e.g., captopril, enalapril)
- Angiotensin II receptor blocker (ARB) (e.g., candesartan, valsartan)

**AND**

**2.2** Diuretic (e.g., hydrochlorothiazide, chlorthalidone)

**AND**

**2.3** Calcium channel blocker (e.g., amlodipine, nifedipine)

**AND**

**2.4** Mineralocorticoid receptor antagonist (MRA) [e.g., eplerenone, spironolactone]

**AND**

**3** - Provider attests other causes of hypertension have been excluded (e.g., secondary causes [e.g., primary hyperaldosteronism], white coat effect, medication nonadherence)

**AND**

**4** - Used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community based program)

**AND**

**5** - Requested drug will be used in combination with at least 3 antihypertensive medications from different classes

**AND**

**6** - Prescribed by or in consultation with a specialist experienced in the treatment of resistant hypertension (e.g., cardiologist, nephrologist)

Product Name: Tryvio

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Patient demonstrates positive clinical response to therapy (e.g., systolic blood pressure [SBP] less than 130 mmHg)

**AND**

**2** - Patient continues to use Tryvio in combination with at least 3 antihypertensive medications from different classes and is adherent to therapy

**AND**

**3** - Requested drug will continue to be used as an adjunct to lifestyle modification (e.g., dietary or caloric restriction, exercise, behavioral support, community based program)

**AND**

**4** - Submission of medical records (e.g., chart notes) or verification of paid claims confirming patient has been previously treated with ALL of the following antihypertensive medications from different classes for an adequate duration (minimum 4 weeks each) at a maximally tolerated dose:

**4.1** One of the following:

- Angiotensin converting enzyme (ACE) inhibitor (e.g., captopril, enalapril)
- Angiotensin II receptor blocker (ARB) (e.g., candesartan, valsartan)

**AND**

**4.2** Diuretic (e.g., hydrochlorothiazide, chlorthalidone)

**AND**

**4.3** Calcium channel blocker (e.g., amlodipine, nifedipine)

**AND**

**4.4** Mineralocorticoid receptor antagonist (MRA) [e.g., eplerenone, spironolactone]

## 2 . Revision History

Date	Notes
9/26/2024	New program

Twyneo (tretinoin-benzoyl peroxide 0.1-3% cream)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-107465
<b>Guideline Name</b>	Twyneo (tretinoin-benzoyl peroxide 0.1-3% cream)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2022
-----------------	----------

## 1 . Criteria

Product Name: Twyneo	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting both of the following:  1.1 Both of the following: <ul style="list-style-type: none"><li>Patient is 9 years of age or older</li></ul>	

- Diagnosis of acne vulgaris

**AND**

**1.2** The patient must have a history of therapeutic failure, contraindication, or intolerance to ALL of the following (verified via paid pharmacy claims or submission of medical records):

- benzoyl peroxide
- topical clindamycin
- topical erythromycin
- topical tretinoin (Brand Retin-A)

## 2 . Revision History

Date	Notes
5/24/2022	New program

Tykerb

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99775
<b>Guideline Name</b>	Tykerb
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Tykerb, generic lapatinib	
Diagnosis	Breast Cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 BOTH of the following:	

**1.1.1** Diagnosis of recurrent or stage IV hormone receptor positive, human epidermal growth factor receptor 2-positive (HER2+) breast cancer

**AND**

**1.1.2** Used in combination with an aromatase inhibitor [e.g., Aromasin (exemestane), Femara (letrozole), Arimidex (anastrozole)]

**OR**

**1.2** BOTH of the following:

**1.2.1** Diagnosis of advanced or stage IV human epidermal growth factor receptor 2-positive (HER2+) breast cancer

**AND**

**1.2.2** Used in combination with ONE of the following:

- Herceptin (trastuzumab)
- Xeloda (capecitabine)

Product Name: Brand Tykerb, generic lapatinib	
Diagnosis	Central Nervous System (CNS) Cancers
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - One of the following:	
1.1 ALL of the following:	
1.1.1 Diagnosis of recurrent, central nervous system (CNS) cancer with metastatic lesions	

**AND**

**1.1.2** Tykerb is active against primary (breast) tumor

**AND**

**1.1.3** Used in combination with Xeloda (capecitabine)

**OR**

**1.2** ALL of the following:

**1.2.1** Diagnosis of recurrent intracranial or spinal ependymoma (excluding subependymoma)

**AND**

**1.2.2** Patient has received previous radiation therapy

**AND**

**1.2.3** Patient has received ONE of the following:

- Gross total or subtotal resection
- Localized recurrence
- Evidence of metastasis (brain, spine, or cerebral spinal fluid)

**AND**

**1.2.4** Used in combination with Temodar (temozolomide)

Product Name:Brand Tykerb, generic lapatinib	
Diagnosis	Chordoma
Approval Length	12 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of epidermal growth factor receptor (EGFR) -positive, recurrent chordoma</p>	

Product Name:Brand Tykerb, generic lapatinib	
Diagnosis	Colon Cancer
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of unresectable, advanced or metastatic colon cancer (Human epidermal growth factor receptor 2 (HER2)-amplified and RAS wild type)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient has not previously been treated with a Human epidermal growth factor receptor 2 (HER2) inhibitor [e.g., Kanjinti (trastuzumab), Perjeta (pertuzumab), Nerlynx (neratinib)]</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient has previously been treated with ONE of the following regimens:</p> <ul style="list-style-type: none"> <li>• Oxaliplatin-based therapy without irinotecan</li> <li>• Irinotecan-based therapy without oxaliplatin</li> <li>• FOLFOXIRI (fluorouracil, leucovorin, oxaliplatin, and irinotecan) regimen</li> <li>• A fluoropyrimidine without irinotecan or oxaliplatin</li> </ul> <p style="text-align: center;"><b>AND</b></p>	

4 - Used in combination with trastuzumab

Product Name:Brand Tykerb, generic lapatinib

Diagnosis Rectal Cancer

Approval Length 12 month(s)

Therapy Stage Initial Authorization

Guideline Type Prior Authorization

**Approval Criteria**

1 - Diagnosis of unresectable, advanced or metastatic rectal cancer (Human epidermal growth factor receptor 2 (HER2)-amplified and RAS wild type)

**AND**

2 - Patient has not previously been treated with a Human epidermal growth factor receptor 2 (HER2) inhibitor [e.g., Kanjinti (trastuzumab), Perjeta (pertuzumab), Nerlynx (neratinib)]

**AND**

3 - Patient has previously been treated with ONE of the following regimens:

- Oxaliplatin-based therapy without irinotecan
- Irinotecan-based therapy without oxaliplatin
- FOLFOXIRI (fluorouracil, leucovorin, oxaliplatin, and irinotecan) regimen
- A fluoropyrimidine without irinotecan or oxaliplatin

**AND**

4 - Used in combination with trastuzumab

Product Name:Brand Tykerb, generic lapatinib

Diagnosis Breast Cancer, Central Nervous System (CNS) Cancers, Chordoma, Colon Cancer, Rectal Cancer

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Tykerb therapy</p>	

Product Name:Brand Tykerb, generic lapatinib	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Tykerb will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.</p>	

Product Name:Brand Tykerb, generic lapatinib	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Tykerb therapy</p>	

## 2 . Revision History

Date	Notes
6/2/2021	Arizona Medicaid 7.1 Implementation

Tysabri (natalizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300302
<b>Guideline Name</b>	Tysabri (natalizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Tysabri	
Diagnosis	Multiple Sclerosis (MS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of multiple sclerosis (MS)	

Product Name:Tysabri	
Diagnosis	Multiple Sclerosis (MS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression)</p>	

Product Name:Tysabri	
Diagnosis	Crohn's Disease (CD)
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of moderately to severely active Crohn's disease (CD)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Crohn's disease has evidence of inflammation (e.g., elevated C-reactive protein [CRP], elevated erythrocyte sedimentation rate, presence of fecal leukocytes)</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a gastroenterologist</p> <p style="text-align: center;"><b>AND</b></p>	

**4** - Submission of medical records (e.g., chart notes) or verification of paid claims documenting BOTH of the following:

**4.1** Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

- 6-mercaptopurine
- azathioprine
- Corticosteroids (e.g., prednisone)
- methotrexate

**AND**

**4.2** History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- A preferred ustekinumab biosimilar

Notes	Note: In CD, discontinue Tysabri in patients that have not experienced therapeutic benefit by 12 weeks of induction therapy, and in patients that cannot discontinue chronic concomitant steroids within six months of starting therapy.
-------	--

Product Name:Tysabri	
Diagnosis	Crohn's Disease (CD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:</p> <ul style="list-style-type: none"> <li>• Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li> </ul>	

- Reversal of high fecal output state

**AND**

**2** - Prescribed by or in consultation with a gastroenterologist

## **2 . Revision History**

Date	Notes
7/3/2025	Updated preferred agents/embedded steps for UC

Uloric

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99501
<b>Guideline Name</b>	Uloric
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Uloric, generic febuxostat	
Approval Length	12 month(s)
Guideline Type	Step Therapy
<b>Approval Criteria</b> 1 - History of failure, contraindication or intolerance to allopurinol (generic Zyloprim)	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Ultomiris (ravulizumab-cwvz)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147673
<b>Guideline Name</b>	Ultomiris (ravulizumab-cwvz)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Ultomiris	
Diagnosis	Paroxysmal Nocturnal Hemoglobinuria (PNH)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of paroxysmal nocturnal hemoglobinuria (PNH)	

**AND**

**2** - Patient is one month of age and older

**AND**

**3** - Prescribed by or in consultation with a hematologist/oncologist

Product Name:Ultomiris

Diagnosis	Paroxysmal Nocturnal Hemoglobinuria (PNH)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Documentation of positive clinical response (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) to therapy

Product Name:Ultomiris

Diagnosis	Atypical Hemolytic Uremic Syndrome (aHUS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes) documenting a diagnosis of atypical hemolytic uremic syndrome (aHUS)

**AND**

**2** - Patient is one month of age and older

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Hematologist
- Nephrologist

Product Name:Ultomiris	
Diagnosis	Atypical Hemolytic Uremic Syndrome (aHUS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response (e.g., hemoglobin stabilization, decrease in the number of red blood cell transfusions) to therapy	

Product Name:Ultomiris	
Diagnosis	Generalized Myasthenia Gravis (gMG)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) documenting both of the following:

- Diagnosis of generalized myasthenia gravis (gMG)
- Patient is anti-acetylcholine receptor (AChR) antibody positive

**AND**

2 - Submission of medical records (e.g., chart notes) or paid claims documenting one of the following:

2.1 Trial and failure, contraindication, or intolerance to two preferred immunosuppressive therapies (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus)

**OR**

2.2 Both of the following:

2.2.1 Trial and failure, contraindication, or intolerance to one preferred immunosuppressive therapy (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus)

**AND**

2.2.2 Trial and failure, contraindication, or intolerance to one of the following:

- Chronic plasmapheresis or plasma exchange (PE)
- Intravenous immunoglobulin (IVIG)

**AND**

3 - Prescribed by or in consultation with a neurologist

Product Name:Ultomiris	
Diagnosis	Generalized Myasthenia Gravis (gMG)
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p>	

Product Name:Ultomiris	
Diagnosis	Neuromyelitis Optica Spectrum Disorder (NMOSD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting both of the following:</p> <ul style="list-style-type: none"> <li>• Diagnosis of neuromyelitis optica spectrum disorder (NMOSD)</li> <li>• Patient is anti-aquaporin-4 (AQP4) antibody positive</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Ophthalmologist</li> </ul>	

Product Name:Ultomiris	
Diagnosis	Neuromyelitis Optica Spectrum Disorder (NMOSD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**2 . Revision History**

Date	Notes
5/23/2024	Added criteria for new indication of NMOSD, added submission of records to all initial auth criteria.

Uplizna (inebilizumab-cdon)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-274202
<b>Guideline Name</b>	Uplizna (inebilizumab-cdon)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Uplizna	
Diagnosis	Neuromyelitis Optica Spectrum Disorder (NMOSD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of neuromyelitis optica spectrum disorder (NMOSD)	

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming patient is anti-aquaporin-4 (AQP4) antibody positive

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Neurologist
- Ophthalmologist

**AND**

**4** - One of the following:

**4.1** Submission of medical records (e.g., chart notes) or paid claims confirming history of failure, contraindication, or intolerance to one of the following preferred rituximab products:

- Ruxience
- Truxima

**OR**

**4.2** For continuation of prior therapy

Product Name:Uplizna	
Diagnosis	Immunoglobulin G4-Related Disease (IgG4-RD)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1** - Submission of medical records (e.g., chart notes) confirming a diagnosis of Immunoglobulin G4-Related Disease (IgG4-RD)

**AND**

**2** - Presence of disease involving two or more organ systems or sites (e.g., Pancreas, Submandibular gland, Lymph node(s), Kidneys, Bile Duct, Lungs or Lacrimal glands)

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims confirming ONE of the following:

**3.1** Patient is currently being treated with a glucocorticoid (e.g., prednisone, methylprednisolone)

**OR**

**3.2** Trial and failure, contraindication or intolerance to a glucocorticoid (e.g., prednisone, methylprednisolone)

<b>Product Name:Uplizna</b>	
Diagnosis	All indications listed above
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	

## 2 . Revision History

Date	Notes
5/29/2025	New program

Urea Cycle Disorder Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-128919
<b>Guideline Name</b>	Urea Cycle Disorder Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	8/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Brand Buphenyl, generic sodium phenylbutyrate, Pheburane	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Both of the following:  1.1 Diagnosis of urea cycle disorder (UCD)	

**AND**

**1.2** One of the following deficiencies:

- carbamylphosphate synthetase (CPS)
- ornithine transcarbamylase (OTC)
- argininosuccinic acid synthetase (AS)

**AND**

**2** - Molecular genetic testing confirms mutations in the CPS1, OTC, or ASS1 gene [2]

**AND**

**3** - Trial and failure, or intolerance to generic sodium phenylbutyrate (applies to Brand Buphenyl and Pheburane only)

**AND**

**4** - Used as an adjunct with dietary protein restriction and, in some cases, dietary supplements (e.g., essential amino acids, arginine, citrulline, protein-free calorie supplements)

**AND**

**5** - Prescribed by or in consultation with a specialist focused on the treatment of metabolic disorders

Product Name: Olpruva, Ravicti

Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## Approval Criteria

1 - Both of the following:

1.1 Diagnosis of urea cycle disorder (UCD)

**AND**

1.2 One of the following deficiencies:

- carbamylphosphate synthetase (CPS)
- ornithine transcarbamylase (OTC)
- argininosuccinic acid synthetase (AS)

**AND**

2 - Molecular genetic testing confirms mutations in the CPS1, OTC, or ASS1 gene [2]

**AND**

3 - Inadequate response to one of the following:

- Dietary protein restriction
- Amino acid supplementation

**AND**

4 - Trial and failure, contraindication, or intolerance to generic sodium phenylbutyrate

**AND**

5 - Used as an adjunct with dietary protein restriction and, in some cases, dietary supplements (e.g., essential amino acids, arginine, citrulline, protein-free calorie supplements)

**AND**

**6** - Prescribed by or in consultation with a specialist focused on the treatment of metabolic disorders

Product Name: Brand Buphenyl, generic sodium phenylbutyrate, Olpruva, Pheburane, Ravicti	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., plasma ammonia and amino acid levels within normal limits)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Used as an adjunct with dietary protein restriction and, in some cases, dietary supplements (e.g., essential amino acids, arginine, citrulline, protein-free calorie supplements)</p>	

## 2 . Revision History

Date	Notes
7/28/2023	Added Olpruva as NP target

Ustekinumab

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-325197
<b>Guideline Name</b>	Ustekinumab
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Preferred: Yesintek SC	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, laboratory values, prescription claims history) documenting ALL of the following:  1.1 Diagnosis of chronic moderate to severe plaque psoriasis (PsO)	

**AND**

**1.2** Greater than or equal to 3% body surface area involvement, palmoplantar, facial, or genital involvement, or severe scalp psoriasis

**AND**

**1.3** History of failure to ONE of the following topical therapies, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., betamethasone, clobetasol, desonide)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Vtama
- Zoryve 0.3% cream

**AND**

**1.4** Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**2** - Patient is 6 years of age or older

**AND**

**3** - Prescribed by or in consultation with a dermatologist

Product Name: Preferred: Yesintek SC	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a dermatologist</p>	

Product Name: Preferred: Yesintek SC	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of active psoriatic arthritis (PsA)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is 6 years of age or older</p> <p style="text-align: center;"><b>AND</b></p>	

**4** - Prescribed by or in consultation with one of the following:

- Rheumatologist
- Dermatologist

Product Name:Preferred: Yesintek SC	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to therapy	
<b>AND</b>	
2 - Prescribed by or in consultation with one of the following:	
<ul style="list-style-type: none"><li>• Rheumatologist</li><li>• Dermatologist</li></ul>	

Product Name:Preferred: Yesintek SC/IV	
Diagnosis	Crohn's Disease
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active Crohn's disease (CD)	

**AND**

**2** - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:

- Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)
- Azathioprine (Imuran)
- 6-mercaptopurine (Purinethol)
- Methotrexate (Rheumatrex, Trexall)

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name: Preferred: Yesintek SC/IV	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Diagnosis of moderately to severely active ulcerative colitis (UC)</p> <p><b>AND</b></p> <p><b>2</b> - Paid claims or submission of medical records (e.g., chart notes) documenting history of failure to ONE of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:</p> <ul style="list-style-type: none"><li>• Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)</li><li>• 6-mercaptopurine (Purinethol)</li><li>• Azathioprine (Imuran)</li><li>• Aminosalicylates (e.g., mesalamine, sulfasalazine)</li></ul>	

**AND**

**3** - Prescribed by or in consultation with a gastroenterologist

Product Name:Preferred: Yesintek SC/IV

Diagnosis	Crohn's Disease (CD), Ulcerative Colitis (UC)
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**AND**

**2** - Prescribed by or in consultation with a gastroenterologist

Product Name:Non-Preferred\*: Ustekinumab SC/IV, Ustekinumab-aekn SC/IV, Ustekinumab-ttwe SC/IV, Imuldosa SC/IV, Otulfi SC/IV, Pyzchiva SC/IV, Selarsdi SC/IV, Stelara SC/IV, Steqeyma SC/IV, Wezlana SC/IV, and newly launched ustekinumab products

Approval Length	Requests for Non-Preferred biosimilars are not approved at this time
-----------------	--

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Per your health plan's criteria, the non-preferred drug is not approved for coverage because the plan's preferred product is Yesintek (ustekinumab-kfce). \*\*Please note: The drug(s) listed above may require additional review.

Notes	*Patients must use preferred ustekinumab biosimilar(s).
-------	---

## 2 . Revision History

Date	Notes
7/16/2025	Yesintek to Preferred, updated embedded steps, updated criteria throughout, updated NP section verbiage.

Vafseo (vadadustat)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-155844
<b>Guideline Name</b>	Vafseo (vadadustat)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Vafseo	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) or paid claims confirming ALL of the following:  1.1 Diagnosis of chronic kidney disease (CKD)	

**AND**

**1.2** Patient has been on dialysis for at least 3 months

**AND**

**1.3** Adequate iron stores confirmed by both of the following:

- Patient's ferritin level is greater than 100mcg/L
- Patient's transferrin saturation (TSAT) is greater than 20%

**AND**

**1.4** Hemoglobin level less than 11 g/dL

**AND**

**1.5** Trial and failure, contraindication or intolerance to one of the following preferred erythropoietin stimulating agents (ESA):

- Epogen (epoetin alfa)
- Retacrit (epoetin alfa-epbx)

**AND**

**2** - Prescribed by or in consultation with one of the following:

- hematologist
- nephrologist

**AND**

**3** - Patient is not on concurrent treatment with an erythropoietin stimulating agent [ESA] (e.g., Aranesp, Epogen, Procrit)

Product Name:Vafseo	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) or paid claims confirming ALL of the following:</p> <p><b>1.1</b> Patient demonstrates positive clinical response to therapy (e.g., increase in hemoglobin)</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.2</b> Hemoglobin level does not exceed 11g/dL</p> <p style="text-align: center;"><b>AND</b></p> <p><b>1.3</b> Adequate iron stores confirmed by both of the following:</p> <ul style="list-style-type: none"> <li>• Patient's ferritin level is greater than 100mcg/L</li> <li>• Patient's transferrin saturation (TSAT) is greater than 20%</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>1.4</b> Trial and failure, contraindication or intolerance to one of the following:</p> <ul style="list-style-type: none"> <li>• Epogen (epoetin alfa)</li> <li>• Retacrit (epoetin alfa-epbx)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is not on concurrent treatment with an erythropoietin stimulating agent [ESA] (e.g., Aranesp, Epogen, Procrit)</p>	

## 2 . Revision History

Date	Notes
9/26/2024	New program

Valchlor

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99693
<b>Guideline Name</b>	Valchlor
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name:Valchlor	
Diagnosis	Primary Cutaneous Lymphomas
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of ONE of the following:	

- Chronic or smoldering T-cell leukemia-lymphoma
- Primary cutaneous marginal zone or follicle center B-cell lymphoma
- Lymphomatoid papulosis (LyP) with extensive lesions
- Mycosis fungoides (MF)-Sezary syndrome (SS)

Product Name:Valchlor	
Diagnosis	Primary Cutaneous Lymphomas
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Valchlor</p>	

Product Name:Valchlor	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Valchlor will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.</p>	

Product Name:Valchlor	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Valchlor therapy</p>	

## 2 . Revision History

Date	Notes
4/8/2021	7/1 Implementation

Valsartan oral solution

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114467
<b>Guideline Name</b>	Valsartan oral solution
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Valsartan oral solution	
Diagnosis	Patients 7 years of age or older
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Patient is 7 years of age or older  <b>AND</b>	

2 - Patient cannot take solid dosage form due to swallowing issues

## 2 . Revision History

Date	Notes
9/26/2022	New program

Vecamyl

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99655
<b>Guideline Name</b>	Vecamyl
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Vecamyl	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderately severe to severe essential hypertension  <b>OR</b>	

2 - Diagnosis of uncomplicated malignant hypertension

Product Name:Vecamyl	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of a positive clinical response to Vecamyl therapy	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Velphoro (sucroferric oxyhydroxide), Auryxia (ferric citrate)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-242217
<b>Guideline Name</b>	Velphoro (sucroferric oxyhydroxide), Auryxia (ferric citrate)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Velphoro, Brand Auryxia, Brand ferric citrate tablet	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following <ul style="list-style-type: none"><li>Diagnosis of hyperphosphatemia</li><li>Diagnosis of End Stage Renal Disease</li></ul>	

**AND**

**2** - One of the following:

- For Velphoro requests: Patient is 9 years of age or older
- For Brand Auryxia/Brand ferric citrate requests: Patient is 18 years of age or older

**AND**

**3** - Adherence to and trial and failure to one of the following at maximum dosages (MUST be verified via paid pharmacy claims or submission of medical records)

- Sevelamer Carbonate at the maximum dosage – 800mg/15 per day
- Sevelamer Powder Packets at maximum dosage – 2.4gm packet 4 per day

**AND**

**4** - For Brand ferric citrate tablet (generic for Auryxia) requests ONLY: history of failure or intolerance to Brand Auryxia

Notes	1. Approval will not be granted for requests based on potential side effects, i.e., constipation 2. Approval will not be granted for submitted prior authorizations based on pill burden. Velphoro and Sevelamer are both taken 3 times a day.
-------	---

## 2 . Revision History

Date	Notes
4/24/2025	Added new Brand (MSC M) ferric citrate tablet, generic for Auryxia as NP; added step through Brand Auryxia.

Velsipity (etrasimod)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-300304
<b>Guideline Name</b>	Velsipity (etrasimod)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Velsipity	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of moderately to severely active ulcerative colitis (UC)	

**AND**

**2** - One of the following:

- Greater than 6 stools per day
- Frequent blood in the stools
- Frequent urgency
- Presence of ulcers
- Abnormal lab values (e.g., hemoglobin, ESR, CRP)
- Dependent on, or refractory to, corticosteroids

**AND**

**3** - Paid claims or submission of medical records (e.g., chart notes) confirming a trial and failure, contraindication, or intolerance to one of the following conventional therapies:

- 6-mercaptopurine
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)
- Azathioprine
- Corticosteroids (e.g., prednisone)

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) confirming history of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

**5** - Prescribed by or in consultation with a gastroenterologist

Product Name:Velsipity

Diagnosis

Ulcerative Colitis (UC)

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient demonstrates positive clinical response to therapy as evidenced by at least one of the following:</p> <ul style="list-style-type: none"> <li>• Improvement in intestinal inflammation (e.g., mucosal healing, improvement of lab values [platelet counts, erythrocyte sedimentation rate, C-reactive protein level]) from baseline</li> <li>• Reversal of high fecal output state</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a gastroenterologist</p>	

## 2 . Revision History

Date	Notes
7/3/2025	Updated preferred agents/embedded steps, updated criteria.

Veltassa

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-136978
<b>Guideline Name</b>	Veltassa
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Veltassa	
Diagnosis	Non-Life Threatening Hyperkalemia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of non-life threatening hyperkalemia	

**AND**

**2** - Patient is 12 years of age or older

**AND**

**3** - Where clinically appropriate, medications known to cause hyperkalemia (e.g. angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker, aldosterone antagonist, non-steroidal anti-inflammatory drugs [NSAIDs]) have been discontinued or reduced to the lowest effective dose

**AND**

**4** - Where clinically appropriate, loop or thiazide diuretic therapy for potassium removal has failed

**AND**

**5** - Patient follows a low potassium diet (less than or equal to 3 grams per day)

**AND**

**6** - History of failure, intolerance, or contraindication to Lokelma

Product Name:Veltassa	
Diagnosis	Non-Life Threatening Hyperkalemia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Patient has a positive clinical response to Veltassa therapy

**AND**

2 - Patient continues to require treatment for hyperkalemia

**AND**

3 - Where clinically appropriate, medications known to cause hyperkalemia (e.g. angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker, aldosterone antagonist, non-steroidal anti-inflammatory drugs [NSAIDs])) have been discontinued or reduced to the lowest effective dose

## 2 . Revision History

Date	Notes
9/26/2024	Added new GPI, added age criterion due to expanded approval

Vemlidy

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-146005
<b>Guideline Name</b>	Vemlidy
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Vemlidy	
Diagnosis	Treatment-Naïve Chronic Hepatitis B Infection
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Patient has a contraindication to entecavir therapy  <b>AND</b>	

**2 - Both of the following:**

- Patient is 6 years of age or older
- Patient weighs at least 25 kg

**Product Name:Vemlidy**

Diagnosis	Treatment-Experienced Chronic Hepatitis B Infection
-----------	---

Approval Length	12 month(s)
-----------------	-------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1 - One of the following:**

**1.1 ALL of the following:**

**1.1.1 Patient is currently on Viread therapy**

**AND**

**1.1.2 ONE of the following:**

- Patient has a creatinine clearance less than 60 mL per minute
- Patient has a diagnosis of osteoporosis

**AND**

**1.1.3 Both of the following:**

- Patient is 6 years of age or older
- Patient weighs at least 25 kg

**OR**

**1.2 Patient is currently on Vemlidy therapy**

## 2 . Revision History

Date	Notes
4/22/2024	Updated age/weight criterion due to expanded indication

Veopoz (pozelimab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-135321
<b>Guideline Name</b>	Veopoz (pozelimab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Veopoz	
Diagnosis	CD55-deficient protein-losing enteropathy (PLE)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting ALL of the following:  1.1 Diagnosis of active CD55-deficient protein-losing enteropathy (PLE), also known as CHAPLE disease	

**AND**

**1.2** Patient has a confirmed genotype of biallelic CD55 loss-of-function mutation

**AND**

**1.3** Patient is 1 year of age or older

**AND**

**1.4** Patient has hypoalbuminemia (serum albumin concentration of  $\leq 3.2$  g/dL)

**AND**

**1.5** Patient has at least one of the following signs or symptoms within the last six months:

- abdominal pain
- diarrhea
- peripheral edema
- facial edema

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Immunologist
- Geneticist
- Hematologist

Product Name: Veopoz	
Diagnosis	CD55-deficient protein-losing enteropathy (PLE)
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g. decrease in albumin transfusions and hospitalizations, normalization of serum IgG concentrations, etc.)</p>	

## 2 . Revision History

Date	Notes
10/23/2023	New program

Veozah (fezolinetant)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-128985
<b>Guideline Name</b>	Veozah (fezolinetant)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2023
-----------------	----------

### 1 . Criteria

Product Name:Veozah	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of moderate to severe vasomotor symptoms due to menopause  <b>AND</b>	

**2** - Submission of medical records (e.g., chart notes, paid claims history) documenting trial and failure, contraindication, or intolerance to both of the following (document drug, date, and duration of trial):

- Menopausal hormone therapy (e.g., Premarin, Bijuva, Estrogel, etc.)
- Non-hormonal therapy (e.g. paroxetine mesylate, venlafaxine, clonidine, etc.)

Product Name: Veozah	
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy (e.g., decrease in frequency and severity of vasomotor symptoms from baseline, etc.)</p>	

## 2 . Revision History

Date	Notes
7/26/2023	New program

Verkazia (cyclosporine ophthalmic emulsion 0.1%)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-107454
<b>Guideline Name</b>	Verkazia (cyclosporine ophthalmic emulsion 0.1%)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Verkazia	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting all of the following:  1.1 Diagnosis of moderate to severe vernal keratoconjunctivitis confirmed by the presence of	

clinical signs and symptoms (e.g., itching, photophobia, giant papillae at the upper tarsal conjunctiva or at the limbus, thick mucus discharge, conjunctival hyperaemia)

**AND**

**1.2** Trial and failure, contraindication, or intolerance to one of the following (verified via pharmacy paid claims or submission of medical records):

- Topical ophthalmic “dual-acting” mast cell stabilizer and antihistamine (e.g., olopatadine, azelastine)
- Topical ophthalmic mast cell stabilizers (e.g., cromolyn)

**AND**

**1.3** Trial and failure, contraindication, or intolerance, for short term use (up to 2 to 3 weeks), of topical ophthalmic corticosteroids (e.g., dexamethasone, prednisolone, fluorometholone) ((verified via pharmacy paid claims or submission of medical records)

**AND**

**2** - Prescribed by or in consultation with ONE of the following:

- Ophthalmologist
- Optometrist

Product Name: Verkazia

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy as evidenced by an improvement in clinical signs and symptoms (e.g., itching, photophobia, papillary hypertrophy, mucus discharge, conjunctival hyperaemia)

## 2 . Revision History

Date	Notes
5/24/2022	New program

Vijoice (alpelisib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-148802
<b>Guideline Name</b>	Vijoice (alpelisib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	7/1/2024
-----------------	----------

### 1 . Criteria

Product Name:Vijoice tablets, Vijoice granules	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of PIK3CA-Related Overgrowth Spectrum (PROS)  <b>AND</b>	

2 - Submission of documentation of mutation in the PIK3CA gene

**AND**

3 - Patient is 2 years of age or older

**AND**

4 - Submission of documentation of severe clinical manifestations (e.g., Congenital Lipomatous Overgrowth, Vascular malformations, Epidermal nevi, Scoliosis/skeletal and spinal [CLOVES], Facial Infiltrating Lipomatosis [FIL], Klippel-Trenaunay Syndrome [KTS], Megalencephaly-Capillary Malformation Polymicrogyria [MCAP])

**AND**

5 - Prescribed by or in consultation with a physician who specializes in the treatment of PROS

Product Name:Vijoice tablets, Vijoice granules	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of documentation of positive clinical response to therapy (e.g., radiological response defined as a $\geq$ 20% reduction from baseline in the sum of target lesion volume)	
<b>AND</b>	
2 - Prescribed by or in consultation with a physician who specializes in the treatment of PROS	

## 2 . Revision History

Date	Notes
6/27/2024	Added new granule formulation

Vitamin B-12

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99535
<b>Guideline Name</b>	Vitamin B-12
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Vitamin B-12	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Provider has submitted lab work documenting a Vitamin B-12 deficiency.	

## 2 . Revision History

Date	Notes
5/20/2021	Arizona Medicaid 7.1 Implementation

Vitamin C

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99532
<b>Guideline Name</b>	Vitamin C
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Vitamin C	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Provider has submitted lab work documenting a Vitamin C deficiency	

## 2 . Revision History

Date	Notes
5/19/2021	7/1 Implementation

Vitamin D

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99533
<b>Guideline Name</b>	Vitamin D
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Vitamin D	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Provider has submitted lab work documenting a Vitamin D deficiency	

## 2 . Revision History

Date	Notes
5/19/2021	7/1 Implementation

Vivjoa (oteseconazole)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-228335
<b>Guideline Name</b>	Vivjoa (oteseconazole)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name: Vivjoa	
Approval Length	4 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of recurrent vulvovaginal candidiasis (RVVC)  <b>AND</b>  2 - Patient is NOT of reproductive potential	

**AND**

**3** - Diagnosis of RVVC confirmed by one of the following:

- Positive potassium hydroxide (KOH) preparation
- Vaginal fungal culture

**AND**

**4** - Patient has experienced 3 or more symptomatic episodes of vulvovaginal candidiasis (VVC) within the past 12 months

**AND**

**5** - Trial and failure, contraindication, or intolerance to one intravaginal product (e.g., clotrimazole, miconazole, tioconazole, terconazole, boric acid)

**AND**

**6** - One of the following:

- Failure of a maintenance course of oral fluconazole defined as 100-mg, 150-mg, or 200-mg taken weekly for 6 months or Inadequate response (defined as greater than or equal to 24 weeks of therapy)
- Recurrence of infection while on maintenance therapy, adverse reaction or contraindication to oral fluconazole

**AND**

**7** - Trial and failure, contraindication or intolerance to Brexafemme (ibrexafungerp)

## 2 . Revision History

Date	Notes
------	-------

4/24/2025	Updated oral fluconazole step, added step through Brexafemme
-----------	--

Vonjo (pacritinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-107466
<b>Guideline Name</b>	Vonjo (pacritinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2022
-----------------	----------

## 1 . Criteria

Product Name:Vonjo	
Diagnosis	Myelofibrosis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ALL of the following:  1.1 Diagnosis of ONE of the following:	

- Primary myelofibrosis
- Post-polycythemia vera myelofibrosis
- Post-essential thrombocythemia myelofibrosis

**AND**

**1.2** Disease is intermediate or high risk

**AND**

**1.3** Pre-treatment platelet count below  $50 \times 10^9$  L

**AND**

**2** - Prescribed by or in consultation with **ONE** of the following:

- Hematologist
- Oncologist

Product Name: Vonjo	
Diagnosis	Myelofibrosis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (e.g., symptom improvement, spleen volume reduction)</p>	

Product Name: Vonjo	
Diagnosis	NCCN Recommended Regimens

Approval Length	12 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - This drug will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B</p>	

## 2 . Revision History

Date	Notes
5/24/2022	New Program

Vonoprazan Containing Agents

--

### Prior Authorization Guideline

<b>Guideline ID</b>	GL-155420
<b>Guideline Name</b>	Vonoprazan Containing Agents
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

**Guideline Note:**

Effective Date:	10/1/2024
-----------------	-----------

### 1 . Criteria

Product Name:Voquezna Dual Pak, Voquezna Triple Pak	
Diagnosis	Helicobacter pylori (H. pylori) Infection
Approval Length	1 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of Helicobacter pylori infection</p> <p style="text-align: center;"><b>AND</b></p>	

**2** - Trial and failure, contraindication, or intolerance to BOTH of the following first line treatment regimens:

- Clarithromycin based therapy (e.g., clarithromycin based triple therapy, clarithromycin based concomitant therapy) [D]
- Bismuth quadruple therapy (e.g., bismuth and metronidazole and tetracycline and proton pump inhibitor [PPI] )

Product Name:Voquezna 20mg tablet

Diagnosis	Helicobacter pylori (H. pylori) Infection
-----------	---

Approval Length	1 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

**1** - Diagnosis of Helicobacter pylori infection

**AND**

**2** - One of the following:

- Used in combination with amoxicillin and clarithromycin for the treatment of H. pylori infection
- Used in combination with amoxicillin for the treatment of H. pylori infection

**AND**

**3** - Trial and failure, contraindication, or intolerance to BOTH of the following first line treatment regimens:

- Clarithromycin based therapy (e.g., clarithromycin based triple therapy, clarithromycin based concomitant therapy) [D]
- Bismuth quadruple therapy (e.g., bismuth and metronidazole and tetracycline and proton pump inhibitor [PPI] )

Product Name:Voquezna 20mg tablet	
Diagnosis	Healing and Relief of Heartburn associated with Erosive Esophagitis
Approval Length	8 Week(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of erosive esophagitis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Used for healing of all grades of erosive esophagitis and relief of heartburn associated with erosive esophagitis</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Trial (of a minimum 8-week supply) and inadequate response (within the last 365 days), contraindication, or intolerance to TWO of the following generic proton pump inhibitors (PPI's):</p> <ul style="list-style-type: none"> <li>• omeprazole</li> <li>• esomeprazole</li> <li>• pantoprazole</li> <li>• lansoprazole</li> <li>• rabeprazole</li> <li>• dexlansoprazole</li> </ul>	

Product Name:Voquezna 10mg tablet	
Diagnosis	Maintenance of Healing and Relief of Heartburn associated with Erosive Esophagitis
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Used to maintain healing and relief of heartburn associated with erosive esophagitis

**AND**

2 - Trial (of a minimum 8-week supply) and inadequate response (within the last 365 days), contraindication, or intolerance to TWO of the following generic proton pump inhibitors (PPI's):

- omeprazole
- esomeprazole
- pantoprazole
- lansoprazole
- rabeprazole
- dexlansoprazole

Product Name:Voquezna 10mg tablet\*

Diagnosis	Relief of Heartburn associated with Non-Erosive Gastroesophageal Reflux Disease
-----------	---

Approval Length	1 month(s)
-----------------	------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of non-erosive Gastroesophageal Reflux Disease

**AND**

2 - Both of the following:

- Patient has history of heartburn for at least 6 months
- Heartburn symptoms are present for at least 4 days during any consecutive 7-day period

**AND**

**3** - Trial (of a minimum 8-week supply) and inadequate response (within the last 365 days), contraindication, or intolerance to TWO of the following generic proton pump inhibitors (PPI's):

- omeprazole
- esomeprazole
- pantoprazole
- lansoprazole
- rabeprazole
- dexlansoprazole

Notes

\*Voquezna 20 mg tablets are not FDA approved for non-erosive gastr oesophageal reflux disease.

## 2 . Revision History

Date	Notes
9/26/2024	Added criteria for new indication

Voxzogo (vosoritide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-137865
<b>Guideline Name</b>	Voxzogo (vosoritide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/15/2023
-----------------	------------

## 1 . Criteria

Product Name:Voxzogo	
Diagnosis	Achondroplasia
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Patient has open epiphyses	

**AND**

**2** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of achondroplasia as confirmed by one of the following: [2, 3]

**2.1** Both of the following:

**2.1.1** Patient has clinical manifestations characteristic of achondroplasia (e.g., macrocephaly, frontal bossing, midface retrusion, disproportionate short stature with rhizomelic shortening of the arms and the legs, brachydactyly, trident configuration of the hands, thoracolumbar kyphosis, and accentuated lumbar lordosis)

**AND**

**2.1.2** Patient has radiographic findings characteristic of achondroplasia (e.g., large calvaria and narrowing of the foramen magnum region, undertubulated, shortened long bones with metaphyseal abnormalities, narrowing of the interpedicular distance of the caudal spine, square ilia and horizontal acetabula, small sacrosciatic notches, proximal scooping of the femoral metaphyses, and short and narrow chest)

**OR**

**2.2** Molecular genetic testing confirmed c.1138G>A or c.1138G>C variant (i.e., p.Gly380Arg mutation) in the fibroblast growth factor receptor-3 (FGFR3) gene

**AND**

**3** - Patient did not have limb-lengthening surgery in the previous 18 months and does not plan on having limb-lengthening surgery while on Voxzogo therapy

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Clinical geneticist
- Endocrinologist
- A physician who has specialized expertise in the management of achondroplasia

Product Name:Voxzogo	
Diagnosis	Achondroplasia
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient continues to have open epiphyses</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy as evidenced by one of the following:</p> <ul style="list-style-type: none"> <li>• Improvement in annualized growth velocity (AGV) compared to baseline</li> <li>• Improvement in height Z-score compared to baseline</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Clinical geneticist</li> <li>• Endocrinologist</li> <li>• A physician who has specialized expertise in the management of achondroplasia</li> </ul>	

Product Name:Voxzogo	
Diagnosis	Idiopathic Short Stature (ISS)
Approval Length	N/A - Requests for non-approvable diagnoses should not be approved
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Requests for coverage for diagnosis of Idiopathic Short Stature (ISS) are not authorized and will not be approved

Notes

Approval Length: N/A - Requests for Idiopathic Short Stature (ISS) should not be approved. Deny as a benefit exclusion.

**2 . Revision History**

Date	Notes
12/15/2023	Updated effective date

Vtama (tapinarof)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-210208
<b>Guideline Name</b>	Vtama (tapinarof)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	3/6/2025
-----------------	----------

## 1 . Criteria

Product Name:Vtama	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes, lab work, imaging, paid claims history) documenting a diagnosis of plaque psoriasis	

**AND**

**2** - Submission of medical records (e.g., chart notes, lab work, imaging, paid claims history) documenting a minimum duration of a 4 week trial and failure, contraindication, or intolerance to TWO of the following topical therapies:

- Corticosteroids (e.g., betamethasone, clobetasol)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Anthralin
- Coal tar

**AND**

**3** - Prescribed by or in consultation with a dermatologist

Product Name:Vtama	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging, paid claims history) documenting positive clinical response to therapy as evidenced by one of the following:	
<ul style="list-style-type: none"><li>• Reduction in the body surface area (BSA) involvement from baseline</li><li>• Improvement in symptoms (e.g., pruritus, inflammation) from baseline</li></ul>	

Product Name:Vtama	
Diagnosis	Atopic Dermatitis (AD)
Approval Length	6 month(s)

Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting a diagnosis of atopic dermatitis</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is 2 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Submission of medical records (e.g., chart notes) or paid claims documenting confirming a trial and failure of a minimum 30-day supply (14-day supply for topical corticosteroids), contraindication, or intolerance to ALL of the following:</p> <ul style="list-style-type: none"> <li>• Medium or higher potency topical corticosteroid</li> <li>• Generic topical calcineurin inhibitor (e.g., tacrolimus ointment)</li> <li>• Eucrisa (crisaborole) ointment</li> <li>• Opzelura (ruxolitinib) cream</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Dermatologist</li> <li>• Allergist</li> <li>• Immunologist</li> </ul>	

Product Name:Vtama	
Diagnosis	Atopic Dermatitis (AD)
Approval Length	12 month(s)
Therapy Stage	Reauthorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes, lab work, imaging, paid claims history) documenting positive clinical response to therapy as evidenced by one of the following:</p> <ul style="list-style-type: none"> <li>• Reduction in the body surface area (BSA) involvement from baseline</li> <li>• Reduction in pruritus severity from baseline</li> <li>• Improvement in quality of life from baseline</li> </ul>	

## 2 . Background

Clinical Practice Guidelines			
Table 1: Relative potencies of topical corticosteroids			
Class	Drug	Dosage Form	Strength (%)
Very high potency	Augmented betamethasone dipropionate	Ointment	0.05
	Clobetasol propionate	Cream, foam, ointment	0.05
	Diflorasone diacetate	Ointment	0.05
	Halobetasol propionate	Cream, ointment	0.05
High Potency	Amcinonide	Cream, lotion, ointment	0.1
	Augmented betamethasone dipropionate	Cream	0.05
	Betamethasone dipropionate	Cream, foam, ointment, solution	0.05
	Desoximetasone	Cream, ointment	0.25
	Desoximetasone	Gel	0.05
	Diflorasone diacetate	Cream	0.05
	Fluocinonide	Cream, gel, ointment, solution	0.05
	Halcinonide	Cream, ointment	0.1
	Mometasone furoate	Ointment	0.1
Triamcinolone acetonide	Cream, ointment	0.5	

Medium potency	Betamethasone valerate	Cream, foam, lotion, ointment	0.1
	Clocortolone pivalate	Cream	0.1
	Desoximetasone	Cream	0.05
	Fluocinolone acetonide	Cream, ointment	0.025
	Flurandrenolide	Cream, ointment	0.05
	Fluticasone propionate	Cream	0.05
	Fluticasone propionate	Ointment	0.005
	Mometasone furoate	Cream	0.1
	Triamcinolone acetonide	Cream, ointment	0.1
Lower-medium potency	Hydrocortisone butyrate	Cream, ointment, solution	0.1
	Hydrocortisone probutate	Cream	0.1
	Hydrocortisone valerate	Cream, ointment	0.2
	Prednicarbate	Cream	0.1
Low potency	Alclometasone dipropionate	Cream, ointment	0.05
	Desonide	Cream, gel, foam, ointment	0.05
	Fluocinolone acetonide	Cream, solution	0.01
Lowest potency	Dexamethasone	Cream	0.1
	Hydrocortisone	Cream, lotion, ointment, solution	0.25, 0.5, 1
	Hydrocortisone acetate	Cream, ointment	0.5-1

### 3 . Revision History

Date	Notes
3/6/2025	Added table for Relative potencies of topical corticosteroids to background section, no change to clinical criteria/intent.

Vyjuvek (beremagene geperpavec-svdt)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-131923
<b>Guideline Name</b>	Vyjuvek (beremagene geperpavec-svdt)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Vyjuvek	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of dystrophic epidermolysis bullosa (DEB)	

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming patient has mutation(s) in the collagen type VII alpha 1 chain (COL7A1) gene

**AND**

**3** - Medication is being used for the treatment of wounds

**AND**

**4** - Patient is 6 months of age or older

**AND**

**5** - Medication will be applied by a healthcare professional

**AND**

**6** - Submission of medical records (e.g., chart notes) confirming wound(s) being treated meet all of the following criteria [2]:

- Adequate granulation tissue
- Excellent vascularization
- No evidence of active wound infection in the wound being treated
- No evidence or history of squamous cell carcinoma in the wound being treated

**AND**

**7** - Prescribed by or in consultation with a dermatologist

Product Name:Vyjuvek

Approval Length

6 month(s)

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting positive clinical response (e.g., decrease in wound size, increase in granulation tissue, complete wound closure)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Wound(s) being treated meet all of the following criteria [2]:</p> <ul style="list-style-type: none"> <li>• Adequate granulation tissue</li> <li>• Excellent vascularization</li> <li>• No evidence of active wound infection in the wound being treated</li> <li>• No evidence or history of squamous cell carcinoma in the wound being treated</li> </ul>	

## 2 . Revision History

Date	Notes
8/29/2023	New program

Vykat XR (diazoxide choline)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-274201
<b>Guideline Name</b>	Vykat XR (diazoxide choline)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Vykat XR	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) confirming a diagnosis of Prader-Willi syndrome (PWS)	

**AND**

**2** - Medication is being used for hyperphagia due to PWS

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting disease is confirmed by the presence of a mutation in chromosome 15 as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)

**AND**

**4** - Patient is 4 years of age or older

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Endocrinologist
- Geneticist
- Specialist knowledgeable in the treatment of Prader-Willi syndrome

Product Name:Vykat XR	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Submission of medical records (e.g., chart notes) demonstrating positive clinical response to therapy (e.g., decreased hunger or thoughts about food, decreased weight or BMI)	

## 2 . Revision History

Date	Notes
5/20/2025	New program

Vyndaqel and Vyndamax

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99867
<b>Guideline Name</b>	Vyndaqel and Vyndamax
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Vyndaqel, Vyndamax	
Diagnosis	Transthyretin (ATTR)-mediated amyloidosis with cardiomyopathy (ATTR-CM)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of transthyretin (ATTR)-mediated amyloidosis with cardiomyopathy (ATTR-CM)	

**AND**

**2** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting ONE of the following:

**2.1** Documentation that the patient has a pathogenic transthyretin (TTR) mutation (e.g., V30M)

**OR**

**2.2** Cardiac or noncardiac tissue biopsy demonstrating histologic confirmation of ATTR amyloid deposits

**OR**

**2.3** Submission of medical records (e.g., chart notes, lab work, imaging) documenting ALL of the following

**2.3.1** Echocardiogram or cardiac magnetic resonance imaging suggestive of amyloidosis

**AND**

**2.3.2** Radionuclide imaging (99mTc-DPD, 99mTc-PYP, or 99m Tc-HMDP) showing grade 2 or 3 cardiac uptake\*

**AND**

**2.3.3** Absence of monoclonal protein identified in serum, urine immunofixation (IFE), serum free light chain (sFLC) assay

**AND**

**3** - Prescribed by, or in consultation, with a cardiologist

**AND**

**4** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting presence of clinical signs and symptoms of cardiomyopathy (e.g., heart failure, dyspnea, edema, hepatomegaly, ascites, angina, etc.)

**AND**

**5** - Submission of medical records (e.g., chart notes, lab work, imaging) documenting BOTH of the following:

**5.1** ONE of the following:

**5.1.1** Patient has New York Heart Association (NYHA) Functional Class I or II heart failure

**OR**

**5.1.2** BOTH of the following:

**5.1.2.1** Patient has New York Heart Association (NYHA) Functional Class III heart failure

**AND**

**5.1.2.2** Patient's cardiopulmonary functional status allows patient to ambulate 100 meters or greater in six minutes or less

**AND**

**5.2** Patient has an N-terminal pro-B-type natriuretic peptide (NT-proBNP) level greater than or equal to 600 picograms/milliliter

**AND**

**6** - One of the following:

**6.1** Paid claims or submission of medical records (e.g., chart notes) verifying patient is not receiving Vyndaqel or Vyndamax in combination with either of the following:

- Onpattro (patisiran)
- Tegsedi ( inotersen)

**OR**

**6.2** If the patient is receiving Vyndaqel/Vyndamax in combination with Onpattro (patisiran) or Tegsedi (inotersen), the physician attests that he/she will coordinate care with other specialist(s) involved in the patient’s amyloidosis treatment plan to determine optimal long term monotherapy\*\* treatment regimen

Notes	NOTE: *May require prior authorization and notification ** Referring to monotherapy with Vyndaqel/Vyndamax, Onpattro, or Tegsedi
-------	---

Product Name:Vyndaqel, Vyndamax	
Diagnosis	Transthyretin (ATTR)-mediated amyloidosis with cardiomyopathy (ATTR-CM)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting that the patient has experienced a positive clinical response to Vyndaqel or Vyndamax (e.g., improved symptoms, quality of life, slowing of disease progression, decreased hospitalizations, etc.)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a cardiologist</p> <p style="text-align: center;"><b>AND</b></p>	

**3** - Submission of medical records (e.g., chart notes) documenting that patient continues to have New York Heart Association (NYHA) Functional Class I, II, or III heart failure

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) verifying patient is not receiving Vyndaqel or Vyndamax in combination with either of the following:

- Onpattro (patisiran)
- Tegsedi (inotersen)

## **2 . Revision History**

Date	Notes
12/9/2021	Added submission of records/paid claims where applicable.

Vyvgart (efgartigimod alfa-fcab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-269198
<b>Guideline Name</b>	Vyvgart (efgartigimod alfa-fcab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Vyvgart, Vyvgart Hytrulo	
Diagnosis	Generalized myasthenia gravis (gMG)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting diagnosis of generalized myasthenia gravis (gMG)	

**AND**

**2** - Patient is anti-acetylcholine receptor (AChR) antibody positive

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims confirming BOTH of the following:

**3.1** Trial and failure, contraindication, or intolerance to two immunosuppressive therapies (e.g., glucocorticoids, azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus)

**AND**

**3.2** Trial and failure, contraindication, or intolerance to one of the following:

- Chronic plasmapheresis or plasma exchange (PE)
- Intravenous immunoglobulin (IVIG) or immunoglobulin (IG) therapy

**AND**

**4** - For Vyvgart Hytrulo, submission of medical records (e.g., chart notes) or paid claims confirming trial and failure or intolerance to Vyvgart IV infusion

**AND**

**5** - Prescribed by or in consultation with a neurologist

Product Name:Vyvgart, Vyvgart Hytrulo	
Diagnosis	Generalized myasthenia gravis (gMG)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy

Product Name:Vyvgart Hytrulo	
Diagnosis	Chronic inflammatory demyelinating polyneuropathy (CIDP)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Submission of medical records (e.g., chart notes) or paid pharmacy claims documenting ALL of the following:

1.1 Diagnosis of chronic inflammatory demyelinating polyneuropathy (CIDP) as confirmed by ALL of the following:

1.1.1 Progressive symptoms present for at least 2 months

**AND**

1.1.2 Symptomatic polyradiculoneuropathy as indicated by ONE of the following:

- Progressive or relapsing motor impairment of more than one limb
- Progressive or relapsing sensory impairment of more than one limb

**AND**

1.1.3 Electrophysiologic findings when THREE of the following four criteria are present:

- Partial conduction block of 1 or more motor nerve
- Reduced conduction velocity of 2 or more motor nerves
- Prolonged distal latency of 2 or more motor nerves

- Prolonged F-wave latencies of 2 or more motor nerves or the absence of F waves

**AND**

**1.2** Trial and failure, contraindication, or intolerance to ONE of the following standard of care treatments:

- Corticosteroids (minimum 3 month trial duration)
- Immunoglobulin
- Plasma exchange

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Immunologist
- Neurologist
- Hematologist

Product Name:Vyvgart Hytrulo	
Diagnosis	Chronic inflammatory demyelinating polyneuropathy (CIDP)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes, lab work, imaging) documenting positive clinical response to therapy (e.g., Improvement in INCAT or aINCAT score)</p> <p><b>AND</b></p> <p><b>2</b> - Prescribed by or in consultation one of the following:</p> <ul style="list-style-type: none"> <li>• Immunologist</li> </ul>	

- Neurologist
- Hematologist

## 2 . Revision History

Date	Notes
5/29/2025	Added new GPI for Vyvgart Hytrulo

Wainua (eplontersen)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-144885
<b>Guideline Name</b>	Wainua (eplontersen)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	4/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Wainua	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of hereditary transthyretin-mediated amyloidosis (hATTR amyloidosis) with polyneuropathy	

**AND**

**2** - Submission of medical records (e.g., chart notes) confirming patient has a transthyretin (TTR) mutation (e.g., V30M)

**AND**

**3** - Submission of medical records (e.g., chart notes) confirming one of the following:

- Patient has a baseline familial amyloidotic polyneuropathy (FAP) stage of 1 or 2
- Patient has a baseline neuropathy impairment score (NIS) greater than or equal to 10 and less than or equal to 130
- Patient has a baseline Karnofsky Performance Status score greater than 50%

**AND**

**4** - Presence of clinical signs and symptoms of the disease (e.g., neuropathy, quality of life)

**AND**

**5** - Patient has not had a liver transplant

**AND**

**6** - Prescribed by or in consultation with a neurologist

Product Name:Wainua	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart note) documenting a positive clinical response to therapy as evidenced by an improvement in clinical signs and symptoms from baseline (e.g., neuropathy, quality of life, lower serum TTR level)

**AND**

2 - Submission of medical records (e.g., chart notes) confirming one of the following:

- Patient continues to have a familial amyloidotic polyneuropathy (FAP) stage of 1 or 2
- Patient continues to have a neuropathy impairment score (NIS) greater than or equal to 10 and less than or equal to 130
- Patient continues to have a Karnofsky Performance Status score greater than 50%

**AND**

3 - Patient has not had a liver transplant

## 2 . Revision History

Date	Notes
3/26/2024	New program

Wakix (pitolisant)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152654
<b>Guideline Name</b>	Wakix (pitolisant)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Wakix	
Diagnosis	Narcolepsy with Cataplexy (i.e., Narcolepsy Type 1)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g. chart notes) documenting ALL of the following:  1.1 Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible)	

**AND**

**1.2** Symptoms of cataplexy are present

**AND**

**1.3** Symptoms of excessive daytime sleepiness (e.g., irrepressible need to sleep or daytime lapses into sleep) are present

**AND**

**2** - Physician attestation to the following: Other causes of sleepiness have been ruled out or treated (including but not limited to obstructive sleep apnea, insufficient sleep syndrome, shift work, the effects of substances or medications or their withdrawal, sleep phase disorder, or other sleep disorders)

**AND**

**3** - Prescribed by one of the following:

- Neurologist
- Psychiatrist
- Sleep Medicine Specialist

Product Name:Wakix	
Diagnosis	Narcolepsy without cataplexy (Narcolepsy Type 2)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

**1** - Submission of medical records (e.g. chart notes) documenting ALL of the following:

**1.1** Diagnosis of narcolepsy as confirmed by sleep study (unless the prescriber provides justification confirming that a sleep study would not be feasible)

**AND**

**1.2** Symptoms of cataplexy are absent

**AND**

**1.3** Symptoms of excessive daytime sleepiness (e.g., irrepressible need to sleep or daytime lapses into sleep) are present

**AND**

**2** - Physician attestation to the following: Other causes of sleepiness have been ruled out or treated (including but not limited to obstructive sleep apnea, insufficient sleep syndrome, shift work, the effects of substances or medications or their withdrawal, sleep phase disorder, or other sleep disorders)

**AND**

**3** - Patient is 6 years of age or older

**AND**

**4** - Paid claims or submission of medical records (e.g., chart notes) confirming BOTH of the following:

**4.1** Trial and failure, contraindication (e.g., age), or intolerance to BOTH of the following:

- armodafinil or modafinil
- Sunosi (solriamfetol)

**AND**

**4.2** Trial and failure, contraindication, or intolerance to an amphetamine (e.g., amphetamine, dextroamphetamine) or methylphenidate based stimulant UNLESS patient is not a candidate

**AND**

**5** - Prescribed by one of the following:

- Neurologist
- Psychiatrist
- Sleep Medicine Specialist

Product Name:Wakix	
Diagnosis	All Indications Listed Above
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient has a reduction in symptoms of excessive daytime sleepiness associated with therapy	

## 2 . Revision History

Date	Notes
8/27/2024	Updated criteria, added age criterion.

Wegovy (semaglutide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152738
<b>Guideline Name</b>	Wegovy (semaglutide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Wegovy	
Diagnosis	Reduction of risk of major adverse cardiovascular events
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Treatment is being requested to reduce the risk of major adverse cardiovascular events	

**AND**

**2** - Patient is 45 years of age or older

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting all the following:

**3.1** BMI (body mass index) greater than or equal to 30 kg/m<sup>2</sup> (kilograms per square meter)

**AND**

**3.2** Established cardiovascular disease as evidenced by one of the following:

**3.2.1** Prior myocardial infarction (MI)

**OR**

**3.2.2** One of the following diagnoses/diagnosis codes:

- I25.2 Old myocardial infarction
- I46.2 Cardiac arrest due to underlying cardiac condition
- I46.8 Cardiac arrest due to other underlying condition
- I46.9 Cardiac arrest, cause unspecified
- Z95.0 Presence of cardiac pacemaker
- Z95.1 Presence of aortocoronary bypass graft
- Z95.2 Presence of prosthetic heart valve
- Z95.3 Presence of xenogenic heart valve
- Z95.4 Presence of other heart-valve replacement
- Z95.5 Presence of coronary angioplasty implant and graft

**AND**

**4** - Used in combination with a reduced calorie diet and increased physical activity

**AND**

**5** - For patients with a history of myocardial infarction, submission of medical records (e.g., chart notes) or paid claims documenting one of the following:

**5.1** Patient is on therapy from each of the following classes (as confirmed by claims history or submission of medical records):

- Cholesterol lowering medication (e.g., statin, PCSK9i)
- Beta blocker (i.e., carvedilol, metoprolol, or bisoprolol)
- Angiotensin-converting enzyme inhibitor (ACE-I)/angiotensin II receptor blocker (ARB)/angiotensin II receptor blocker neprilysin inhibitor (ARNI)
- Antiplatelet (e.g., aspirin, clopidogrel)

**OR**

**5.2** Patient has a history of intolerance or contraindication to all of the following therapeutic classes (please specify intolerance or contraindication):

- Cholesterol lowering medication (e.g., statin, PCSK9i)
- Beta blocker (i.e., carvedilol, metoprolol, or bisoprolol)
- Angiotensin-converting enzyme inhibitor (ACE-I)/angiotensin II receptor blocker (ARB)/angiotensin II receptor blocker neprilysin inhibitor (ARNI)
- Antiplatelet (e.g., aspirin, clopidogrel)

**AND**

**6** - Patient does NOT have diagnosis of diabetes or HgA1c greater than or equal to 6.5%

**AND**

**7** - Patient does NOT have New York Heart Association class IV heart failure

**AND**

**8** - Patient does NOT have end-stage renal disease

Product Name:Wegovy

Diagnosis

Reduction of risk of major adverse cardiovascular events

Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - BMI (body mass index) greater than or equal to 30 kg/m<sup>2</sup> (kilograms per square meter)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Used in combination with a reduced calorie diet and increased physical activity</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient does NOT have diagnosis of diabetes or HgA1c greater than or equal to 6.5%</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Patient does NOT have New York Heart Association class IV heart failure</p> <p style="text-align: center;"><b>AND</b></p> <p>5 - Patient does NOT have end-stage renal disease</p>	

Product Name:Wegovy	
Diagnosis	Weight Loss
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Wegovy when used solely for the treatment of weight loss is excluded and is to be denied as a benefit exclusion</p>	

## 2 . Revision History

Date	Notes
8/29/2024	New program

Xalkori

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99695
<b>Guideline Name</b>	Xalkori
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name:Xalkori	
Diagnosis	Inflammatory Myofibroblastic Tumor (IMT)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Diagnosis of inflammatory myofibroblastic tumor (IMT) with anaplastic lymphoma kinase (ALK) translocation

Product Name:Xalkori	
Diagnosis	Non-Small Cell Lung Cancer (NSCLC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of non-small cell lung cancer (NSCLC)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Disease is ONE of the following:</p> <ul style="list-style-type: none"> <li>• Metastatic</li> <li>• Recurrent</li> <li>• Advanced</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Tumor is anaplastic lymphoma kinase (ALK)-positive</li> <li>• Tumor is ROS1-positive</li> <li>• Tumor is positive for mesenchymal-epithelial transition (MET) amplification</li> <li>• Tumor is positive for MET exon 14 skipping mutation</li> </ul>	

Product Name:Xalkori	
Diagnosis	Central Nervous System (CNS) Cancers
Approval Length	12 month(s)
Therapy Stage	Initial Authorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of metastatic brain cancer from non-small cell lung cancer (NSCLC)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - ONE of the following:</p> <ul style="list-style-type: none"> <li>• Tumor is anaplastic lymphoma kinase (ALK)-positive</li> <li>• Tumor is ROS1-positive</li> </ul>	

Product Name: Xalkori	
Diagnosis	Anaplastic Large Cell Lymphoma
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of anaplastic large cell lymphoma</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Tumor is anaplastic lymphoma kinase (ALK)-positive</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Disease is relapsed or refractory</p>	

Product Name:Xalkori	
Diagnosis	Inflammatory Myofibroblastic Tumor (IMT), Non-Small Cell Lung Cancer (NSCLC), Central Nervous System (CNS) Cancers, Anaplastic Large Cell Lymphoma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Patient does not show evidence of progressive disease while on Xalkori therapy</p>	

Product Name:Xalkori	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Xalkori will be approved for uses supported by The National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium with a Category of Evidence and Consensus of 1, 2A, or 2B.</p>	

Product Name:Xalkori	
Diagnosis	NCCN Recommended Regimens
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Documentation of positive clinical response to Xalkori therapy

## 2 . Revision History

Date	Notes
4/8/2021	7/1 Implementation

Xdemvy (lotilaner)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-285207
<b>Guideline Name</b>	Xdemvy (lotilaner)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Xdemvy	
Approval Length	2 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting ALL of the following:</p> <p>1.1 Diagnosis of Demodex blepharitis</p> <p style="text-align: center;"><b>AND</b></p>	

**1.2** Patient exhibits one of the following signs of Demodex infestation

- Collarettes
- Eyelid margin erythema
- Eyelash anomalies (e.g., eyelash misdirection)

**AND**

**1.3** Patient is experiencing symptoms or architectural changes associated with Demodex infestation (e.g., burning, tearing, itching, foreign body sensation, eyelashes missing, eyelashes growing inward)

**AND**

**1.4** Inadequate response to at least a 12-week trial of BOTH of the following:

- tea tree-oil
- eyelid scrub

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Ophthalmologist
- Optometrist

## 2 . Revision History

Date	Notes
5/30/2025	Updated embedded step

Xeljanz, Xeljanz XR (tofacitinib)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-152802
<b>Guideline Name</b>	Xeljanz, Xeljanz XR (tofacitinib)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2024
-----------------	-----------

## 1 . Criteria

Product Name:Xeljanz tablets or Xeljanz XR tablets	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of moderately to severely active rheumatoid arthritis (RA)	

**AND**

**2** - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

**3** - Prescribed by or in consultation with a rheumatologist

Product Name: Xeljanz tablets or Xeljanz XR tablets	
Diagnosis	Rheumatoid Arthritis (RA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Documentation of positive clinical response to therapy	
<b>AND</b>	
<b>2</b> - Prescribed by or in consultation with a rheumatologist	

Product Name: Xeljanz tablets or Xeljanz XR tablets	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of active psoriatic arthritis

**AND**

2 - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to a 3 month trial of methotrexate at the maximally indicated dose within the last 6 months, unless contraindicated or clinically significant adverse effects are experienced

**AND**

3 - Prescribed by or in consultation with one of the following:

- Rheumatologist
- Dermatologist

Product Name: Xeljanz tablets or Xeljanz XR tablets	
Diagnosis	Psoriatic Arthritis (PsA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p><b>AND</b></p> <p>2 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"><li>• Rheumatologist</li></ul>	

- Dermatologist

Product Name: Xeljanz tablets or Xeljanz XR tablets	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of moderately to severely active ulcerative colitis (UC)</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Submission of medical records (e.g., chart notes) or paid claims documenting history of failure to one of the following conventional therapies at maximally indicated doses within the last 3 months, unless contraindicated or clinically significant adverse effects are experienced:</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., prednisone, methylprednisolone, budesonide)</li> <li>• 6-mercaptopurine (Purinethol)</li> <li>• Azathioprine (Imuran)</li> <li>• Aminosalicylates (e.g., mesalamine, sulfasalazine)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with a gastroenterologist</p>	

Product Name: Xeljanz tablets or Xeljanz XR tablets	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with a gastroenterologist

Product Name: Xeljanz tablets or Xeljanz XR tablets

Diagnosis	Ankylosing Spondylitis (AS)
-----------	-----------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Diagnosis of active ankylosing spondylitis

**AND**

2 - Trial and failure, contraindication, or intolerance to TWO nonsteroidal anti-inflammatory drugs (NSAIDs) (e.g., ibuprofen, naproxen)

**AND**

3 - Prescribed by or in consultation with a rheumatologist

Product Name: Xeljanz tablets or Xeljanz XR tablets

Diagnosis	Ankylosing Spondylitis (AS)
-----------	-----------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p>	

Product Name: Xeljanz tablets and oral solution	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of active polyarticular course juvenile idiopathic arthritis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - One of the following:</p> <p>    <b>3.1</b> Both of the following:</p> <p>        <b>3.1.1</b> Submission of medical records (e.g., chart notes) or paid claims documenting trial and failure, contraindication, or intolerance to one of the following nonbiologic DMARDs:</p>	

- leflunomide
- methotrexate

**AND**

**3.1.2** Submission of medical records (e.g., chart notes) or paid claims documenting history of failure, contraindication, or intolerance to all of the following (applies to oral solution ONLY):

- A preferred adalimumab biosimilar
- Enbrel (etanercept)
- Xeljanz or Xeljanz XR (tofacitinib) tablets
- Orencia (abatacept)

**OR**

**3.2** Submission of medical records (e.g., chart notes) or paid claims documenting patient is currently on the requested therapy

Product Name: Xeljanz tablets and oral solution	
Diagnosis	Polyarticular Juvenile Idiopathic Arthritis (PJIA)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Prescribed by or in consultation with a rheumatologist</p>	

## 2 . Revision History

Date	Notes
9/24/2024	Xeljanz XR now preferred, removed embedded step/bypass for XR formulation

Xenazine

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99657
<b>Guideline Name</b>	Xenazine
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Brand Xenazine, generic tetrabenazine	
Diagnosis	Chorea associated with Huntington's Disease
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of chorea in patients with Huntington's disease	

Product Name:Brand Xenazine, generic tetrabenazine
--

Diagnosis	Tardive Dyskinesia (Off Label)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Diagnosis of tardive dyskinesia</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - One of the following:</p> <p>2.1 Patient has persistent symptoms of tardive dyskinesia despite a trial of dose reduction, tapering, or discontinuation of the offending medication</p> <p style="text-align: center;"><b>OR</b></p> <p>2.2 Patient is not a candidate for a trial of dose reduction, tapering, or discontinuation of the offending medication</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Neurologist</li> <li>• Psychiatrist</li> </ul>	

Product Name: Brand Xenazine, generic tetrabenazine	
Diagnosis	Tardive Dyskinesia (Off Label)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

Product Name: Brand Xenazine, generic tetrabenazine

Diagnosis	Tourette's syndrome (off-label)
-----------	---------------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Initial Authorization
---------------	-----------------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Patient has tics associated with Tourette's syndrome

**AND**

2 - History of failure, contraindication, or intolerance to Haldol (haloperidol)

**AND**

3 - Prescribed by or in consultation with one of the following:

- Neurologist
- Psychiatrist

Product Name: Brand Xenazine, generic tetrabenazine

Diagnosis	Tourette's syndrome (off-label)
-----------	---------------------------------

Approval Length	12 month(s)
-----------------	-------------

Therapy Stage	Reauthorization
---------------	-----------------

Guideline Type	Prior Authorization
----------------	---------------------

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Xenleta

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99529
<b>Guideline Name</b>	Xenleta
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Xenleta	
Diagnosis	Community-acquired bacterial pneumonia
Approval Length	7 Day(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - One of the following:  1.1 For continuation of therapy upon hospital discharge	

**OR**

**1.2** As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

**1.3** All of the following:

**1.3.1** Diagnosis of community-acquired bacterial pneumonia (CABP)

**AND**

**1.3.2** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Xenleta

**AND**

**1.3.3** History of failure, contraindication, or intolerance to three of the following antibiotics:

- Amoxicillin
- A macrolide
- Doxycycline
- A fluoroquinolone
- Combination therapy with amoxicillin/clavulanate or cephalosporin AND a macrolide or doxycycline

Product Name: Xenleta\*

Diagnosis

Off-Label Uses

Guideline Type

Prior Authorization

**Approval Criteria**

**1** - One of the following:

**1.1** For continuation of therapy upon hospital discharge

**OR**

**1.2** As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

**1.3** The medication is being prescribed by or in consultation with an infectious disease specialist

Notes

\*Approval Duration: Based on provider recommended treatment durations, not to exceed 6 months

## 2 . Revision History

Date	Notes
5/18/2021	7/1 Implementation

Xenpozyme (olipudase alfa)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-131958
<b>Guideline Name</b>	Xenpozyme (olipudase alfa)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Xenpozyme	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) confirming diagnosis of acid sphingomyelinase deficiency (ASMD)*	

**AND**

**2** - Disease confirmed by ONE of the following: [2]

**2.1** Molecular genetic testing confirms biallelic pathogenic variants in the SMPD1 (sphingomyelin phosphodiesterase-1) gene

**OR**

**2.2** Residual acid sphingomyelinase activity that is less than 10% of controls (in peripheral blood lymphocytes or cultured skin fibroblasts)

**AND**

**3** - Submission of medical records (e.g., chart notes) documenting patient has non-central nervous system manifestations of ASMD

**AND**

**4** - Prescribed by or in consultation with ONE of the following:

- Metabolic disease specialist
- Geneticist

Notes	*Acid Sphingomyelinase Deficiency is also known as Niemann-Pick Disease types A, A/B, and B [1]
-------	---

Product Name: Xenpozyme	
Approval Length	24 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., decrease in spleen size, decrease in liver size, increase in platelet count, improved lung function)

## 2 . Revision History

Date	Notes
8/29/2023	Added new GPI for 4 mg strength

Xermelo

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99658
<b>Guideline Name</b>	Xermelo
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Xermelo	
Diagnosis	Carcinoid Syndrome Diarrhea
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of carcinoid syndrome diarrhea	

**AND**

**2** - Diarrhea is inadequately controlled with somatostatin analog therapy (e.g., octreotide, Sandostatin LAR, Somatuline Depot)

**AND**

**3** - Used in combination with somatostatin analog therapy (e.g., octreotide, Sandostatin LAR, Somatuline Depot)

Product Name: Xermelo	
Diagnosis	Carcinoid Syndrome Diarrhea
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Documentation of positive clinical response to Xermelo	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Xhance (fluticasone nasal spray)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-147691
<b>Guideline Name</b>	Xhance (fluticasone nasal spray)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	6/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Xhance	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Diagnosis of one of the following: <ul style="list-style-type: none"><li>Chronic rhinosinusitis with nasal polyps (CRSwNP)</li><li>Chronic rhinosinusitis without nasal polyps (CRSsNP)</li></ul>	

**AND**

2 - History of failure to generic fluticasone nasal spray

## 2 . Revision History

Date	Notes
5/23/2024	New program

Xolair (omalizumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-146025
<b>Guideline Name</b>	Xolair (omalizumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Xolair	
Diagnosis	Allergic Asthma
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of documentation (e.g., chart notes) confirming diagnosis of moderate to severe persistent allergic asthma	

**AND**

**2** - Submission of documentation (e.g., chart notes, lab values) confirming a positive skin test or in vitro reactivity to a perennial aeroallergen

**AND**

**3** - One of the following:

**3.1** Both of the following:

- Patient is 12 years of age or older
- Submission of documentation (e.g., chart notes, lab values) confirming pre-treatment serum immunoglobulin (Ig)E level between 30 to 700 IU/mL

**OR**

**3.2** Both of the following:

- Patient is 6 years to less than 12 years of age
- Submission of documentation (e.g. chart notes, lab values) confirming pre-treatment serum immunoglobulin (Ig)E level between 30 to 1300 IU/mL

**AND**

**4** - Paid claims or submission of documentation (e.g., chart notes) confirming patient is currently being treated with ONE of the following, unless there is a contraindication or intolerance to these medications:

**4.1** Both of the following:

- High-dose inhaled corticosteroid (ICS) (e.g., greater than 500 mcg fluticasone propionate equivalent/day)
- Additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol ], tiotropium)

**OR**

**4.2** One maximally-dosed combination ICS/LABA product (e.g., Advair [fluticasone propionate/salmeterol], Symbicort [budesonide/formoterol], Breo Ellipta [fluticasone/vilanterol])

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Pulmonologist
- Allergist/immunologist

Product Name: Xolair

Diagnosis	Allergic Asthma
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of documentation (e.g., chart notes) confirming a positive clinical response to therapy (e.g., reduction in exacerbations, improvement in forced expiratory volume in 1 second [FEV1], decreased use of rescue medications)

**AND**

**2** - Paid claims or submission of documentation (e.g., chart notes) confirming patient continues to be treated with an inhaled corticosteroid (ICS) (e.g., fluticasone, budesonide) with or without additional asthma controller medication (e.g., leukotriene receptor antagonist [e.g., montelukast], long-acting beta-2 agonist [LABA] [e.g., salmeterol], tiotropium) unless there is a contraindication or intolerance to these medications

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Pulmonologist

- Allergist/immunologist

Product Name:Xolair	
Diagnosis	Chronic Spontaneous Urticaria (CSU)
Approval Length	3 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of documentation (e.g., chart notes) confirming diagnosis of chronic spontaneous urticaria</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Persistent symptoms (itching and hives) for at least 4 consecutive weeks despite titrating to an optimal dose with a second generation H1 antihistamine (e.g., cetirizine, fexofenadine), unless there is a contraindication or intolerance to H1 antihistamines</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Paid claims or submission of documentation (e.g., chart notes) confirming concurrent use with an H1 antihistamine, unless there is a contraindication or intolerance to H1 antihistamines</p> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Paid claims or submission of documentation (e.g., chart notes) confirming patient has tried and had an inadequate response or intolerance to at least TWO of the following additional therapies:</p> <ul style="list-style-type: none"> <li>• Doxepin</li> <li>• H1 antihistamine</li> <li>• H2 antagonist (e.g., famotidine, cimetidine)</li> <li>• Hydroxyzine</li> </ul>	

- Leukotriene receptor antagonist (e.g., montelukast)

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Allergist/immunologist
- Dermatologist

Product Name:Xolair	
Diagnosis	Chronic Spontaneous Urticaria (CSU)
Approval Length	6 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Patient's disease status has been re-evaluated since the last authorization to confirm the patient's condition warrants continued treatment</p> <p><b>AND</b></p> <p><b>2</b> - Submission of documentation (e.g., chart notes) confirming patient has experienced at least one of the following:</p> <ul style="list-style-type: none"> <li>• Reduction in itching severity from baseline</li> <li>• Reduction in the number of hives from baseline</li> </ul>	

Product Name:Xolair	
Diagnosis	Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

## **Approval Criteria**

1 - Patient is 18 years of age or older

**AND**

2 - Submission of documentation (e.g., chart notes) confirming ONE of the following:

2.1 ALL of the following:

2.1.1 Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) defined by ALL of the following:

2.1.1.1 TWO or more of the following symptoms for greater than or equal to 12 weeks duration:

- Mucopurulent discharge
- Nasal obstruction and congestion
- Decreased or absent sense of smell
- Facial pressure or pain

**AND**

2.1.1.2 ONE of the following:

- Evidence of inflammation on paranasal sinus examination or computed tomography (CT)
- Evidence of purulence coming from paranasal sinuses or ostiomeatal complex

**AND**

2.1.1.3 The presence of nasal polyps

**AND**

2.1.2 ONE of the following:

- Patient has required prior sino-nasal surgery
- Patient has required systemic corticosteroids in the previous 2 years

**AND**

**2.1.3** Patient has been unable to obtain symptom relief after trial of ALL of the following agents/classes of agents:

- Nasal saline irrigations
- Intranasal corticosteroids (e.g. fluticasone, mometasone, triamcinolone, etc.)
- Antileukotriene agents (e.g. montelukast, zafirlukast, zileuton)

**OR**

**2.2** ALL of the following:

**2.2.1** Diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP)

**AND**

**2.2.2** Patient is currently on Xolair therapy

**AND**

**3** - Patient will receive Xolair as add-on maintenance therapy in combination with intranasal corticosteroids

**AND**

**4** - Patient is NOT receiving Xolair in combination with another biologic medication [e.g., Dupixent (dupilumab), Nucala (mepolizumab)]

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Otolaryngologist
- Allergist
- Pulmonologist

Product Name:Xolair	
Diagnosis	Chronic Rhinosinusitis with Nasal Polyps (CRSwNP)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of documentation (e.g., chart notes, lab values) confirming a positive clinical response to Xolair therapy</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient will continue to receive Xolair as add-on maintenance therapy in combination with intranasal corticosteroids</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Patient is NOT receiving Xolair in combination with another biologic medication [e.g., Dupixent (dupilumab), Nucala (mepolizumab)]</p> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with one of the following:</p> <ul style="list-style-type: none"> <li>• Otolaryngologist</li> <li>• Allergist</li> <li>• Pulmonologist</li> </ul>	

Product Name:Xolair	
Diagnosis	IgE-Mediated Food Allergy
Approval Length	20 Week(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - One of the following:

1.1 Submission of documentation (e.g., chart notes, lab values) confirming both of the following:

1.1.1 Diagnosis of IgE Mediated Food Allergy as evidenced by one of the following:

- Positive skin prick test (defined as greater than or equal to 4 mm wheal greater than saline control) to food
- Positive food specific IgE (greater than or equal to 6 kU/L)
- Positive oral food challenge, defined as experiencing dose-limiting symptoms at a single dose of less than or equal to 300 mg of food protein

**AND**

1.1.2 Clinical history of IgE Mediated Food Allergy

**OR**

1.2 Submission of documentation (e.g., chart notes, lab values) confirming patient has a history of severe allergic response, including anaphylaxis, following exposure to one or more foods

**AND**

2 - Patient is 1 year of age or older

**AND**

**3** - Used in conjunction with food allergen avoidance

**AND**

**4** - Submission of documentation (e.g., chart notes, lab values) confirming both of the following:

- Baseline (pre-Xolair treatment) serum total IgE level is greater than or equal to 30 IU/mL and less than or equal to 1850 IU/mL
- Dosing is according to serum total IgE levels and body weight

**AND**

**5** - Prescribed by or in consultation with one of the following:

- Allergist
- Immunologist

Product Name:Xolair

Diagnosis	IgE-Mediated Food Allergy
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

**1** - Submission of documentation (e.g., chart notes, lab values) confirming a positive clinical response to therapy e.g., reduction of type 1 allergic reactions, including anaphylaxis, following accidental exposure to one or more foods)

**AND**

**2** - Used in conjunction with food allergen avoidance

**AND**

**3** - Submission of documentation (e.g., chart notes, lab values) confirming that dosing will continue to be based on body weight and pretreatment total IgE serum levels (Note: Dose should only be adjusted during therapy due to significant changes in patient body weight)

**AND**

**4** - Prescribed by or in consultation with one of the following:

- Allergist
- Immunologist

## 2 . Background

### Clinical Practice Guidelines

**The Global Initiative for Asthma Global Strategy for Asthma Management and Prevention: Table 1. Low, medium and high daily doses of inhaled corticosteroids in adolescents and adults 12 years and older [3]**

Inhaled corticosteroid	Total Daily ICS Dose (mcg)		
	Low	Medium	High
Beclometasone dipropionate (pMDI, standard particle, HFA)	200-500	> 500-1000	> 1000
Beclometasone dipropionate (DPI or pMDI, extrafine particle*, HFA)	100-200	> 200-400	> 400
Budesonide (DPI, or pMDI, standard particle, HFA)	200-400	> 400-800	> 800
Ciclesonide (pMDI, extrafine particle*, HFA)	80-160	> 160-320	> 320
Fluticasone furoate (DPI)	100		200

Fluticasone propionate (DPI)	100-250	> 250-500	> 500
Fluticasone propionate (pMDI, standard particle, HFA)	100-250	> 250-500	> 500
Mometasone furoate (DPI)	Depends on DPI device – see product information		
Mometasone furoate (pMDI, standard particle, HFA)	200-400		> 400
<p>DPI: dry powder inhaler; HFA: hydrofluoroalkane propellant; ICS: inhaled corticosteroid; N/A: not applicable; pMDI: pressurized metered dose inhaler (non-chlorofluorocarbon formulations); ICS by pMDI should be preferably used with a spacer *See product information.</p> <p><b><i>This is not a table of equivalence</i></b>, but instead, suggested total daily doses for the ‘low’, ‘medium’ and ‘high’ dose ICS options for adults/adolescents, based on available studies and product information. Data on comparative potency are not readily available and therefore this table does NOT imply potency equivalence. Doses may be country -specific depending on local availability, regulatory labelling and clinical guidelines.</p> <p>For new preparations, including generic ICS, the manufacturer’s information should be reviewed carefully; products containing the same molecule may not be clinically equivalent.</p>			

### 3 . Revision History

Date	Notes
4/23/2024	New (updated) UM for Xolair, updated criteria for all approved indications. Added new GPs for SC formulations.

Xolremdi (mavorixafor)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-150126
<b>Guideline Name</b>	Xolremdi (mavorixafor)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Xolremdi	
Diagnosis	WHIM syndrome
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Submission of medical records (e.g., chart notes) documenting all of the following:  1.1 Diagnosis of WHIM (warts, hypogammaglobulinemia, infections and myelokathexis) syndrome	

**AND**

**1.2** Patient has genotype confirmed variant of CXCR4 as detected by an FDA-approved test or a test performed at a facility approved by Clinical Laboratory Improvement Amendments (CLIA)

**AND**

**1.3** Patient has an absolute neutrophil count (ANC) less than or equal to 500 cells / $\mu$ L

**AND**

**2** - Patient is 12 years of age or older

**AND**

**3** - Prescribed by or in consultation with one of the following:

- Immunologist
- Hematologist
- Geneticist
- Allergist

Product Name: Xolremdi	
Diagnosis	WHIM syndrome
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy (e.g., improvement in ANC, reduction in infections)

## 2 . Revision History

Date	Notes
7/23/2024	New program

Xopenex Respules

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99502
<b>Guideline Name</b>	Xopenex Respules
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Brand Xopenex inhalation soln, generic levalbuterol inhalation soln	
Approval Length	12 month(s)
Guideline Type	Step Therapy
<b>Approval Criteria</b>  1 - The patient has a history of failure, contraindication, or intolerance to treatment with albuterol inhalation solution	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Xphozah (tenapanor)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-139344
<b>Guideline Name</b>	Xphozah (tenapanor)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Xphozah	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of hyperphosphatemia in chronic kidney disease  <b>AND</b>	

2 - Patient is on dialysis

**AND**

3 - Submission of medical records (e.g., chart notes) or paid claims confirming trial and inadequate response (minimum 30-day supply), contraindication or intolerance to ALL of the following:

- calcium carbonate
- calcium acetate
- sevelamer carbonate

Product Name:Xphozah	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	
<b>AND</b>	
2 - Trial and inadequate response (minimum 30-day supply), contraindication or intolerance to ALL of the following::	
<ul style="list-style-type: none"><li>• calcium carbonate</li><li>• calcium acetate</li><li>• sevelamer carbonate</li></ul>	

## 2 . Revision History

Date	Notes
------	-------

1/23/2024	New program
-----------	-------------

Xromi (hydroxyurea)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-241268
<b>Guideline Name</b>	Xromi (hydroxyurea)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Xromi	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of sickle cell anemia  <b>AND</b>	

2 - Patient has moderate to severe painful crises

**AND**

3 - Patient is 6 months of age or older

**AND**

4 - One of the following:

4.1 Patient is unable to swallow solid oral dosage forms (e.g., oral tablet, capsule) due to one of the following:

- Age
- Physical impairment (e.g., difficulties with motor or oral coordination)
- Dysphagia
- Patient is using a feeding tube or nasal gastric tube

**OR**

4.2 Dosage requirements cannot be met with a solid dosage form

Notes	Note: PA is required for patients greater than 10 years of age
-------	--

Product Name: Xromi	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy	
Notes	Note: PA is required for patients greater than 10 years of age

## 2 . Revision History

Date	Notes
4/24/2025	New program

Xuriden

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99660
<b>Guideline Name</b>	Xuriden
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

### 1 . Criteria

Product Name:Xuriden	
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of a hereditary orotic aciduria	

Product Name:Xuriden
----------------------

Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Documentation of positive clinical response to Xuriden therapy</p>	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Medicaid SP to Medicaid Arizona SP for 7/1

Yorvipath (palopegteriparatide)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-161661
<b>Guideline Name</b>	Yorvipath (palopegteriparatide)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	1/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Yorvipath	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting all of the following: <b>1.1</b> Diagnosis of hypoparathyroidism	

**AND**

**1.2** Requested drug is not being used in the setting of acute post-surgical hypoparathyroidism

**AND**

**1.3** Patient has achieved albumin-corrected serum calcium of at least 7.8 mg/dL using calcium and active vitamin D (e.g., calcitriol) treatment

**AND**

**2** - Prescribed by or in consultation with an endocrinologist

Product Name:Yorvipath	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting that patient demonstrates positive clinical response to therapy as evidenced by maintenance of normalized calcium levels compared to baseline	

## 2 . Revision History

Date	Notes
12/4/2024	New program

Zeposia (ozanimod)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-329259
<b>Guideline Name</b>	Zeposia (ozanimod)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	8/1/2025
-----------------	----------

## 1 . Criteria

Product Name:Zeposia	
Diagnosis	Multiple Sclerosis (MS)
Approval Length	12 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Diagnosis of multiple sclerosis (MS)	

**AND**

**2** - Submission of medical records (e.g., chart notes) or verification of paid claims confirming a history of failure, contraindication, or intolerance to a trial of at least TWO of the preferred alternatives:

- Avonex
- Brand Copaxone
- generic dalfampridine
- generic dimethyl fumarate
- generic fingolimod
- Kesimpta
- Ocrevus/Ocrevus Zunovo
- Rebif
- generic teriflunomide
- Tysabri

Product Name:Zeposia	
Diagnosis	Multiple Sclerosis (MS)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
1 - Patient demonstrates positive clinical response to therapy (e.g., stability in radiologic disease activity, clinical relapses, disease progression)	

Product Name:Zeposia	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 Week(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Diagnosis of moderately to severely active ulcerative colitis (UC)

**AND**

2 - Submission of medical records (e.g., chart notes) or verification of paid claims documenting BOTH of the following:

2.1 Trial and failure, contraindication, or intolerance to ONE of the following conventional therapies:

- 6-mercaptopurine
- Aminosalicylate (e.g., mesalamine, olsalazine, sulfasalazine)
- Azathioprine
- Corticosteroids (e.g., prednisone)

**AND**

2.2 History of failure, contraindication, or intolerance to ALL of the following:

- A preferred adalimumab biosimilar
- infliximab
- Xeljanz (tofacitinib) oral tablet (IR or XR)
- A preferred ustekinumab biosimilar

**AND**

3 - Prescribed by or in consultation with a gastroenterologist

Product Name:Zeposia	
Diagnosis	Ulcerative Colitis (UC)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy

**AND**

2 - Prescribed by or in consultation with a gastroenterologist

**2 . Revision History**

Date	Notes
7/16/2025	Updated preferred agents/embedded steps, updated criteria throughout. MS: Added Ocrevus Zunovo as prerequisite option.

Zimhi (naloxone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114472
<b>Guideline Name</b>	Zimhi (naloxone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Zimhi	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - History of failure, or intolerance to preferred naloxone products (e.g., Brand Narcan nasal spray, Kloxxado, preferred naloxone injections)	

## 2 . Revision History

Date	Notes
9/26/2022	New program

Zinplava (bezlotoxumab)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-133807
<b>Guideline Name</b>	Zinplava (bezlotoxumab)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2023
-----------------	-----------

## 1 . Criteria

Product Name:Zinplava	
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting ALL of the following:</p> <p>1.1 Used for the reduction of the recurrence of Clostridium difficile infection (CDI)</p> <p style="text-align: center;"><b>AND</b></p>	

**1.2** Patient is 1 year of age or older

**AND**

**1.3** Used in combination with antibacterial drug treatment for CDI [e.g., oral Vancocin (vancomycin), Flagyl (metronidazole), or Dificid (fidaxomicin)]

**AND**

**1.4** Patient has one or more of the following risk factors associated with CDI recurrence:

- One or more prior episodes of CDI in the previous 6 months
- Immunocompromised
- Chronic dialysis
- Inflammatory bowel disease
- Continued use of non-CDI antimicrobials after diagnosis of CDI and/or after CDI treatment

**AND**

**2** - Prescribed by or in consultation with one of the following:

- Infectious disease specialist
- Gastroenterologist

## **2 . Revision History**

Date	Notes
9/26/2023	New Program

Zolgensma (onasemnogene abeparvovec-xioi)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-124879
<b>Guideline Name</b>	Zolgensma (onasemnogene abeparvovec-xioi)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2023
-----------------	----------

## 1 . Criteria

Product Name:Zolgensma	
Approval Length	1 Time Authorization in Lifetime
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - The mutation or deletion of genes in chromosome 5q resulting in one of the following: [1-8, A]  1.1 Homozygous gene deletion or mutation of SMN1 gene (e.g., homozygous deletion of exon 7 at locus 5q13)	

**OR**

**1.2** Compound heterozygous mutation of SMN1 gene (e.g., deletion of Survival of Motor Neuron 1 [SMN1] exon 7 [allele 1] and mutation of SMN1 [allele 2])

**AND**

**2** - One of the following:

**2.1** Both of the following: [1-5]

**2.1.1** Diagnosis of diagnosis of SMA Type 0, I or Type II spinal muscular atrophy (SMA) confirmed by a neurologist with expertise in the treatment of SMA

**AND**

**2.1.2** Patient is less than or equal to 2 years of age

**OR**

**2.2** Both of the following:

**2.2.1** Diagnosis of SMA based on the results of SMA newborn screening

**AND**

**2.2.2** Patient has 3 copies or less of Survival of Motor Neuron 2 (SMN 2)

**AND**

**3** - Patient is not dependent on either of the following:

- Invasive ventilation or tracheostomy
- Use of invasive ventilation beyond use of naps and nighttime sleep

**AND**

**4** - Submission of medical records (e.g., chart notes, laboratory values) documenting patient's anti-AAV9 antibody titers are less than or equal to 1:50 [1]

**AND**

**5** - Patient is not to receive concomitant SMN modifying therapy (e.g. Spinraza)

**AND**

**6** - Prescribed by a neurologist with expertise in the diagnosis of SMA

**AND**

**7** - Patient has never received Zolgensma treatment in their lifetime

## 2 . Revision History

Date	Notes
4/20/2023	Added new GPIs, no changes to criteria.

Zontivity

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99503
<b>Guideline Name</b>	Zontivity
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Zontivity	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - ONE of the following: <ul style="list-style-type: none"><li>History of myocardial infarction (MI)</li><li>Peripheral arterial disease (PAD)</li></ul>	

**AND**

**2** - Patient does not have a history of ONE of the following:

- Stroke
- Transient ischemic attack (TIA)
- Intracranial hemorrhage (ICH)

**AND**

**3** - Patient does not have active pathological bleeding

## **2 . Revision History**

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Zortress - ARIZONA

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99504
<b>Guideline Name</b>	Zortress - ARIZONA
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
-----------------	-----------

## 1 . Criteria

Product Name:Zortress	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - Kidney transplant rejection prophylaxis in patients at low-moderate immunologic risk  <b>OR</b>  2 - Liver transplant rejection prophylaxis	

## 2 . Revision History

Date	Notes
3/11/2021	Bulk Copy C&S Arizona Standard to Medicaid Arizona Standard for 7 /1 go live

Zoryve (roflumilast)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-154427
<b>Guideline Name</b>	Zoryve (roflumilast)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	9/5/2024
-----------------	----------

## 1 . Criteria

Product Name:Zoryve 0.3% cream	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of plaque psoriasis	

**AND**

**2** - Patient is 6 years of age or older

**AND**

**3** - Submission of medical records (e.g., chart notes) or paid claims history documenting a minimum duration of a 4 week trial and failure, contraindication, or intolerance to TWO of the following topical therapies (trial must be from two different classes):

- Corticosteroids (e.g., betamethasone, clobetasol)
- Vitamin D analogs (e.g., calcitriol, calcipotriene)
- Tazarotene
- Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)
- Anthralin
- Coal tar

**AND**

**4** - Prescribed by or in consultation with a dermatologist

Product Name: Zoryve 0.3% cream	
Diagnosis	Plaque Psoriasis (PsO)
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	
<b>1</b> - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by one of the following:	
<ul style="list-style-type: none"><li>• Reduction in the body surface area (BSA) involvement from baseline</li><li>• Improvement in symptoms (e.g., pruritus, inflammation) from baseline</li></ul>	

Product Name:Zoryve 0.15% cream	
Diagnosis	Atopic Dermatitis
Approval Length	6 month(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p><b>1</b> - Submission of medical records (e.g., chart notes) documenting a diagnosis of mild to moderate atopic dermatitis</p> <p style="text-align: center;"><b>AND</b></p> <p><b>2</b> - Patient is 6 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p><b>3</b> - Submission of medical records (e.g., chart notes) or paid claims history documenting trial and failure of a minimum 30-day supply (14-day supply for topical corticosteroids), contraindication, or intolerance to BOTH of the following</p> <ul style="list-style-type: none"> <li>• ONE topical corticosteroid [e.g., mometasone furoate, fluocinolone acetonide (generic Synalar), fluocinonide]</li> <li>• ONE topical calcineurin inhibitor [e.g., pimecrolimus (generic Elidel), tacrolimus (generic Protopic)]</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p><b>4</b> - Prescribed by or in consultation with a dermatologist</p>	

Product Name:Zoryve foam	
Diagnosis	Seborrheic Dermatitis
Approval Length	6 month(s)
Therapy Stage	Initial Authorization

Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) documenting a diagnosis of seborrheic dermatitis</p> <p style="text-align: center;"><b>AND</b></p> <p>2 - Patient is 9 years of age or older</p> <p style="text-align: center;"><b>AND</b></p> <p>3 - Submission of medical records (e.g., chart notes) or paid claims history documenting a minimum duration of a 4 week trial and failure, contraindication, or intolerance to TWO of the following topical therapies (trial must be from two different classes):</p> <ul style="list-style-type: none"> <li>• Corticosteroids (e.g., betamethasone, clobetasol)</li> <li>• Antifungals (e.g., ciclopirox, ketoconazole)</li> <li>• Calcineurin inhibitors (e.g., tacrolimus, pimecrolimus)</li> </ul> <p style="text-align: center;"><b>AND</b></p> <p>4 - Prescribed by or in consultation with a dermatologist</p>	

Product Name: Zoryve foam	
Diagnosis	Seborrheic Dermatitis
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p>	

1 - Submission of medical records (e.g., chart notes) documenting positive clinical response to therapy as evidenced by improvement from baseline for one of the following:

- Scaling
- Erythema
- Pruritis
- Body surface area (BSA) involvement

## 2 . Revision History

Date	Notes
9/5/2024	Removed "initial auth" from AD indication section

Ztalmy (ganaxolone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-114155
<b>Guideline Name</b>	Ztalmy (ganaxolone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	10/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Ztalmy	
Approval Length	6 month(s)
Therapy Stage	Initial Authorization
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Submission of documentation (e.g., chart notes) confirming diagnosis of cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD)	

**AND**

**2** - Patient has a mutation in the CDKL5 gene

**AND**

**3** - Patient is 2 years of age or older

**AND**

**4** - Patient is experiencing motor seizures (e.g., bilateral tonic, generalized tonic-clonic, bilateral clonic, atonic, focal, or bilateral tonic-clonic)

**AND**

**5** - One of the following:

**5.1** Trial and failure, contraindication, or intolerance to two preferred anticonvulsants (e.g., valproic acid, levetiracetam, lamotrigine)

**OR**

**5.2** For continuation of prior therapy

**AND**

**6** - Prescribed by or in consultation with a neurologist

Product Name:Ztalmy	
Approval Length	12 month(s)
Therapy Stage	Reauthorization
Guideline Type	Prior Authorization

**Approval Criteria**

1 - Documentation of positive clinical response to therapy as evidenced by a reduction in the frequency of seizures from baseline

**2 . Revision History**

Date	Notes
9/20/2022	New Program

Zunveyl (benzgalantamine)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-241267
<b>Guideline Name</b>	Zunveyl (benzgalantamine)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	5/1/2025
-----------------	----------

### 1 . Criteria

Product Name:Zunveyl	
Approval Length	12 month(s)
Guideline Type	Prior Authorization
<b>Approval Criteria</b> 1 - Diagnosis of mild to moderate dementia of the Alzheimer's type  <b>AND</b>  2 - Patient is 18 years of age or older	

**AND**

**3** - Trial and failure (of a minimum 30-day supply), contraindication, or intolerance to ALL of the following:

- generic galantamine
- generic rivastigmine
- generic donepezil

## **2 . Revision History**

Date	Notes
4/24/2025	New program

Zurzuvae (zuranolone)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-139356
<b>Guideline Name</b>	Zurzuvae (zuranolone)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	2/1/2024
-----------------	----------

## 1 . Criteria

Product Name:Zurzuvae	
Approval Length	14 Day(s)
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - One of the following:</p> <p>1.1 Diagnosis of severe postpartum depression (PPD)</p> <p style="text-align: center;"><b>OR</b></p>	

**1.2** Both of the following:

**1.2.1** Diagnosis of mild to moderate postpartum depression (PPD)

**AND**

**1.2.2** Trial and failure, contraindication or intolerance to at least one oral SSRI or SNRI (e.g., escitalopram, duloxetine)

**AND**

**2** - Patient is 18 years of age or older

**AND**

**3** - Onset of symptoms in the third trimester or within 4 weeks of delivery

**AND**

**4** - Prescriber attests that the patient has been counseled and has agreed to adhere to the following: Will follow instructions to not drive or operate machinery until at least 12 hours after taking each dose of Zurzuvae for the duration of the 14-day treatment course and that patients are informed that they may not be able to assess their own driving competence or the degree of driving impairment caused by Zurzuvae

## **2 . Revision History**

Date	Notes
1/23/2024	New program

Zynteglo (betibeglogene autotemcel)

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-116186
<b>Guideline Name</b>	Zynteglo (betibeglogene autotemcel)
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	11/1/2022
-----------------	-----------

## 1 . Criteria

Product Name:Zynteglo	
Approval Length	1 Time Authorization in Lifetime
Guideline Type	Prior Authorization
<p><b>Approval Criteria</b></p> <p>1 - Submission of medical records (e.g., chart notes) confirming diagnosis of transfusion-dependent beta-thalassemia as confirmed by the presence of a mutation at both alleles of the <math>\beta</math>-globin gene (i.e., <math>\beta^0/\beta^0</math>, <math>\beta^0/\beta^+</math>, <math>\beta^+/\beta^+</math>, <math>\beta^0/\beta^E</math>)</p> <p style="text-align: center;"><b>AND</b></p>	

**2** - One of the following:

- Patient has a history of transfusions of at least 100 mL/kg/year of packed red blood cells (pRBCs)
- Patient requires 8 or more red blood cell (RBC) transfusions per year

**AND**

**3** - Patient is 4 years of age or older [A]

**AND**

**4** - Patient is ineligible for an allogeneic hematopoietic stem cell transplant with an HLA-identical sibling donor [B]

**AND**

**5** - Provider attests that patient is clinically stable and eligible to undergo hematopoietic stem cell transplant (HSCT) and has not received any prior gene therapy or HSCT

**AND**

**6** - Patient has obtained a negative test result for all of the following prior to cell collection:

- Hepatitis B virus (HBV)
- Hepatitis C virus (HCV)
- Human T-lymphotrophic virus 1 & 2 (HTLV-1/HTLV-2)
- Human immunodeficiency virus (HIV)

**AND**

**7** - Patient is able to provide an adequate number of cells to meet the minimum recommended dose of  $5 \times 10^6$  CD34+ cells/kg

**AND**

**8** - Patient does not have any of the following [1-4]:

- Severely elevated iron in the heart (e.g., patients with cardiac T2\* less than 10 msec by MRI)
- Advanced liver disease
- MRI results of the liver demonstrating liver iron content greater than or equal to 15 mg/g (unless biopsy confirms absence of advanced disease)

**AND**

**9** - Both of the following:

- Iron chelation therapy (e.g., deferoxamine, deferasirox) will be discontinued for at least 7 days prior to initiating myeloablative conditioning therapy
- Prophylactic HIV anti-retroviral medications (e.g., Truvada, Descovy) or hydroxyurea will be discontinued for at least one month prior to mobilization (or for the expected duration for elimination of those medications)

**AND**

**10** - Prescribed by a stem cell transplant specialist

**AND**

**11** - Patient has never received Zynteglo treatment in their lifetime

## **2 . Endnotes**

- A. The safety and efficacy of Zynteglo in children less than 4 years of age have not been established. [1]
- B. Per consultant feedback, Zynteglo should be reserved for patients who do not have an HLA-identical sibling for an allogeneic hematopoietic stem cell transplant. [5]

## **3 . Revision History**

Date	Notes
10/28/2022	New Program

Zyvox

--

## Prior Authorization Guideline

<b>Guideline ID</b>	GL-99578
<b>Guideline Name</b>	Zyvox
<b>Formulary</b>	<ul style="list-style-type: none"><li>Medicaid - Arizona (AZM, AZMREF, AZMDDD)</li></ul>

### Guideline Note:

Effective Date:	12/9/2021
P&T Approval Date:	
P&T Revision Date:	

## 1 . Criteria

Product Name: Brand Zyvox*, generic linezolid*	
Diagnosis	Labeled Uses
Guideline Type	Prior Authorization
<b>Approval Criteria</b>  1 - One of the following:  1.1 For continuation of therapy upon hospital discharge	

**OR**

**1.2** As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

**1.3** BOTH of the following:

**1.3.1** ONE of the following diagnoses:

- Nosocomial pneumonia
- Community-acquired pneumonia
- Skin and skin structure infections (complicated and uncomplicated)

**AND**

**1.3.2** Infection caused by an organism that is confirmed to be or likely to be susceptible to treatment with Zyvox

**OR**

**1.4** Invasive infection caused by or likely to be caused by vancomycin-resistant *Enterococcus faecium* (VRE)

Notes	*Approval Duration: For vancomycin-resistant <i>Enterococcus faecium</i> , authorization will be issued for 28 days. For osteomyelitis, authorization will be issued for the requested duration, not to exceed 6 weeks. All other approvals will be issued for 14 days.
-------	---

Product Name: Brand Zyvox*, generic linezolid*	
Diagnosis	Off label Uses
Guideline Type	Prior Authorization
<b>Approval Criteria</b>	

1 - For continuation of therapy upon hospital discharge

**OR**

2 - As continuation of therapy when transitioning from intravenous antibiotics that are shown to be sensitive to the cultured organism for the requested indication

**OR**

3 - The medication is being prescribed by or in consultation with an Infectious Disease specialist

Notes

\*Approval Duration: Based on provider recommended treatment durations, not to exceed 6 months.

## 2 . Revision History

Date	Notes
7/21/2021	Update guideline